

THE UNIVERSITY OF LINCOLN

**The behaviour and ecology of adult common bream *Abramis brama* (L.) in  
a heavily modified lowland river**

A Thesis submitted for the Degree of Doctor of Philosophy

By

Christopher John Gardner B.Sc. (Hons)

September 2013

## CONTENTS

<b>Publications and outputs</b>	<b>5</b>
<b>List of tables</b>	<b>6</b>
<b>List of figures</b>	<b>7</b>
<b>Acknowledgements</b>	<b>12</b>
<b>Abstract</b>	<b>13</b>
<b>1. Rivers, fish and human interventions</b>	<b>15</b>
1.1 General introduction	15
1.2 Objectives	17
1.3 A ‘natural’ lowland riverine ecosystem	17
1.4 Anthropogenic impacts on lowland riverine ecosystems	18
1.5 Fish ecology and rivers	21
1.6 Common bream ecology	23
1.7 Studying spatio-temporal behaviour of fishes with telemetry techniques	29
1.8 Proposed study	33
<b>2. The ecology of the lower River Witham</b>	<b>35</b>
2.1 Objectives	35
2.2 The lower River Witham	35
2.2.1 Background	35
2.2.2 History	36
2.2.3 Catchment and land use	39
2.2.4 Water quality	39
2.2.5 Biological quality	40
2.2.6 Flow regime and flood events	41
2.2.7 Water level management	43
2.2.8 Trent-Witham-Ancholme water transfer scheme	43
2.2.9 Morphology	43
2.2.10 Habitat assessments	45
2.2.11 Barriers to fish migration	49
2.2.12 River uses	49
2.2.13 Conservation	49
2.2.14 Fishery assessments	50
2.2.15 Water Framework Directive classification	57
2.3 Discussion	57
2.3.1 Water quality	57
2.3.2 Biological quality	58
2.3.3 Habitat assessments	58
2.3.4 Fishery assessments	58
<b>3. The effects of surgical implantation of a telemetry transmitter into the peritoneal cavity and translocation on the temporal-spatial behaviour of common bream</b>	<b>61</b>
3.1 Introduction	61

3.2	Material and methods	62
3.2.1	Study area	62
3.2.2	Sampling procedures	62
3.2.3	Tag implantation protocol	69
3.2.4	Tracking	76
3.2.5	Recapture and translocation	76
3.2.6	Data handling and statistical analysis.	78
3.3	Results	79
3.3.1	Recaptured fish	80
3.3.2	Effects of tagging procedure	82
3.3.3	Effects of translocation	83
3.4	Discussion	86
3.4.1	Tagging protocol	86
3.4.2	Recaptured fish	88
3.4.3	Shoaling behaviour	89
3.4.4	Fish that left the study area	89
3.4.5	Effects of tagging procedure	90
3.4.6	Effects of translocation	90
3.4.7	Possible experimental bias	91
<b>4.</b>	<b>Seasonal movements with shifts in lateral and longitudinal habitat use</b>	<b>93</b>
4.1	Introduction	93
4.2	Material and methods	93
4.2.1	Study area	93
4.2.2	Sampling and tagging procedures	95
4.2.3	Environmental data	96
4.2.4	Data handling	96
4.2.5	Statistical analysis	97
4.3	Results	97
4.3.1	Monthly distance and monthly longitudinal distribution	97
4.3.2	Use of lateral habitat	100
4.4	Discussion	101
4.4.1	Monthly distance and monthly longitudinal distribution	101
4.4.2	Use of lateral habitat	106
<b>5.</b>	<b>Seasonal water level manipulations for flood risk management influences home-range size</b>	<b>109</b>
5.1	Introduction	109
5.2	Material and methods	110
5.2.1	Study area	110
5.2.2	Sampling and tagging procedures	111
5.2.3	Data handling and statistical analysis	111
5.3	Results	114
5.3.1	Effect of artificial water level manipulations	114
5.4	Discussion	118
5.4.1	Home-range size	118
5.4.2	Effect of artificial water level manipulations	118
<b>6.</b>	<b>Social networks in common bream</b>	<b>121</b>
6.1	Introduction	121
6.2	Material and methods	125

6.3	Data handling and statistical analysis	125
6.4	Results	127
6.5	Discussion	132
<b>7.</b>	<b>Conclusions, management implications and further research</b>	<b>134</b>
7.1	Rehabilitation of lowland rivers	134
7.2	Conclusions	135
7.2.1	Review of rivers, fish, human interventions, telemetry and common bream	135
7.2.2	The ecology of the lower River Witham	135
7.2.3	Effects of tagging and translocation	136
7.2.4	Seasonal movements and shifts in habitat use	136
7.2.5	Home-range size and the effects of artificial water level manipulations	138
7.2.6	Social networks	139
7.3	Recommendations for further research	139
	<b>References</b>	<b>141</b>



## **Publications and outputs**

### **Scientific Papers**

Gardner, C.J., Deeming, D.C. & Eady, P.E. (2013) Seasonal movements with shifts in lateral and longitudinal habitat use by common bream *Abramis brama* (L.) in a heavily modified lowland river. *Fisheries Management and Ecology* **20**, 315-325. doi: 10.1111/fme.12014

Gardner, C.J., Deeming, D.C. & Eady, P.E. (2013) Seasonal water level manipulation for flood risk management influences home-range size of common bream *Abramis brama* L. in a lowland river. *Rivers Research and Applications*, doi: 10.1002/rra.2727

Gardner, C.J., Deeming, D.C., Eady, P.E. & Wellby, I. (in prep) The effects of surgical implantation of a telemetry transmitter into the peritoneal cavity and translocation on the spatio-temporal behaviour of common bream. *Fisheries Research*,

### **Conference proceedings**

Gardner, C.J. (2009) Using acoustic telemetry to target improvements to the lower River Witham, Lincolnshire. *Proceedings of the 40<sup>th</sup> Institute of Fisheries Management Conference* (eds: Dolben, I. & Axford, S.) 99-104 pp.

### **Presentations**

Gardner, C.J. (2006) Bream Tracking in the River Witham. Environment Agency Anglian Region Fisheries Science Symposium.

Gardner, C.J. (2007) Bream Tracking in the River Witham. Environment Agency Anglian Region Fisheries Science Symposium.

Gardner, C.J. (2008) Bream Tracking in the River Witham. Grantham Angling Association Annual General Meeting.

Gardner, C.J. (2008) Bream Tracking in the River Witham. Environment Agency National Fisheries Science Symposium.

Gardner, C.J. (2008) Bream Tracking in the River Witham. Post-graduate showcase, Lincoln University.

Gardner, C.J. (2009) Lower River Witham Water Level Management; Attempting Regime Change. Environment Agency Fisheries Science Symposium.

Gardner, C.J. (2009) Bream Tracking in the River Witham. Lincoln University.

Gardner, C.J. (2009) Using Acoustic Telemetry to Target Improvements to the Lower River Witham, Lincolnshire. Institute of Fisheries Management Conference.

Gardner, C.J. (2010) Using Acoustic Telemetry to Target Improvements to the Lower River Witham, Lincolnshire. Environment Agency area management team.

### **Conference posters**

Gardner, C.J. (2009) Bream Tracking in the River Witham. Institute of Fisheries Management Conference.

### **Press articles**

Gardner, C.J. (2009) Bream Tracking in the River Witham. 'Carp and Coarse Angler' Magazine, Mpress Ltd. Romford.

Gardner, C. J. (2010) Bream Tracking in the River Witham. Institute of Fisheries Management 'Fish' Magazine, IFM Nottingham.

<b>List of tables</b>	<b>Page</b>
Table 1.1	Fishes of modified European floodplain rivers have been classified into guilds by their habitat requirements. Simplified from Schiemer & Waidbacher (1992) and Nunn <i>et al.</i> (2007).
	22
Table 2.1	Results from linear regression analysis of Environment Agency water quality data.
	39
Table 2.2	Results from linear regression analysis of macroinvertebrate indices (Environment Agency data) against date.
	41
Table 2.3	Results from River Habitat Surveys Surveys (Environment Agency data).
	46
Table 2.4	Summary of hydro-acoustic survey results.
	54
Table 2.5	Water Framework Directive classifications for the lower River Witham (Environment Agency, unpublished data).
	57
Table 3.1	Details of common bream tagged and tracked during the study. Individual tag number, tag type, capture method, date tagged, capture site, mass (kg), fork length (mm), age determined by scale analysis, gender, nominal tag delays, tag life, length of dataset collected (days), projected tag expiration date and whether the tag automatically switched itself off (y) or ran until batteries ran flat (n).
	65-8
Table 3.2	Mann-Whitney U tests results for comparisons of TDM and LR during ten day period post- surgery between freshly tagged (T) and control (C; previously tagged) fish during all five tagging events. Significant at < 0.005 following Bonferroni correction.
	83
Table 3.3	Mann-Whitney U tests results for comparisons of TDM and LR between fish captured and translocated ~35 km downstream, and control fish not captured or translocated.
	85
Table 5.1	The number of individual fish and the number of days between ten day home-range calculation periods used to determine the effects of seven independent artificial water level manipulations.
	113
Table 5.2	Two sample t-test results for adjacent month pairings around artificial water level manipulations. Levels are manipulated biannually in spring (1 <sup>st</sup> April) and autumn (1 <sup>st</sup> November). Asterisks denote significance at < 0.004 following Bonferroni Correction.
	117
Table 6.1	Sample size and sampling periods for the analysis of social networks for each group.
	126

<b>List of figures</b>	<b>Page</b>
Figure 1.1	The lower reaches of the unmodified River Rees, South Island New Zealand, with complex meander and diverse flow patterns creating lentic and lotic habitat types (authors own photo). 18
Figure 1.2	Common bream distribution map redrawn from Wheeler (1978). 24
Figure 1.3	‘Coarse fish migration occurrence causes and implications’ reproduced from Lucas <i>et al.</i> (1998) modified from Northcote (1978). 30
Figure 2.1	Map of the East Midlands and Lincolnshire. The lower River Witham flows from Lincoln to Boston and discharges into The Wash. 35
Figure 2.2	Plan of the lower River Witham in 1762 from Wheeler (1990). 38
Figure 2.3	Map of land use within the River Witham Catchment Flood Management Plan (CFMP) area (Environment Agency, 2008). 40
Figure 2.4	Mean daily flow ( $\text{m}^3 \text{s}^{-1}$ ) of the Barlings Eau at Langworth bridge, this tributary generates the largest flow input into the lower river. 42
Figure 2.5	Langworth bridge gauging station on the Barlings Eau, a tributary of the lower river, during normal conditions (left) and the summer flooding (right) in June 2007 (authors own photos). 43
Figure 2.6	Mean daily river level of the lower River Witham at Kirkstead Bridge from 1999-2010. 44
Figure 2.7	The heavily modified lower River Witham looking downstream from Dogdyke (below the confluence with the River Sleas), canalised and straightened with high floodbanks preventing connection with the floodplain (authors own photo taken at winter retention level). 44
Figure 2.8	The heavily modified lower River Witham looking downstream from Tattershall Bridge, canalised and straightened with high floodbanks preventing connection with the floodplain (authors own photo taken at winter retention level). 45
Figure 2.9	Aerial photograph of the lower River Witham, upstream half of the study area Bardney – Tattershall Bridge (maps.google.co.uk). 47
Figure 2.10	Aerial photograph of the lower River Witham, downstream half of the study area Tattershall Bridge – Boston (maps.google.co.uk). 47
Figure 2.11	A depth profile of the lower River Witham. 48
Figure 2.12	The contribution made by each species to density population estimates for years 1994-2009. 51
Figure 2.13	The contribution made by each species to biomass population estimates for years 1994-2009. 51
Figure 2.14	A comparison of mean ( $\pm$ confidence intervals) fish biomass estimates for fish $> 99 \text{ mm}$ recorded during the surveys 1994-2009. 52
Figure 2.15	A comparison of mean ( $\pm$ confidence intervals) fish density estimates for fish $> 99 \text{ mm}$ recorded during the surveys 1994-2009. 52

Figure 2.16	Common bream lengths from seine netting surveys 1994-2009.	53
Figure 2.17	Fish densities recorded during the spring 2008 hydro-acoustic survey.	55
Figure 2.18	Fish densities recorded during the autumn 2008 hydro-acoustic survey.	56
Figure 3.1	Map of the lower River Witham between Lincoln and Boston, the study area runs from the old river loop upstream of Bardney to the tidal limit at Boston.	63
Figure 3.2	Frequency distribution of the length of the tagged common bream.	64
Figure 3.3	Seine net catch of 990 common bream caught in a single 100 m haul on 3 <sup>rd</sup> December 2007 downstream of Bardney Bridge. Mobile tracking revealed the presence of several previously tagged fish in the vicinity which enabled the area to be targeted with seine netting (authors own photo).	64
Figure 3.4	Anaesthetic and recovery tanks (authors own photo).	71
Figure 3.5	Removing scales from the incision site (authors own photo).	71
Figure 3.6	Making the incision (authors own photo).	72
Figure 3.7	Tag insertion (V9 model; authors own photo).	72
Figure 3.8	Closing the incision with a single suture (authors own photo).	73
Figure 3.9	Using Orabase™ protective paste to provide a temporary barrier (authors own photo).	73
Figure 3.10	Post-surgery fish were placed in the recovery tank until they had recovered their righting reflex and were actively swimming (authors own photo).	74
Figure 3.11	Antibiotic injection of 1 ml Baytril at 2.5% directly behind the dorsal fin was given while the fish was recovering. The injection was made on the same line as the erect last dorsal ray between the two main muscle blocks (authors own photo).	74
Figure 3.12	Fish were retained in-river in holding nets for an hour after the last fish was tagged, before collective release as a group (authors own photo).	75
Figure 3.13	Releasing a tagged common bream on 10 <sup>th</sup> October 2010 at Bardney Bridge (authors own photo).	75
Figure 3.14	Receiver coverage and sample size of tracked fish. Grey bars represent the number of VR2 and VR2W acoustic receivers used to track the fish and, solid black line with black diamond markers, the number of fish tracked during the study period. No tracking took place during July-September 2008, as existing tags had expired and tagging new fish was delayed until water temperatures reduced during the autumn / winter months due to the benefits to fish recovery at higher dissolved oxygen levels.	76
Figure 3.15	Identifying and isolating recaptured tagged fish on the banks of the Sincil Dyke. Fish were placed in tanks and tagged fish identified with a VR100 (Vemco, Nova Scotia, Canada)	

	mobile tracking unit (authors own photo).	77
Figure 3.16	Releasing a translocated fish at Langrick Bridge 35km downstream from capture site (authors own photo).	78
Figure 3.17	Data collected each month. Solid black line and black square markers is the total number of detections collected, dark grey line with triangular markers is the number of detections per tagged fish present and the light grey line with diamond markers is the number of detections per receiver in each month of the study.	80
Figure 3.18	Projected sample size compared with the actual sample size. Solid black line with black square markers represents the projected sample size assuming no fish left the study area, grey line with grey diamond markers is the observed number of tagged fish within the study area.	81
Figure 3.19	Incision site of fish 6073 - 51 days post surgery. Note the very clean advanced healing and incision closure, suture still present (authors own photo).	81
Figure 3.20	The mean distance moved ( $\pm$ SE) for all five tagging events over the ten day period after tagging and release of freshly tagged fish (n=61) and control tracked fish (n=55).	82
Figure 3.21	The mean linear range ( $\pm$ SE) for all five tagging events over the ten day period after tagging and release of freshly tagged fish (n=61) and control tracked fish (n=55). Positive values denote ranges upstream whilst negative values denote ranges downstream.	82
Figure 3.22	Fish plots for A; tag ID 12255 (male), B; 12257 (female) and C; ID 12266 (male) showing behaviour before and after translocation. Return journeys to Sincil Dyke shown in red, which took 11, 6 and 24 days after release at Langrick Br respectively.	84
Figure 3.23	The mean total distance moved ( $\pm$ SE) over concurrent five day periods up to twenty five days after release for the translocated group (n=3) and non-translocated control group (n=7).	85
Figure 3.24	The mean linear range ( $\pm$ SE) over concurrent five day periods up to twenty five days after release for the translocated group (n=3) and non-translocated control group (n=7). Positive values denote ranges upstream whilst negative values denote ranges downstream.	86
Figure 4.1	Map of study area; the lower River Witham, showing the river zone number (1-7) to which receivers were allocated for the 'monthly longitudinal distribution' analysis. Place names are also included, tributaries are labelled in italics and capture and release sites are labelled by capital letters (see Table 1). Insert (top right) is an expanded view of the study area showing Bardney Lock 'z' which is generally impassable to fish, the old river loop that is fed from the main navigation channel at an impassable weir 'w', and the Sincil Dyke which contains more than one receiver ('x' and 'y').	94
Figure 4.2	Flow ( $\text{m}^3 \text{s}^{-1}$ ; black line) gauged on the Barlings Eau at	

	Langworth bridge (~6 km upstream of the study area) and river level (m ODN; grey line) recorded at Kirkstead bridge, near the Timberland Delph confluence (see Figure 5.1) approximate midpoint of study area.	96
Figure 4.3	Mean ( $\pm$ SE) 'Monthly distance' (km; grey columns) in relation to mean monthly air temperature ( $^{\circ}$ C; black line). Gap between July and September 2008 represents period of no tagged fish in the study.	98
Figure 4.4	Mean ( $\pm$ SE) 'Monthly distance' (km; grey columns) in relation to mean monthly air temperature ( $^{\circ}$ C; black line), averaged over the twelve month cycle.	99
Figure 4.5	'Monthly longitudinal distribution': zone 1 (upstream) to zone 7 (downstream).	99
Figure 4.6	'Monthly longitudinal distribution' averaged over the twelve month cycle for each zone of the river: zone 1 (upstream) to zone 7 (downstream).	100
Figure 4.7	'Lateral habitat use' mean proportion of total detections recorded in the shallow side-channels (black), the deeper Sincil Dyke (light grey) and the main river channel (dark grey).	103
Figure 4.8	'Lateral habitat use' mean proportion of total detections recorded in the shallow side-channels (black), the deeper Sincil Dyke (grey) and the main river channel (dark grey) averaged over the twelve month cycle.	103
Figure 4.9	Daily detections per tagged fish present, recorded in the shallow tributaries (grey columns) and mean daily water temperature ( $^{\circ}$ C; black line).	104
Figure 4.10	Daily detections per tagged fish present, recorded in the Sincil Dyke tributary (grey columns) and mean daily flow recorded at Langworth Bridge on the Barlings Eau ( $\text{m}^3\text{s}^{-1}$ ; black line).	105
Figure 4.11	Pre-spawning behaviour, males guarding marginal territories, observed in Nocton Delph May 2007 (authors own photo).	106
Figure 4.12	Aggressive interaction between two males occupying adjacent marginal territories, Nocton Delph May 2008 (authors own photo).	107
Figure 5.1	Water level (metres above Ordinance Datum Newlyn, m ODN) recorded at Kirkstead Bridge (midpoint of study area) during two of the seven artificial water level manipulations, illustrating the periods of flux in levels when changes are made and the selection of ten day sampling period's pre- and post-level manipulations. A; level manipulation in autumn 2007 (summer level to winter level), 5 days between sampling periods due to level flux, (i) = ten day sampling period pre-manipulation and (ii) = ten day sampling period post-manipulation. B; level manipulation in spring 2008 (winter level to summer level), 18 days between sampling periods due rainfall events, (iii) = ten day sampling period pre-manipulation and (iv) = ten day sampling period post-manipulation. Shaded areas represent periods of transition.	112
Figure 5.2	Mean seasonal 100 % (dark grey) and 90% (light grey) home-	

	ranges ( $\pm$ SE) for common bream in spring (n=92), summer (n=32), autumn (n=86) and winter (n=98). Common lettering indicate statistical equivalence following <i>post-hoc</i> Tukey test ( $P < 0.05$ ).	115
Figure 5.3	Mean 90% home-ranges ( $\pm$ SE) for common bream in January (n=75), February (n=79), March (n=98), April (n=89), May (n=83), June (n=47), July (n=29), August (n=21), September (n=19), October (n=44), November (n=40), December (n=72). Months during winter level retention period are coloured light grey and months during summer level retention period are dark grey. Common lettering indicate statistical difference following two sample t-test (significance at $< 0.004$ following Bonferroni correction) of adjacent month pairings around artificial water level manipulations (Table 5.2).	115
Figure 5.4	Mean 100% (solid symbols, solid line) and 90% (open symbols, broken line) home-ranges ( $\pm$ SE) for individual common bream tracked during the ten days pre- and post-artificial water level manipulations. Level manipulations occurred bi-annually in spring (n=56; winter level to summer level) and autumn (n=24; summer level to winter level).	116
Figure 5.5	Mean total recorded distance moved ( $\pm$ SE) for individual common bream tracked during the ten days pre- and post-artificial water level manipulations. Level manipulations occurred bi-annually in spring (n=56; winter level to summer level) and autumn (n=24; summer level to winter level).	117
Figure 6.1	Social network structure (mean number of interaction days per month) for the 7 groups (A-G Table 6.1) constructed. Connecting lines are weighted to indicate the number of interaction days. Females are indicated by open nodes, males by shaded nodes. Numbers next to nodes are the individual fish (tag) ID numbers.	127-9
Figure 6.2	Mean monthly weighted-degree per group ( $\pm$ SE). Weighted-degree represents how many other individuals each individual is connected to, taking account of the number of interactions between individuals.	130
Figure 6.3	Mean monthly weighted-degree over the 12-month cycle ( $\pm$ SE) for all groups.	131
Figure 6.4	The influence of time from tagging event on the mean monthly weighted-degree for all groups ( $\pm$ SE).	131
Figure 6.5	Frequency histogram of the overall weighted-degree for all fish.	132
Figure 7.1	Metheringham delph which was disconnected from the main river in the 1990s. The delph could be reconnected to the lower river to provide additional spawning, refuge and nursery habitat. In this photograph the main river runs perpendicular to the delph, behind the position of view (authors own photo).	137
Figure 7.2	The lower river at Chapel Hill at winter retention level with marginal habitat high, dry and exposed to frost. Varying water levels prohibit the establishment of permanent marginal habitat vegetation essential for fish and aquatic wildlife (authors own photo).	139

## **Acknowledgments**

I am extremely grateful to my supervisors Dr Charles Deeming and Dr Paul Eady for their continuous support and advice, and to Dr Tom Pike of Lincoln University for his assistance in constructing the social networks presented in Chapter 6. Many thanks to Ian Wellby of Blueroot for advice and help with fieldwork and to Chris Reeds and Dr Chris Extence of The Environment Agency for managerial support and funding that allowed this work to take place. Environment Agency fisheries staff Dave May, Phil Thornton, Jake Reeds, Wal Potter, John Smith, Keith Bone, Andy Beal, Ellie Chapman, Reuben Page, Andy Hindes, Rupert Bucknall, Alan Henshaw, Dave Green, Robert Pitkin and Dan Mainwaring all provided invaluable help with fieldwork. I also wish to thank Dr Martyn Lucas of Durham University, Jim Lyons, Paul Hyatt, Andy Hindes, Richard Chadd & Dr Graeme Peirson of The Environment Agency and Dr Jon Bolland & Professor Ian Cowx of Hull International Fisheries Institute for their constructive comments on experimental design, early findings, thesis structure, data analysis and manuscripts. The views expressed in this thesis are those of the author and not necessarily those of The Environment Agency.



## Abstract

Anthropogenic influences on riverine ecosystems have led to numerous impacts that have compromised fish communities. Large European lowland rivers have been amongst the most impacted aquatic environments globally. However, paradoxically, these environments have also been amongst the least studied, due the inherent difficulties of studying fish in large, turbid, dynamic environments. Common bream *Abramis brama* (L.) populations are amongst those impacted by reduced lateral and longitudinal connectivity that are a consequence of modifications to large lowland rivers in Europe. Thus common bream in the lower River Witham, Lincolnshire, UK, were selected as a case study to examine how fish utilize these river systems.

A literature review was carried out to assess the influence of human intervention on riverine ecosystems and their impacts on fish communities. In addition the use of telemetry techniques for studying fish populations was reviewed.

An ecological assessment of the lower River Witham was carried out using data available from The Environment Agency monitoring and hydrological systems and data specifically collected during this study. The history of this highly modified lowland river was reviewed to give perspective, timescale and context to the degree of modification that has been undertaken on the river and its consequences. Chemical and biological water quality indicators showed improvements in the last twenty years. Fisheries data identified three species that had become locally extirpated in the lower river since the 1800s. The river is now roach dominated, but common bream are still present, possibly due to available lateral spawning habitats that provide surrogates for, and are functionally similar to, the lentic floodplain waterbodies of natural riverine ecosystems. Hydroacoustic surveys show that fish communities are aggregated and favour the upstream half of the lower river where the channel is more heterogeneous. Recent changes in river character have made traditional fishery assessments by seine netting less effective.

Data collected during fish tracking studies are most valuable when the tagged fish are behaving naturally, thus the effects of tag attachment should not impact on the behaviour or well being of the fish. Acoustic telemetry was used to investigate the impacts of tagging, surgical inter-peritoneal implantation, and translocation of common bream. Fish were recaptured following surgery and the incision site photographed to assess healing, which was clean in all cases. All recaptured fish appeared to be behaving normally as they were part of large shoals of fish. On five separate occasions fish were tagged when other previously tagged fish were present allowing their behaviour to be compared in terms of the distances moved and linear range; there were no significant differences between the two groups indicating the tagging procedure to have no detectable impact on the short-term behaviour of the fish at the resolution of the tracking undertaken. Three recaptured fish were translocated ~35 km downstream; these fish appeared to exhibit homing behaviour, returning to the capture site between 6-24 days.

Their level of activity (in terms of distance moved) did not significantly differ from non-translocated fish, but their linear range was larger.

The distribution and habitat use of adult common bream was assessed by tracking their longitudinal and lateral movements over forty-three months, producing a dataset of over 3.1 million detections. A positive relationship between activity and temperature was revealed, with common bream moving greater cumulative distances during the warmer months and occupying a greater longitudinal proportion of the main channel. The occupancy of tributaries was related to temperature, with common bream entering shallow tributaries during rising temperatures in the spring, whilst a deeper, slow-flowing tributary was used more frequently during the cooler autumn/winter months. During the autumn, occupancy of this deeper tributary was positively related to flow rate, suggesting that common bream use this as a refuge from high main channel flow.

The home range concept is at the centre of theoretical models to explain the spatio-temporal behaviour of a wide range of animals including fish. There are lots of different ways of expressing home ranges, including areas, but since rivers are principally linear systems, most range is reflected linearly. Home ranges were calculated for 100% and 90% of locations. Seasonal differences were identified with fish occupying larger home ranges during the spring and summer opposed to autumn and winter. Artificial water level manipulations, management actions for the control of flood risk, affected home range size with fish occupying smaller ranges at reduced winter levels than at increased summer levels.

Analysis of when and how often individuals within a population interact with one another provides a method to study the social organisation of animals with the potential to reveal ecologically significant aspects that would otherwise have remained hidden. Network analysis and social network theory were used to examine the social interactions of the tracked fish. The extent of sociality appears to follow a normal distribution, such that there was no evidence of discrete classes of social/non-social fish. Fish were more social immediately after tagging and less social as time progressed, as such little evidence could be found to support the hypothesis that these groups of fish remained in each others' company for extended periods. However, more active individuals were found to be more socially connected, but home-range size did not significantly affect sociality.

The functions of these behaviours are discussed along with potential management and rehabilitation strategies for the lower River Witham and other heavily modified lowland rivers in order to meet good ecological potential/status under the Water Framework Directive (2000/60/EEC).

## **Chapter 1 Rivers, fish and human interventions.**

### **1.1 General introduction**

Riverine ecosystems represent complex systems which drain water falling on the earth's surface and conduct it to the sea (Welcomme, 1994). These are considered to be amongst the most human-degraded ecosystems worldwide (Malmquist & Rundle, 2002; Huckstorf *et al.*, 2008). River modification to facilitate human civilization, has led to flow regulation, channelisation and habitat degradation which has impacted fish populations (Welcomme, 1994; Pinder, 1997). Large lowland rivers support a significant proportion of the world's fish diversity (Huckstorf *et al.*, 2008) and the majority of such environments in Europe have been modified to some extent (Cowx & Welcomme, 1998). The long-term degradation of rivers can disrupt the longitudinal and lateral river continuum which can separate individuals from specialised / essential resources or habitats and which can result in population fragmentation; small populations are more prone to extinction than large ones and with fragmentation, recolonisation from other populations is less likely (Cain *et al.*, 2011). Anadromous species, such as sturgeon *Acipenser* spp. and Atlantic salmon *Salmo salar* (L.), are often lost locally, species diversity declines and the sustainable use of resources is lost (Junkwirth, 1998).

Rehabilitation of riverine ecosystems is currently being driven in the European Union by the Water Framework Directive (2000/60/EEC). This legislation establishes a framework for the protection of groundwater, inland surface, estuarine (transitional), and coastal waters. Objectives are to prevent water ecosystem deterioration, to protect and to enhance the status of water resources and, most importantly for rehabilitation, is to achieve a 'good ecological status' for all waters, by 2015 (Borja & Elliot, 2007). The directive also requires 'artificial waterbodies' (those created by human activity) and 'heavily modified waterbodies' (a water body subjected to physical alterations by human activity, which substantially changes its hydro-geomorphological character) to achieve at least 'good ecological potential' by 2015 (Borja & Elliot, 2007). The ecological potential of a waterbody represents the degree to which the quality of the waterbody's aquatic ecosystem approaches the maximum it could achieve, given the heavily modified and artificial characteristics of the waterbody that are necessary for the use or for the protection of the wider environment (UK TAG, 2008). This ecological quality is based upon the status of the biological (phytoplankton, macroalgae, macrophytes, benthos and fishes), hydro-morphological and physico-chemical qualities of the waterbody (Borja & Elliot, 2007).

Wolter (2001b) has argued long-term studies of fish communities in European lowland rivers are lacking. In particular there is a paucity of knowledge about the spatio-temporal behaviour and ecology of cyprinid fish in heavily modified large lowland rivers. This may be a result of problems in sampling adult fishes in large rivers (Wolter & Freyhof, 2004), which has made data on habitat use and movements scarce. Aquatic telemetry offers the opportunity for new insights into the ecology of fishes and acoustic tracking is ideally suited for studying these large lowland riverine ecosystems. Using the lower River Witham as a case study, the overall aim of this study was to attempt to fill knowledge gaps in the long-term movements and habitat

use of cyprinids with a view to informing management strategies to conserve and restore fish stocks. The study was divided into topics that are addressed in Chapters 1 to 7. Specific objectives and hypotheses are stated at the beginning of each chapter.

**Chapter 1** reviews the current literature on fish ecology in riverine ecosystems, anthropogenic impacts on lowland rivers, the use of aquatic telemetry and the ecology of common bream.

**Chapter 2** uses ecological data to assess the current status of the lower River Witham calling upon historic and recent data on fish populations and how these have changed over time.

**Chapter 3** describes an investigation of the effects of tagging and long range translocation on the spatial temporal behaviour of common bream in the lower River Witham.

**Chapter 4** describes an investigation in to the distribution and habitat use of eighty three adult common bream that were tracked in the main channel and tributaries over a four year study period, with particular emphasis on identifying spawning and refuge habitats.

**Chapter 5** describes an investigation into the effects of season and water level management on the home range size of adult common bream.

**Chapter 6** describes an investigation into the social interactions of the tracked fish.

**Chapter 7** summarises how our improved understanding of common bream ecology can be placed in the context of targeted management actions aimed at meeting good ecological potential/status under the Water Framework Directive (2000/60/EC).

The information in this thesis is intended to inform long-term management and rehabilitation strategies of large heavily modified lowland riverine ecosystems, to minimise potential impacts and maximise potential enhancements. Thus helping to improve and enhance fish communities in the future, when pressures on such systems are likely to be even greater than they are today.

## **1.2 Objectives**

The overall objective of this chapter is to review current literature relevant to understanding fish ecology in riverine ecosystems and how anthropogenic effects have shaped lowland rivers worldwide. The use of aquatic telemetry is also discussed with an emphasis on its application to the study of common bream in the lower River Witham, Lincolnshire, UK. The specific objectives are to:

1. Describe a 'natural' riverine ecosystem.
2. Summarise human pressures on lowland rivers and how these have shaped large rivers;
3. Provide an overview of the ecology of fish in riverine ecosystems;
4. Summarise common bream ecology, with reference to management and rehabilitation strategies aimed at meeting good ecological potential/status in line with the EU Water Framework Directive (2000/60/EC).
5. Discuss the use of telemetry techniques and their applicability to study fish in the lower River Witham;
6. Outline the aims of this study.

### 1.3 A 'natural' lowland riverine ecosystem

Rivers transport run-off from precipitation and the amount of run-off is dependent on the timing, intensity and duration of rainfall events. During periods of high precipitation river discharge will increase to the point where the river floods, spreading across onto the land. Junk *et al.* (1989) referred to these areas of land which are routinely inundated in this manner as the aquatic / terrestrial transition zone or the floodplain. The lower reaches of rivers offer lateral habitats, within the floodplain which are exploited by fish during seasonal flood pulses (Junk *et al.*, 1989). Unmodified large rivers consist of two components; diverse active channel(s) through which water flows at all times and, the floodplain over which water flows during times of flood, seasonally altering between terrestrial and aquatic phases with a range of residual lakes and swamps (Welcomme, 1994). Natural erosion and deposition processes cause diversity in both components. River channels meander from side to side directing the path of the main flow from one side of the channel to the other, with local geology and topography governing the route rivers follow. Where the main flow swings towards the bank erosion occurs and where the main flow swings away from the bank material is deposited causing sinuosity of the channel (Ward *et al.*, 1999). Meander arms grow giving rise to more complex meander patterns increasing the channel length (Figure 1.1). Highly developed meanders are cut off through erosion at their neck, giving rise to oxbow lakes that can become detached from the main channel only reconnecting during times of flood. These same processes of erosion and deposition shape the floodplain, creating ecotones between the terrestrial and aquatic environments, with flooding and fluvial dynamics maintaining diverse lentic, lotic and semi-aquatic habitat types (Ward *et al.*, 1999). Fluvial action may create other habitat features such as channels, dead arms, side-channels and marshes, while terrestrialsation remoulds habitats during periods of low flow (Ward *et al.*, 1999). Deposition and erosion, energised by floods, are dynamic processes which are at equilibrium resulting in a consistent channel length with many natural habitat types and features which are in a constant state of flux.

According to Ward & Stanford (1995) river systems consist of three spatial dimensions: longitudinal (river – river or tributary), vertical (river - aquifer) and lateral (river - floodplain); and the fourth dimension: time. Interactions between these dimensions encompass the long-term development of a river system and the shorter term processes of the hydrological cycle. The seasonally variable lateral and longitudinal connectivity is a key ecological feature of fish species in lowland river floodplain systems (Jungwirth, 1998) with the diversity of floodplain habitat providing vital species-specific niches for fishes to complete their life cycles (Copp & Peñáz, 1988; Copp, 1989). For example, Bolland *et al.* (2012) investigated assemblages of juvenile fishes by electro-fishing in ten artificial floodplain waterbodies of the lowland River Trent, UK, with variable hydrological connectivity, in which waterbodies ranged between being permanently connected to the main river channel and only connected during elevated levels. The degree of connectivity of waterbodies was positively correlated with species diversity and fish communities in poorly connected waterbodies were most dissimilar to communities in the main



**Figure 1.1** The lower reaches of the unmodified River Rees, South Island New Zealand, with complex meander and diverse flow patterns creating lentic and lotic habitat types (authors own photo).

river channel, demonstrating the importance of variable lateral connectivity in lowland riverine ecosystems.

#### **1.4 Anthropogenic impacts on lowland riverine ecosystem**

River floodplains have been at the centre of human settlement from the origins of civilization (Welcomme, 1985). Rivers provide a host of goods and services for society *e.g.* drainage, water resources, food, power generation and recreation. Thus society has modified river channels and floodplains for our own purposes. Navigation, power generation and water storage has led to the construction of dams, locks, weirs and impoundments causing barriers, which prevent upstream access for migratory fish and reduce the quantity of specialist habitats available for spawning (Welcomme, 1985). These impacts have been long understood; in 1215 the Magna Carta demanded the removal of numerous weirs along the River Thames so that migratory Atlantic salmon and sea trout *Salmo trutta* L. could pass upstream to access their spawning grounds, however there was little response to this edict (Mann, 1988). Modification of the river discharge below dams also impacts on both water quality and flow regime (Mann, 1988). Industrial, agricultural and residential development in the floodplain has led to reductions in the size and function of floodplains. Increased human occupation of the floodplain has been driven by demands for rich agricultural land, access to river channels for transport, power generation and a general need for living space. This has led to the progressive deforestation of floodplains allowing their conversion from semi-aquatic to terrestrial habitats, stabilisation of watercourses

and the drainage of swamps and floodplain waterbodies (Welcomme, 1994). Flood defences are built to protect people's lives, property and businesses so as to sustain economic activity. River modification for flood risk management aims to contain rivers through channalisation; the straightening, deepening and widening of natural river channels to increase capacity and the building of flood banks to exclude floodwaters from land and property (Flemming, 2002). Increasing drainage and minimising the area inundated significantly affects the rivers natural features and thus adversely impacts habitat diversity in the floodplain. Flood risk management also impairs the natural hydrograph necessary to maintain and shape habitat features within the river and floodplain alike (Junk *et al.*, 1989). Abstraction from both ground and surface waters for industrial and domestic supply and water transfer schemes modify flow regimes and have had widespread effects on fish populations. These include low flows, which affect water quality and migratory behaviour, thermal discharges, mortalities by entrainment into pumps and impingement on intake screens (Mann, 1988) and translocation of fish species (juveniles and viable eggs can be transported unintentionally with water) to new environments impacting on native populations (Gozlan *et al.*, 2010).

Direct interventions on lowland river systems are prevalent, with changes in river morphology generally shifting systems from lentic habitats, such as are found in backwaters, downstream of point bars, or behind overhangs and snags, towards lotic environments devoid of shelter and nursery space with reduced longitudinal and lateral connectivity (Welcomme, 1994). Loss of habitat diversity leads to reduced populations, local extirpation of species and other impacts such as lower genetic variability within fish populations (*e.g.* Wolter, 1999) potentially placing them at an evolutionary disadvantage. River modification generally reduces habitat diversity with fewer off-channel areas (Copp, 1997) which provide low velocity environments. Slack waters within a lowland river system are particularly important for zooplankton and micro-benthic assemblages and as a refuge for adult, larvae and juvenile fish (Humphries *et al.*, 2006). Floodplain and off-channel waterbodies provide essential habitats for fishes acting as both nursery grounds and refuges (*e.g.* Heermann & Borchertding, 2006). Without an adequate number of well-dispersed, off-channel areas along the river's course, young fish will be unable to find refuge during winter and spring floods and will be eventually swept downstream, possibly to the sea (Copp, 1997).

Hydroelectric power stations can have direct effects on fish through entrainment (*e.g.* Hadderingh & Bakker, 1998; Turnpenny, 1998); hydro-peaking, water diversion and poor quality water discharge from reservoirs can have negative consequences for aquatic habitats and organisms (Jungwirth, 1998). Management action on regulated rivers can impact on fish recruitment. For example, weed cutting to reduce flood risk through improved conveyance can reduce spawning and nursery habitat (Mann, 1988) and zooplanktonic densities impacting on fish growth and abundance (Garner *et al.*, 1996).

Lowland rivers are particularly susceptible to modification; relative higher flows in the lower reaches give better potential for power generation and their locality to sea ports leads to increased industrialisation and associated impacts. Also, actions upstream have implications

downstream; silt, pollutants and nutrients introduced or disturbed in the upper reaches are transported and can accumulate in the lower reaches. Run-off from agricultural land and sewage discharges has enriched lowland rivers leading to elevated levels of the plant nutrients nitrogen and phosphorus (*e.g.* Pinder *et al.*, 1997) leading to eutrophication, the effects of which (*e.g.* turbid waters, algal blooms, changes in plant and fish communities) may be problematic (Edmondson, 1991; O'Sullivan, 1995).

The impacts of these interventions on fish communities centre on the loss of, and the reduced ability of fishes to exploit, specialised habitats required for the completion of their life-cycle. River regulation is considered one of the main causes for declines in river fish populations (Peter, 1998; Northcote, 1998). Of nearly two hundred European freshwater fish species, sixty-seven are now considered to be threatened by human activity (Northcote, 1998). Major causes have been identified for forty-eight of these and over half are associated with obstructions to migratory pathways at dams, weirs or other alterations in river channel features (Northcote, 1998). Fish communities in unmodified large rivers are highly diverse, reflecting the structural diversity and habitat richness of connected floodplains (Schiemer, 2000) meaning the structure of the fish population reflects its overall quality. For example the relative abundances of specifically adapted floodplain species can indicate the quality and extent of lateral connectedness (Schiemer, 2000).

There are many examples of river modification leading to a decline in freshwater fish populations in Europe and elsewhere. The Great Ouse, a large lowland system in the east of England, has been dredged and channelised with the installation of numerous locks and sluices to maintain water levels in the interests of power generation, navigation and land drainage. This has reduced lateral habitats and caused deviation from a natural flow regime. Fish populations have changed, with a once diverse cyprinid community becoming roach dominated, with a notable decline in once abundant common bream, and the extinction of the burbot *Lota lota* (L.) (Pinder, 1997) and the European sturgeon *Acipenser sturio* L. (Pinder *et al.*, 1997). Recruitment of other limnophilic and rheophilic species like silver bream *Blicca bjoerkna* (L.), tench *Tinca tinca* (L.) and rudd *Scardinius erythrophthalmus* (L.) appeared low and their distribution sporadic when compared to a similar unmodified habitat the River Biebrza in Poland (Copp, 1990). Similarly, Wolter (2001a) surveyed nineteen modified (primarily for navigation) lowland waterways in Germany, which were found to be dominated by roach *Rutilus rutilus* (L.) and perch *Perca fluviatilis* L. Areas of artificial shoreline structures like riprap or sheet piling lead to reduced species richness, diversity and abundance of intolerant species. Similarly, Wolter & Vilcinskis (1997) carried out electro-fishing surveys of eleven lowland rivers in Germany. Two trends could be identified in the fish assemblage owing to human influences; an increase in the dominance of roach and perch, and a decrease in species number. In particular, intolerant fish with a preference for sand or gravel substrata and current diversity had become rare and endangered. The influence of habitat modification on fish assemblages was analysed by Pilcher *et al.* (2004). Biomass, density and species diversity were compared between parallel channelised (modified for navigation) and natural (unmodified) side loop sections of the Rivers



Lee and Stort in Eastern England. All three parameters were significantly lower in channelised sections than in natural sections. Native species such as the common bream, European eel *Anguilla anguilla* (L.), barbel *Barbus barbus* (L.), pike *Esox lucius* L., chub *Leuciscus cephalus* (L.), dace *Leuciscus leuciscus* (L.), perch and roach were more abundant in the natural river sections. Channelised sections were absent of the rheophilic barbel whilst favoured by the limnophilic tench.

River modification is extensive globally (see Welcomme, 1994) and future pressures are likely to put further stress on these environments. Human population growth will increase demand for agricultural output and increased development will lead to yet further modification of river corridors. The effect of climate change is likely to increase rainfall intensity leading to increased flood risk (Law, 2002) which will in turn increase demand for flood defence schemes and river management. Climate change is predicted to cause lower summer flows and consequently increased residence times which will increase nutrient, organic and biological contaminant concentrations with the potential to compromise water quality and increase phytoplankton growth (Johnson *et al.*, 2009). Thus, rivers in the southern UK may become a less favourable habitat for some species of fish, such as the Atlantic salmon, and may give rise to the temporary loss of ecosystem services, such as providing water of acceptable quality for abstraction (Johnson *et al.*, 2009). Measures to reduce carbon dioxide production have led to resurgence in small scale low-head hydropower (Environment Agency, unpublished data) which has the potential to fragment riverine ecosystems. However, such developments do present numerous opportunities to have improvements for fish passage brought in as mitigation measures agreed prior to their approval. Increasing oil prices will increase the cost of importing food and goods, producing pressure to grow more food locally which may result in increased fertilizer use and water abstraction.

### **1.5 Fish ecology and rivers**

Fish have evolved life history strategies that suit their environment. For example, Atlantic salmon construct gravel nests called redds in which they lay their eggs. Redds prevent the eggs from being washed downstream in the high flow environment in which they reproduce (Gallagher *et al.*, 2008). By contrast in lowland rivers common bream attach their eggs to aquatic plants in the lentic off-channel areas. Amongst fishes, reproductive strategies are diverse *e.g.* non-guarders; open substratum spawners and brood hidiers; guarders; substratum choosers and nest spawners; and bearers; external and internal, and this preference has been used to classify species into ecoethological guilds (Balon, 1975, 1981). Table 1.1 gives a simplified classification of fishes of modified European floodplain rivers into guilds by their habitat requirements (from Schiemer & Waidbacher, 1992; Nunn *et al.*, 2007).

**Table 1.1** Fishes of modified European floodplain rivers have been classified into guilds by their habitat requirements. Simplified from Schiemer & Waidbacher (1992) and Nunn *et al.* (2007).

Guild	Spawning, nursery and adult habitat requirement	Example species
Rheophilic	Flowing water to spawn and are usually found in fast-flowing, well oxygenated stretches of river where substratum is characterised by sand or gravel	Dace, chub & barbel
Limnophilic	Low velocity habitats such as floodplain waterbodies, back waters and oxbow lakes	Tench & common bream
Eurytopic	Less strict requirements for spawning and are thus able to establish populations in a wide range of waterbodies	Perch & roach

Upland reaches of rivers are characterised by cool spring water and steep gradients resulting in water which is both fast flowing and highly aerated. As rivers progress towards the lowland they become larger, gradients reduce, slowing flows and allowing deposition of silt which increases aquatic plant growth (Mann, 1988). Huet (1959) described how river habitats progress to this ilk; the trout zone has stony substrate with variable width and depth. This changes into the grayling zone; larger deeper streams with pronounced riffle-pool sequencing and bottom substrate a finer material than the trout zone. Both of these zones are predominantly habitats for salmonid fish species, phenotypic adaptation to these environments make for streamlined body shapes to minimise energy expenditure in high velocity currents, high dissolved oxygen concentrations and a relatively low temperature (Phillips & Rix, 1985). Next comes the barbel zone; fast flowing sections and quiet backwaters with increased aquatic plants, this zone is dominated by rheophilic fish species. Finally the bream zone; characterised by deep turbid water with slow currents found in lowland rivers and canals with extensive macrophyte growth (Mann, 1988), here limnophilic species dominate. Not all of these zones need occur in every river, nor every species in each zone, but these key habitat types have provided a practical basis for fisheries management (Mann, 1988).

In the dynamic aquatic environment, swimming is essential to survival. Mobility is necessary for predator avoidance, foraging and to satisfy daily and seasonal habitat requirements. River fishes constantly endure swimming against the flow simply to hold a stationary position. The aquatic environment, although dense, affords a low friction, weightlessness medium that coupled with fishes' streamlined body morphology allows almost constant travel at minimal energetic cost. As such, freshwater fish have the potential to be very mobile animals with the ability to move over large distances and exhibit daily and seasonal patterns of movements (see reviews by Northcote, 1984 and Lucas & Baras, 2001). Winter and Fredrich (2003) note some very large movements for cyprinids; black sea roach *Rutilus frisii* (N.) > 1000 km (Nikol'skii, 1961), Colorado squawfish (pikeminnow) *Ptychocheilus lucius* G. up to 400 km (Tyus, 1985), Nase *Chondrostoma nasus* (L.) 446 km (Steinmann *et al.*, 1937),

barbel 318 km (Steinmann *et al.*, 1937) and ide *Leuciscus idus* (L.) > 200km (De Leeuw & Winter, 2008) which demonstrate the potential these fishes have to move over large distances.


There have been many definitions of migration in fisheries science starting with Heape's (1931) distinction between migration "*A class of movement which impels migrants to return to the region from which they have migrated*" and emigration "*movements which entail change of environments, but which do not involve return to the original area of habitat*". Three types of migration were recognised: *alimentary* migration, in search of food; *climatic* migration, in search of more suitable environmental conditions; and *gametic* migration, for reproduction. Northcote (1998) defined migration as "*movements involving regular cyclic alternation between different habitats used for spawning, feeding, or survival*". While Lucas and Baras (2001) consider migrations to be "*out of the ordinary larger than usual movements for a specific purpose, usually seeking out specialised habitats*", while movements are simply "*everyday changes in the animals location*". Out of the ordinary movements not directly associated with migratory behaviour are called excursions and represent a little studied aspect of freshwater fish behaviour (Muhlfeld & Marotz, 2005; Knight *et al.*, 2008).

Migratory life-history strategies of fish have also been used to classify species, according to the following scheme:

1. **diadromous** fish travel between salt and fresh water. (Greek: '*Dia*' is between; Greek: '*Dromous*' is moving in a specific manner) There are three types of diadromous fish:
    - **anadromous** fish live in the sea, but return to freshwater in order to breed (Greek: '*Ana*' is up; The noun is "*anadromy*")
    - **catadromous** fish live in fresh water, but breed in the sea (Greek: '*Cata*' is down)
    - **amphidromous** fish move between fresh and salt water during some part of life cycle, but not for breeding (Greek: '*Amphi*' is both)
  2. **potamodromous** fish migrate within fresh water only. (Greek: '*Potamos*' is river)
  3. **oceanodromous** fish migrate within salt water only. (Greek: '*Oceanos*' is ocean)
- (Lucas & Baras, 2001)

## 1.6 Common bream ecology

The common bream, also known as bronze bream, is a predominantly benthic feeding cyprinid found throughout western and central Europe and into Asia (Figure 1.2; Wheeler, 1978; Maitland & Campbell, 1992). They are classified as potamodromous and limnophilic. Predominantly a shoaling species (Backiel & Zawiska, 1968; Phillips & Rix, 1985) common bream favour lowland, slow moving fresh and brackish waters with their distribution being limited by the conditions necessary for reproduction and embryonic development. Water temperature should not exceed 28 °C. Oxygen content requirements are different for specific life stages and should be no less than 1.8 mg/L for larvae, 5 mg/L for embryos and 0.3 mg/L for adults. Common bream are tolerant of salinity of up to 2.8‰ (Backiel & Zawiska, 1968). Heut's (1959) classification of rivers referred to the lower reaches of rivers as the 'bream zone' due to this species preference for and prevalence in this type of river habitat.



**Figure 1.2** Common bream distribution map redrawn from Wheeler (1978).

*“Quiet waters usually in the lower stretches of the streams (including ditches, canals and large rivers) make up the bream zone. In these waters current is slight, summer water temperature high and dissolved oxygen often fairly low. The water is often turbid and water depth may exceed two metres. Larger streams in the bream zone typically flow through a wide valley and often rivers of this type have been dredged and straightened to aid navigation. Calm-water cyprinids, especially bream, carp, and tench predominate, along with an abundance of roach” (Huet, 1959).*

Exceptionally deep bodied with a strong lateral compression adult common bream are easy to identify. They have dark grey, dark silver or brown body colouration, depending on water clarity, with dark brown fins. The tail is deeply forked with pointed lobes, and common bream have a long anal fin and a short but tall dorsal fin set back behind the pelvic fins (Phillips & Rix, 1985). The head is small, the back rather humped and the body is very slimy and covered by large strong cycloid scales that can be used for age determination (Maitland & Campbell, 1992). The age of a fish can be calculated by counting the number of rings, or “annuli”, laid down in a fish scale. During the colder winter month’s fish grow more slowly and scale deposits are laid down close together, forming a dark ring similar to a tree trunk. Rapid summer growth results in thinner deposits and the formation of a wider, lighter ring across the scale. These light and dark rings depict a history of growth over a period of time and therefore represent the age of the fish (Bagenal, 1973; Environment Agency, undated). Juveniles are silvery in colour, and can be confused with the smaller silver bream, however lateral scale (less than 60), dorsal (12) and anal (25-31) fin ray counts can be used for accurate identification and common bream also have a snout longer than the diameter of the eye (Wheeler, 1978; Phillips & Rix, 1985). Unlike many cyprinids, adult common bream are easy to sex determine according to the external features (Kennedy & Fitzmaurice, 1968). Rough texture of the skin and raised tubercles on the head, back and shoulders identify the males during the spawning season, although evidence of tubercles can be visible throughout the year. Common bream can grow up to 80 cm, although average adult size is between 25-45 cm. They are relatively long lived with individuals reaching

over 20 years of age (Maitland & Campbell, 1992), with exceptional individuals having been aged at 32 years (Backiel & Zawisza, 1968).

The common bream also readily hybridises with other cyprinids, especially with the roach to produce the commonest cyprinid hybrid in Great Britain (Cowx, 1983). Common bream x rudd hybrids (Maitland & Campbell, 1992) and common bream x silver bream hybrids are also common in some areas (Wheeler, 1978; Matondo *et al.*, 2009). These hybrids can sometimes be fertile and may breed with either species with the resulting offspring being nearer in character to the parent species (Cowx, 1983; Phillips & Rix, 1985). Hybridisation occurs when there is an overlap of spawning season and habitat. Changes in river morphology, canalisation and reduced lateral and longitudinal connectivity have the potential to increase the incidence of hybridisation by reducing the number of potential spawning sites (Cowx, 1983).

Molls (1999) describes an age-dependant habitat shift in the life-cycle of common bream between the main river Rhine, Germany, and its oxbows. Adult common bream of over 5 years of age migrated into oxbows to spawn before returning to the main river, although some remained resident. Young of the year and one year old juveniles were abundant in these oxbows until they emigrated to the main river where they remained until reaching maturity. No immature common bream between the ages of 2 and 5 years old were found in the oxbows, yet these were plentiful in the main river. This habitat shift may be related with feeding strategies, as these change with age and season.

Juveniles feed predominantly on planktonic prey items such as copepods, cladocera (Winfield *et al.*, 1983), cyclops, rotifers and diatoms (Backiel & Zawisza, 1968). By contrast adults favour benthic feeding (Cowx, 1983) with a mouth that can be extended as a feeding tube (Backiel & Zawisza, 1968), but will exploit locally and seasonally abundant pelagic food sources (Lammens, 1985; Schulz & Berg, 1987; Breukelaar *et al.*, 1994). Stomach contents of adult common bream from the Norfolk Broads were found to include Cladocerans (planktonic water flea) (31.0%) and Copepods (mostly planktonic and some benthic) (28.7%) and insect midge larve and pupae (13.1%) (benthic) suggesting both benthic and midwater planktonic feeding behaviour (Hartley, 1947). Goldspink (1978) confirms the general benthophagic habit of the species with oligochaeta *Limnodrilis* spp. (a benthic worm), chironomid larve *Tanytarsus* spp. and *Pentapedilium* spp. and the occurrence of large numbers of cladocera (water fleas) being a common features in the diet of large (>25 cm) common bream. Large common bream (total length 30-35 cm), feed pelagically by pump filter feeding, a series of rapid suction movements in which the premaxilla is protruded 100-200 times a minute briefly interspersed with closed protrusion movements when prey are filtered through the buccal cavity and swallowed. Smaller common bream (total length) 15-20 cm behave in a similar fashion while feeding pelagically, but are also particulate feeders, attacking prey individually, when prey density is low (Lammens, 1985). While feeding on benthic macroinvertebrates common bream tilt downwards from between 60 to 90 degrees (Svendsen *et al.*, 2005) and can cause the re-suspension of sediments, increasing turbidity and nutrient release into the water column (Breukelaar *et al.*, 1994), to such an extent that shoals of feeding fish can often be located by

spotting areas of churned up suspended mud and silt (Phillips & Rix, 1985). As such, the removal of common bream and other cyprinids has been used as a restoration technique (biomanipulation) in efforts to restore eutrophic lakes (Perrow *et al.*, 1997). Their removal not only prevents nutrient release from disturbed sediments, but also reduces predation on zooplankton thus promoting the numbers of these algae grazing animals which reduces blooms and restores clear water conditions that allows macrophyte establishment (Lammens, 1999). In the same vein, this benthic feeding behaviour by common bream has also prompted changes in the ecological quality of receiving watercourses following their introduction as non-natives (Volta *et al.*, 2013).

Feeding is reduced in both the spawning season and during the winter when water temperatures drop below 4-5 °C although common bream have been known to feed at temperatures as low as 0.5 °C (Backiel & Zawisza, 1968). Sudden drops in temperature are likely to stop feeding although once fish become acclimatised to lower temperatures, feeding at a reduced rate is likely to continue. Metabolic rate decreases at cooler temperatures (Hölker, 2006), slowing to 25% of that at 20 °C at 5 °C (Backiel & Zawiska, 1968),

Hickley and Dexter (1979) constructed expected lengths-at-age growth curves for common bream, demonstrating early fast growth reaching lengths of over 100mm in the first year, over 150 mm in their second year and ~200mm in their fourth year when they reach sexual maturity. Growth rates will be dependent on prey availability and thus vary from river to river and varies with gender. Marciak (1972) positively correlated growth rates with the standing crop biomass of invertebrates in mud. Sex variation in growth rates were observed in the River Exe, Devon UK, with males exhibiting rapidly declining growth rates after six years which was not paralleled in the female growth curve. This is thought to be a strategy for maximizing reproductive capacity as female fecundity increases with size (Cowx, 1983). Sexual maturity varies considerably with climate, ranging from between three to ten years, and is determined by 'physic logical age' being reached at lengths between 14-30 cm, with males frequently reaching maturity before females of the same age (Backiel & Zawisza, 1968). Described as semi-migratory, common bream of the Black Caspian and Azov seas are known to exhibit large migrations of up to 100 km during spring and autumn. The spring time spawning migration and over-wintering autumn migrations are both from brackish sea feeding areas into the lower reaches and deltas of large rivers (Backiel & Zawisza, 1968). Common bream spawning migrations in freshwater (Whelan, 1983; Schulz & Berg, 1987; Prignon *et al.*, 1998; Molls; 1999; Grift *et al.*, 2001; Lilja *et al.*, 2003) occur in May-July and can vary from 5-60 km (Lucas, 1998). These are triggered by rising water temperatures (Prignon *et al.*, 1998; Lilja *et al.*, 2000). Schulz & Berg (1987) observed long distance movements occurring more frequently during spawning time and increased shoaling behaviour (spawning aggregations) with movements between spawning sites up to 10 km apart in Lake Constance, Germany. Whelan (1983) showed that aggregations of common bream at a single spawning site on the River Suck, Ireland, broke up into four separate shoals when they returned to their feeding grounds up to 59 km away. Prignon *et al.* (1998) observed fish captures from 1989 to 1994 at the Tailfer Dam on the River

Meuse in Belgium. Movements of mature common bream occurred during mid-May to July when water temperature varied between 13-23 °C. Lilja *et al.* (2000) used echo-sounding and net traps to study fish migrations in a channel between a large mesotrophic and a small eutrophic lakes in Finland. Capture of common bream in the net trap peaked when water temperatures reached between 14-16 °C.

Spawning occurs at high water temperatures in the middle of spring - early summer after gonad maturation has taken place. The oocyte development is often asynchronous amongst fish of different sizes and ages (Rinchard & Kestemont, 1996; Fredrich *et al.*, 2003) which can fraction spawning into several temporal groups. Common bream generally spawn once a year, but have been known to spawn twice or three times near the limit of their southerly range (Backiel & Zawisza, 1968; Shikhshabekov, 1972). Spawning takes place when water temperatures range between 16-18 °C (Backiel & Zawisza, 1968) or reach more than 15 °C (Maitland & Campbell, 1992). Spawning is polygamous *i.e.* occurring repeatedly between different males and females (Backiel & Zawisza, 1968; Poncin *et al.*, 1996) in shallow weedy areas (Maitland & Campbell, 1992). The males are ready to spawn first, linger on the spawning grounds longer than females, becoming aggressive defending small territories from rival males (Maitland & Campbell, 1992; Giles, 1994). Other spawning strategies, *e.g.* lek and sneaking may be adopted by different males during spawning (Poncin *et al.*, 1996). Females can produce on average 100,000-200,000 eggs (although this can range from 2,000-900,000) and spawn several times until all their eggs are laid. Spawning behaviour is easily identified; fish are aggressive with much splashing and bow-waving. Common bream spawn in sheltered places where the water is either still or the current is weak most often in depths from 20-80 cm, although can range from 9 cm to 17 metres (Backiel & Zawisza, 1968). Where rivers are allowed to meander, common bream have been shown to spawn in the slack water of oxbows (Molls 1999; Borcharding *et al.* 2002). The temperature of the water is important, with first spawning occurring in shallower warmer water. In environments with variable water levels, common bream often spawn on inundated areas and success is dependent on hydrometeorological conditions (Backiel & Zawisza, 1968). Common bream are phytophilous and are known to spawn on macrophyte substrata (Pinder, 1997) and marginal reed beds (Giles, 1994). Eggs are between 1-2 mm in diameter and yellow in colour once fertilised outside the body. Natural fertilisation of eggs is high 90-100% (Backiel & Zawisza, 1968), they adhere to plants and hatch in about 5-10 days depending on water temperature (Maitland & Campbell, 1992), or from 1430-1890 degree hours at 20-22 °C (Backiel & Zawisza, 1968).

Once eggs hatch they develop quickly from immobile yolk-sac fed 4-5 mm long larvae to actively swimming fully developed fish of 20 mm feeding on zooplankton, green algae and diatoms after twenty days (Backiel & Zawisza, 1968). Through much of this development they are free drifting with any currents and reside near the shore and amongst plants in lentic habitats (*e.g.* Copp & Peñáz, 1988). Mortality of larvae is high, usually between 95-99% (Backiel & Zawisza, 1968). Fish then enter an adolescent phase lasting from 2-8 years, dependant on local

conditions, the basic period of development and organogenesis ends at the length of 75-107 mm and they attain their final body shape when longer than 14 cm (Backiel & Zawisza, 1968).

In central and eastern Europe, common bream are an important commercial species and are caught in large numbers by traps and seine nets, while it is a popular sporting fish in many areas of Great Britain (Maitland & Campbell, 1992). Natural predators of adult common bream include the otter *Lutra lutra* L. (Erlinge, 1968; Copp & Roche, 2003) and common seal *Phoca vitulina* L. (authors observation in the River Witham), juveniles will be taken by piscivorous fish such as pike and perch and birds such as cormorant *Phalacrocorax carbo carbo* (L.) (Britton & Shepherd, 2005) and grey heron *Ardea cinerea* L. (Feunteun & Marion, 1994). Common bream larger than 20 cm are attacked rarely and only by very large predatory fish (Backiel & Zawisza, 1968) or mammals, due to their lateral compression which makes them difficult to ingest whole.

Common bream have the potential to be a very mobile species. Large spawning migrations have been observed (see above) and common bream have also been observed making autumnal movements into drainage ditches of floodplain lakes in the lower Rhine, Germany, for over-wintering when water temperatures dropped below 10 °C (Borcherding *et al.*, 2002). Slow swimming is characterised by pectoral propulsion while faster swimming is gradually dominated by caudal propulsion. Body angle becomes more level as swimming speed increases, with slow swimming usually being associated with a head down posture (Svendsen *et al.*, 2005). Swimming speeds increase with length, 0.66 m/s for 12-16 cm long fish, 0.90 m/s for 24-28 cm long fish (Backiel & Zawiska, 1968). Swimming speed can be maintained for considerable lengths of time, with large quick movements being observed by both Langford (1979, 1981) and Lyons & Lucas (2002) using telemetry techniques. Floy tagging of 2,763 adult common bream re-released into the river Suck, Ireland, resulted in 110 recaptures, 35 were recorded at 2-59 km from the original tagging site while 75 were recaptured on site (Whealan, 1983). Goldspink (1978) marked 4,619 adult common bream with operculum chicken wing tags (Goldspink & Banks, 1971) in Tjeukemeer, a large shallow lake in the Netherlands. The majority of fish recaptures were from the same lake in which they were marked, but some fish also travelled through connecting channels to adjoining water bodies and were recorded up to 60 km from the release point. Homing behaviour has also been observed by common bream (Vostradovsky, 1975) and following translocation by Malinin (1971), Langford (1979, 1981) and Goldspink (1978).

Movements can be triggered by environmental factors. Alabaster & Robertson (1961) observed common bream, roach and perch behaviour in very large convoluted tanks forming a 500 foot (~150 m) long channel with many side bays where temperature and dissolved oxygen could be monitored and manipulated. Observations of roach were made for two weeks prior to any manipulations being made, in which time the fish moved routinely. During the daytime roach were only observed in two bays of the tank, at night fish were observed in these same bays but just after dawn the roach became very active and swam over considerable lengths of the tank's channel, this coincided with minimum recorded values of both temperature and



dissolved oxygen. Similar pronounced activity of perch and common bream was observed at dawn and also dusk. Shoals of both perch and common bream broke up in the evening into smaller groups, sometimes as small as a single fish, but at dawn both species came together forming a single group which remained intact throughout the day, demonstrating diurnal activity behaviours in common bream. These observations were maintained even when temperature was artificially maintained, suggesting these behaviour are a response to light levels as opposed to environmental variables. However, changes in temperature and dissolved oxygen concentrations did cause movements independent of these diurnal patterns in all three species. When fish were exposed to changing temperature or dissolved oxygen concentration they became more active and increased their range of swimming, tending to move away from changing conditions towards areas where conditions were nearer those to which they were acclimatized before conditions were manipulated.


### **1.7 Studying spatio-temporal behaviour of fishes with telemetry techniques**

For some large oceanic species annual migration circuits extend to as much as 10,000 km (Arnold & Dewar, 2001) and the migrations made by the Atlantic salmon and the European eel can be dramatic and relatively obvious. Humans have been exploiting these migrations to and from the sea where fish have congregated at natural barriers for thousands of years (Lucas & Baras, 2001) and specifically designed man made barriers and traps have been used to increase fishing efficiency. Scientific research on the spatio-temporal behaviour of fishes has been biased towards salmonids and has concentrated on their return migrations through natal freshwater environment required to complete their life cycle. This has been driven by the commercial value of salmon fisheries (Radford *et al.*, 1991); Peirson *et al.* (2001) placed a capital value of salmon fishing rights in England and Wales at £128 million.

However, potamodromous migrations of fishes can be subtle and difficult to detect, as such scientific study in this area has been limited. Migrations by freshwater fish occur between three types of habitat feeding / growth, spawning and refuge and as such migrations can be classified as feeding, refuge and spawning dependant on the destination motivation (Figure 1.3; Lucas *et al.*, 1998b). Spawning migrations have been observed in many cyprinids, for example barbel (Lucas & Batley, 1996) and chub (Fredrich *et al.*, 2003), these being triggered by increases in water temperature (Waidbacher & Haidvogel, 1998). Migrations to over-wintering habitats have also been observed in cyprinids *e.g.* in ide (Winter & Fredrich, 2003), and migrations between adult and juvenile habitat types have also been observed *e.g.* in common bream (Molls 1999).

The timing and size of such migrations will be species specific and environmentally dependent. In environments with an abundance of suitable habitat types, migrations are likely to be smaller than where habitat types are limited and fish are forced to travel longer distance in search of a certain habitat type.

Modern telemetry technology offers researchers a very powerful tool for studying the long-term movements of individual fishes, with high temporal and spatial resolution (see reviews by



**Figure 1.3** ‘Coarse fish migration occurrence causes and implications’ reproduced from Lucas *et al.* (1998b) modified from Northcote (1978).

Lucas & Baras, 2001; Jellyman, 2009). The use of electronic tags has been one of the most important advances in the study of the behaviour and distribution of fishes in the freshwater environment (Lucas & Baras, 2001). Beginning with the use of a simple coloured ribbon attached to the tails of Atlantic salmon to observe localised movements as far back as the 17<sup>th</sup> century (McFarlane *et al.*, 1990; Lucas & Baras, 2001) to modern day satellite technology monitoring movements of bluefin tuna *Thunnus thynnus* (L.) on a near global scale (Sibert *et al.*, 2006) the tagging and tracking of fish has developed as researchers have strived to gain more detailed knowledge of behaviour.

Battery operated acoustic and radio tags have a limited life span, as battery life is limited. There is a trade-off with tag life and size, limiting the duration of study if tag size is to be kept appropriate for use with anything but the largest of fishes. Radio transmitters are a popular method of studying animal behaviour (see Cochran, 1980; Priede & Swift, 1992; Cooke, 2008) and are suitable for fish research in low conductivity (< 500  $\mu\text{S}/\text{cm}$ ) environments (Cooke *et al.*, 2012) having been used frequently (*e.g.* Lucas & Batley, 1996; Huber & Kirchhofer, 1998; Kuiskova *et al.*, 2009). Acoustic or ultrasonic tracking dates back to its development by Trefethen-Parker (1956) and Henderson *et al.* (1966) and is usually applied in the marine environment as it is suitable for use in highly conductive environments where radio telemetry is not. However, it has rarely been used with cyprinids, for example with common carp *Cyprinus*

*carpio* L. (Johnson & Hasler, 1977) and common bream (Langford, 1974, 1979 & 1981; Schulz & Berg, 1987; Lyons & Lucas, 2002). Passive Integrated Transponders (PIT) telemetry has the advantage that no battery power is required within the tag itself and so offers a more long-term option. However PIT telemetry requires the tag to pass within close proximity (~1 m) to aerial receivers and so its application is usually restricted to streams and small rivers or where fish are forced to pass through narrow structures like fish passes (see Lucas *et al.*, 1999, Lucas *et al.*, 2000; Skov *et al.*, 2005). PIT, radio and acoustic tags are coded so individuals can be identified and distinguished from each other.

Tracking methods vary, and fishes may be tracked manually on foot (*e.g.* Lucas & Batley, 1996) or from a boat (*e.g.* Lyons & Lucas, 2002; Kuiskova *et al.*, 2009) or remotely with fixed receivers. Manual tracking studies rely on regular visits to the field and the physical location of each tagged individual to be determined and recorded. This technique tends to have better spatial coverage, assuming access to the whole study area is possible, and resolution by using triangulation of the signal to give an accurate ( $\pm <10$  m) position of the transmitter and thus individual. However unless individual tagged fish are monitored intensively data on fish locations can only be collected relatively infrequently thus lacking temporal resolution of fish location. By contrast fixed receivers offer data with complete temporal coverage, as they can operate continuously, but with limited spatial coverage and resolution as data will only be recorded when an individual is within range of a receiver and then this will only indicate that the tagged individual was within the range of the receiver, which may be several hundred metres. Previous tracking studies have relied heavily on manual tracking techniques which are both restrictive and labour intensive. Manual tracking is usually restricted to daylight hours and for relatively short periods which has the potential to introduce bias (Clough & Ladle, 1997; Baras, 1998). Manual tracking studies also vary in the temporal frequency of locations, for example; Kuiskova *et al.* (2009) recorded the position of radio tracked ide once every three hours on one day a week, over a 12 month period. While Lyons & Lucas (2002) located tagged common bream on at least three separate days each week, a total of 24 occasions, over 55 days, data collection occurred over four equal length time periods and on two occasions selected individuals were monitored throughout a 24 hour cycle. The time interval between fish locations may overlook certain movements, especially those diel movements to and from resting and feeding places (Clough & Ladle, 1997), and can lead to underestimations of both home range size and mobility (Ovidio *et al.*, 2000). Through sub-sampling, multiple daily locations for radio-tracked brown trout *Salmo trutta* L. down to single observations at 2, 4, 7 and 14 day intervals, Ovidio *et al.* (2000) demonstrated reductions of accuracy from 13% (2 day interval) to 48% (14 day interval) for home range size and 28% (2 day interval) to 82% (14 day interval) for mobility estimations. Ovidio *et al.* (2000) concluded that any interval of locations probably has the potential to produce bias compared to the actual movements of trout and that fish should be ideally monitored continually to obtain the most accurate results. The popular arrangement of a single mobile receiver is also restrictive, making the tracking of many individuals at the same time problematic (Lucas & Baras, 2001). Manual tracking may also impact on behaviour (*e.g.*

Winter, 1983; Lyons & Lucas 2002) as human presence is required close to the animal being tracked is necessary, which may be detected and influence the behaviour of the fish. It also generates the possibility of numerous errors and inconsistencies in field data recording and signal interpretation both within the same study and particularly between studies.

Alternatively, tracking can be conducted automatically with fixed receivers (*e.g.* Westerberg & Lagenfelt, 2008) which record the date and time tags are detected. Acoustic telemetry has recently benefited from technological advances. Automatic data logger acoustic receivers have been developed (Lacroix & Voegeli, 2000) and can be used to record fish movements 24 hours a day, 7 days a week. These have been used in the marine/estuarine environment to study Atlantic salmon smolts (Moore *et al.*, 2008) and European eel (Westerberg & Lagenfelt, 2008). In the freshwater environment they have been used to study sea trout, Atlantic salmon (Bendall & Moore, 2008) and European eel (Aarestrup *et al.*, 2008). These receivers standardise data capture to a certain extent but factors such as turbidity, macrophyte growth, background noise, algae and fouling can impede acoustic propagation and reception, reducing the detection range of tags locally. Continual monitoring with fixed receiver stations covering all of the 24 hour cycle has rarely been practiced in the freshwater environment or with cyprinids (see Lucas & Frear 1997; Lucas *et al.*, 1999; Lucas *et al.*, 2000; De Leeuw & Winter, 2008).

Cyprinid species form a large proportion of fish communities in European lowland rivers, but relatively little is known about their spatio-temporal behaviour in such environments (Lucas & Baras, 2001). Those studies that have been carried out on cyprinids have tended to concentrate on rheophilic species such as chub (Allouche *et al.*, 1999; Fredrich *et al.*, 2003), barbel (Baras, 1992; Baras, 1995; Baras & Philippart, 1989; Kalpers *et al.*, 1989; Baras & Cherry, 1990; Lucas & Batley 1996; Lucas & Frear, 1997; Baras, 1997) and dace (Clough & Ladle, 1997; Clough & Beaumont, 1998) in relatively small heterogeneous riverine environments. Interestingly some of the very first telemetry studies did look at limnophilic species in large environments, such as carp in Lake Mendota, USA (Johnson & Hasler, 1977), common bream in the rivers Witham and Thames (Langford, 1974, 1979, 1981) and common bream in lake Constance (Schulz & Berg, 1987), but recently research on limnophiles has been limited.

The timing of studies is crucial if seasonal differences (*e.g.* in habitat use) are to be detected, ideally studies should be carried out over more than one 12 month cycle so that seasonal patterns can be confirmed by repeated observations. Sample size is also paramount, with partial migrations (Chapman *et al.*, 2012a; Chapman *et al.*, 2012b) being observed in many fish species. Sample sizes should be sufficient enough to demonstrate behaviours at the population level rather than just the individual. However, most cyprinid telemetry studies have been confined to just a few individuals during a relatively short period of a few months (*e.g.* Winter & Fredrich, 2003) or a single 12 month cycle (*e.g.* Fredrich *et al.*, 2003). Studies covering more than a single annual cycle include; Winter & Fredrich (2003) where forty-nine ide were radio-tracked on a weekly basis over four years in two large lowland rivers in Germany & The Netherlands and De Leeuw & Winter (2008) who tracked 245 barbel, chub, ide and nase over three years in

rivers Meuse and lower Rhine in the Netherlands with transponders and fixed receivers. Studies investigating seasonal changes in behaviour have used tags that operate throughout the year (*e.g.* Lucas & Batley, 1996) while others have tracked fish in batches throughout the course of a year (*e.g.* Baade & Fredrich, 1998).

If a long-term telemetry study were undertaken in the lower River Witham, the ideal target species would be common bream. Common bream are prevalent in the lower river and would therefore be relatively easy to capture. Common bream are also large enough to accommodate a tag of sufficient battery life to gather data through different seasons allowing study of behaviours for different ecological functions like spawning and over-wintering. Studying roach would make an interesting comparison, but the capture of fish large enough to carry transmitters would be erratic. Common bream have also been shown to be affected by river modification (see Pinder *et al.*, 1997) because they use lateral habitats for spawning.

### **1.8 Proposed study**

It is proposed that a long-term passive telemetry study on the lower River Witham could yield valuable data on seasonal habitat use of fish within a heavily modified lowland river. Common bream would be the ideal study species as they are large enough to carry tags with suitable battery life, they are a typical species of lowland fish communities including the lower River Witham and their life-cycle within lowland rivers relies on habitat features that are impacted by river modification.

Large, lowland rivers often incorporate important recreational fisheries (Lyons *et al.*, 2002) and are often heavily influenced by human activity, resulting in uniform systems with regard to both physical form and water flow (Brookes, 1988) that present many challenges to fish populations (Linfield, 1985). As described above, direct interventions on lowland river systems for the purposes of navigation, power generation, flood control, land drainage and flow regulation are prevalent, with changes in river morphology generally shifting systems from lentic towards lotic environments (Welcomme, 1994) through reduced longitudinal and lateral connectivity. The impacts of these interventions on fish communities centre on the reduced ability of fishes to exploit specialised habitats required for the completion of their life-cycles (Pinder, 1997). Thus, understanding lateral habitat function is crucial to the implementation of effective management actions and targeted rehabilitation projects. In Europe, lowland systems are dominated by cyprinid fishes but studies of the seasonal movements and habitat use in such environments are relatively rare.

The lower river Witham, in eastern England, is a uniform man-made channel that has been re-engineered many times since its inception during Roman occupation. Over the last two thousand years the lower river Witham has been managed for the purposes of navigation and land drainage, and has been straightened, widened and deepened to its current course (Wheeler, 1990). This environment presents fish populations with pressures such as the absence of a functional floodplain, large floodwater discharge and high velocity flows in the lower reaches (Environment Agency, 2008) leading to flush out (Linfield, 1985) and poor in-river and

marginal habitat. Intensive use of adjacent agricultural land results in high nutrient and silt levels from run-off leading to eutrophication and the potential to reduce water quality. River levels are artificially manipulated for drainage purposes resulting in fluctuating water levels that inhibit the establishment of permanent marginal habitat. In addition, two side-channels (Metheringham delph and Branston delph) have been made inaccessible to fishes and there is pressure to disconnect other side-channels as they are deemed unnecessary for land drainage yet require regular investment in order to maintain them. However, very little is known about the value of the side-channels to fish species, but anecdotal evidence would indicate that common bream use the side-channels for spawning, and common bream have been shown to utilise lateral habitats in European rivers (Molls, 1999).

Tracking is an established technique for monitoring and understanding fish behaviour (Lucas & Baras, 2001; Jellyman, 2009). Tracking studies have revealed homing behaviour by common bream (Langford *et al.*, 1979; Lyons & Lucas, 2002) and barbel (Lucas *et al.*, 1998a), as well as diurnal patterns of behaviour (Baade & Fredrich, 1998; Lyons & Lucas, 2002), annual spawning migrations (Fredrich *et al.*, 2003; Lucas & Batley, 1996), patterns of fish activity (Baade & Fredrich, 1998; Baras & Cherry, 1990) and the quantification of home range occupancy and site fidelity (Baras & Cherry, 1990; Winter & Fredrich, 2003).

Here the proposed study would determine the extent of seasonal activity patterns of common bream within the lower river Witham over an extended period, so repeated seasonal patterns of behaviour can be observed. In particular, the use of side channel habitat in relation to the time of year and the environmental variables of temperature and flow rate will be examined, in order to inform effective management of this lowland river system. It is hypothesised that the main river channel and the different tributary habitats would be used at different times of the year and in relation to changing environmental conditions.

Recently recreational fishery performance has deteriorated, indicated by match catch data (Environment Agency, unpublished data), and fishery survey results have been inconsistent, indicating reduced fish stocks. Water Framework Directive fishery classifications have indicated the lower river is not in favourable status, highlighting the need for rehabilitation. River rehabilitation schemes require data to target improvements; this study of common bream habitat use will provide such data. Also, the lower river Witham is subject to some of the most intensive river management practices employed in Europe, this study will present the opportunity to study how these practices affect fish behaviour and habitat use.

## Chapter 2 The ecology of the lower River Witham

### 2.1 Objectives


The overall objective of this chapter is to summarise the ecological status of the lower River Witham as a case study for heavily modified lowland rivers and to provide context for chapters 3-7. The specific objectives are to:

1. Provide an overview of the lower River Witham and its catchment area with regard to ecological and morphological status, human activities and influences, water quality and flow.
2. Assess the current status of the fish populations in the river in relation to human pressures and how these have changed over time.

### 2.2 The lower River Witham

#### 2.2.1 Background

The River Witham's source is located near South Witham ( $52^{\circ}45'44''\text{N}$ ;  $0^{\circ}37'38''\text{W}$ ), Lincolnshire UK, approximately 15 km south of Grantham. From here the river flows north towards Lincoln where it turns eastwards through the Lincoln Gap, an area between two raised pieces of land. The river then flows in a south-easterly direction across the lowland valley floor to the tidal outfall at Grand Sluice Boston ( $52^{\circ}58'53''\text{N}$ ;  $0^{\circ}1'46''\text{W}$ ), where it discharges into The Wash (Figure 2.1). In total the main river length is approximately 106 km. The total catchment area is 2010 km<sup>2</sup> and contains numerous watercourses.



**Figure 2.1** Map of the East Midlands and Lincolnshire. The lower River Witham flows from Lincoln to Boston and discharges into The Wash (source; Environment Agency).

The River Till is a major tributary joining the river at Lincoln, other major tributaries joining the river downstream of Lincoln include the Barlings Eau, Sincil Dyke, Duckpool Drain, Billingham Skirth, River Bain and River Slea. The lower river is classed as the reach between Lincoln and the tidal limit at Boston. This case study will focus on the lower river from Bardney lock to the tidal limit at Boston approximately 40 km of open continuous reach. The principal towns in the area are; Lincoln, Boston, Grantham, Sleaford, and Horncastle.

### 2.2.2 History

The city of Lincoln derives its name from Brayford Mere, the ancient British word for a deep pool being *Lynn*, hence Lincoln being the settlement by the deep pool (Wheeler, 1990). Other possible origins of the name Lincoln are similar, *Lindu* being ‘dark pool’ or *Lindun* being ‘fort on a hill by a pool’ both in the Brythonic language of the Celtic people. Lincoln became an important site during Roman times, it being at the junction between the Roman roads Ermine Street, between London and York, and Fosse Way, which led to Leicester and Bath. A Roman city, Lindum Colonia, and fort were built at Lincoln, one of only four in Britain, denoting its importance at that time.

Before Roman occupation the geography of the area was very different to what exists now. Traces of an Iron Age settlement, discovered in 1972, on the site of Lincoln by Brayford Pool, show that the site predates Roman involvement. The Romans conquered this part of Britain in AD 48 and set about changing the nature of the whole region with a series of massive engineering projects that would transform the area from swamp to fertile agricultural land and incorporating modern transport networks of roads and river navigations. William Henry Wheeler’s account from 1868, republished in 1990, gives the following description of the lie of the land and the extent of the Roman drainage works;

*“The Witham and Til discharged their water into the Lynn (Brayford Mere), and the swampy and low ground was frequently flooded by the overflowing of the Trent... The overflow from Brayford Mere would be either by a natural water course along the line of the Fosdyke, or through the gap in the high land below the high part of the city, into the meres on the south, where another great mere, extending from Washingborough to Chapel Hill, into which the Langworthy [Barlings Eau], the Bane [River Bain] and the Sleas [River Sleas] discharged their waters. The outlet for this mere was a winding tidal creek, extending through the marshes below Chapel Hill to the Scalp at Fishtoft, which has since become the channel of the Witham. In order to drain these meres and swampy grounds, the Romans either enlarged an old waterway, or cut the channel to the Trent, now known as the Fosdyke, the prefix of this name being the Roman word for an embanked cutting or ditch... For the drainage of the mere below Lincoln, the Cardyke... was cut, skirting the higher ground on the west side of the Black Sluice level, which was a continuation of the same mere and ran southwards to Thurlby and thence to the Welland and Nene... For drainage of Brayford Mere and of the swampy ground round Lincoln and to afford a better outlet for the Witham water, a straight cut was made by the Romans through the gap in the cliff below the city to Shortferry and thence, skirting the high land, to the*



*tidal creek at Chapel Hill. This cut now constitutes the channel of the river Witham. This catchwater drain intercepted the water of Langworthy and all the other brooks and water courses which formerly passed their water into the mere. The absence of all winding in the course of the channel between Lincoln and Chapel Hill, and the fact that portions of the fen lie on the east side, between it and the high land, indicate that this river was never a natural stream... There is no record of the name by which it was known during the Roman period. Since the Saxon times it has been known as the Witham. The word Witham is probably derived from Wye-om, or river plain”.*

Recent genetic analysis of roach has suggested that the Witham was once a tributary of the Trent (Environment Agency, unpublished data). These waterways are now linked by the Fosdyke canal, but data from the genetic study suggests that the Witham and Trent were linked prior to the construction of the Fosdyke. Wheeler (1990) suggests that perhaps the fosdyke was built along the course of a former watercourse that would have drained the upper Witham. Hence, it is likely the River Witham now flows to the east, whereas it used to flow to the west with the watershed to the east of Lincoln being separate.


Water drained from the upper river need not to pass through the river channel in the centre of Lincoln. Water can be diverted down the Sincil Dyke, which leaves the river about a mile upstream of the city centre before rejoining the main channel just below Bardney Lock. There is no record of when this drain was built, but it can only have been built for the purposes of diverting floodwater away from the city, and appears to be of either Roman or Norman origin (Wheeler, 1990).

The course of the lower River Witham has changed and been re-engineered many times over the preceding two thousand years, often as a result of Acts of Parliament with the objective of improving the navigation of the river or limiting flooding (Wheeler, 1990). Soon after the Roman works connecting Lincoln and the Wash, the Witham became one of the chief transport links with the continent, exporting corn and importing wine. During the reign of William the Conqueror, Lincoln was one of the most important cities in England, especially for the wool trade, with commerce on a parallel with that of London. Flooding events in 1014 led to alterations to the river, the terminal course was diverted from Bicker Haven in the Welland estuary to discharge into The Haven at Boston, giving rise to the growth of this port through the 12<sup>th</sup> and 13<sup>th</sup> centuries. However, with no maintaining authority the channel deteriorated due to siltation, so much so that navigation became difficult and a petition was sent to the King in 1342. In 1369 the centre for the wool trade was transferred to Boston due to the poor state of the river. Petitions of both the King(s) and Courts continued for many years until during the reign of Henry VII, approximately 1500, a commission decided that a sluice was needed to prevent sea water carrying sediment up the channel on the flooding tide and work began. This project was the first of a series of improvements and developments that allowed the continued draining of the fenland in the area and maintaining the navigation on the Witham (Wheeler, 1990).

The Act for the Improvement of the Navigation 1671 states that at this time the navigation was still in a state of decay, and a report by James Scribo in 1733 states that winter flooding

below Chapel Hill was commonplace and that without improvements it would not be long before the navigation would be entirely lost and draining of agricultural land also rendered impractical (Wheeler, 1990). To remedy the situation he proposed an improvement scheme that included a straightening in the channel from Tattershall through to Boston. In 1744 Mr John Grundy of Spalding prepared a scheme for restoring the navigation of the river between Boston and Lincoln, and for draining the lowlands. A year later a Mr Daniel Coppin proposed a similar scheme. But it was not until 1752 that meetings between landowners were held and a course of action decided on. The Witham Act 1762 created the Witham Drainage General Commissioners, who exercised jurisdiction over the drainage by the River Witham, its tributaries and main drains within the six districts of the Witham. The banks were embanked and the water drained off the lands by mechanical means so that they became suitable for cultivation. 1764 saw the start of the construction of the Grand Sluice at Boston, replacing the previous structure that had long since proved inadequate (Wheeler, 1990). This maintained the height of water above Boston to near high tide level; prior to this the tidal influence would have reached as far upstream as Chapel Hill. At this time the river had a far more natural appearance than the present. A plan made in 1762 by John Grundy shows the section between Boston and Tattershall (Figure 2.2) containing numerous meanders.

These bends were lost in accordance with John Grundy's proposals; the river was straightened, widened and deepened to its approximate current course. By 1830 the lower River Witham, as it is today, had taken shape. Further cycles of floods, reports and works occurred throughout the 1800s (Wheeler, 1990) and continue to the present day (Environment Agency,



**Figure 2.2** Plan of the lower River Witham in 1762 from Wheeler (1990).

2008). Evidence of the old river's course can still be seen today as anomalies visible on aerial photographs, especially around the Langrick Bridge area.

### 2.2.3 Catchment & land use

Lincolnshire is one of Britain's prime agricultural counties (Figure 2.3), with a long tradition of agriculture and horticulture, which remains strong today (Environment Agency, 2008). The draining of the fens has left rich fertile peat soils allowing the county to have up to six times more agricultural development than any other county in the UK with food production valued in the region of £300 million (NFU, 2008). Arable farming requires well drained soils. The peat soils, where the best agricultural land is located, retain water. To support agriculture on the peat a network of drainage ditches, pumping stations and raised watercourses have been constructed. This network maintains dry conditions in an area that would otherwise be very wet, thus facilitating farming (Environment Agency, 2008).

The Agricultural Land Classification (ALC) provides a measure for assessing the quality of farmland, grades are related to climate, topography, drainage, soil characteristics and other site factors, and the interactions between them. Grades 1 and 2 are the most versatile and productive land for agriculture, Grade 3 is considered moderate for agricultural production and Grades 4 and 5 are poor. Almost half (46%) of the agricultural land in the catchment can be classed as Grade 1 or 2 (this tends to be found in the south east of the catchment on the peat soils) and most of the remaining land (53%) classed as Grade 3. Therefore, the majority of the River Witham catchment area is used to produce arable crops, although historically only the free draining soils would have been suitable (Environment Agency, 2008).


### 2.2.4 Water Quality

Water quality samples have been taken by the Environment Agency and its predecessors on an approximately monthly basis at two sampling points, Tattershall bridge and Langrick bridge since 1980. To assess how water quality has changed in the lower river since 1990, data for dissolved oxygen, suspended solids, biological oxygen demand, total organic nitrogen and total phosphorus were analysed against date with linear regression. Results are presented in Table 2.1.

**Table 2.1** Results from linear regression analysis of Environment Agency water quality data.

Determinant	Sampling site	Period	Slope	F (df)	P	R <sup>2</sup>
Dissolved oxygen	Tattershall Bridge	1981-09	1.37	1.87 (1,329)	0.173	0.01
Suspended solids	Langrick Bridge	1989-09	-2.96	8.75 (1,357)	0.003*	0.02
Biological oxygen demand	Langrick Bridge	1990-09	-6.31	39.82 (1,345)	<0.0001*	0.10
Total organic nitrogen	Langrick Bridge	1990-09	-1.80	3.25 (1,351)	0.072	0.01
Total phosphorus	Langrick Bridge	1990-09	-5.89	34.64 (1,132)	<0.0001*	0.21

\* Significant at < 0.01 following Bonferroni correction.



**Figure 2.3** Map of land use within the River Witham Catchment Flood Management Plan (CFMP) area (Environment Agency, 2008).

Thus over the sampling periods there has been a significant reduction in suspended solids, biological oxygen demand and total phosphorus, while dissolved oxygen has increased but non-significantly, suggesting that water quality has improved.

### 2.2.5 Biological Quality

Four sampling points, three on the main river at Bardney, Tattershall and Langrick and one on the tributary Sincil Dyke have been sampled by kick sampling (Chadd, 2010) approximately

twice yearly (Spring and Autumn) by the Environment Agency and its predecessors since 1986-87.

Indices have been developed to summarise the taxa present in a macroinvertebrate sample. The use of freshwater macroinvertebrates as biological indicators is well established and a range of indices have been developed based on invertebrate community response to specific environmental pressures (Extence *et al.*, 2011). Average score per taxon (ASPT), number of taxa (NTAXA) and British monitoring working party (BMWP) scores were used to assess sensitivity to organic pollution in the river macroinvertebrate samples. Higher scores indicate a relatively higher proportion of the more pollution sensitive taxa present, usually indicating higher ecological status. This is due to taxa sensitive to organic enrichment being progressively lost resulting in lower overall diversity, but a higher abundance of tolerant animals, such as worms (Oligochaeta) and non-biting midges (Chironomidae). Temporal trends were analysed with linear regression and spatial differences with ANCOVA. Results from the linear regression analysis are presented in Table 2.2.

**Table 2.2** Results from linear regression analysis of macroinvertebrate indices (Environment Agency data) against date.


Sampling site	Index	Mean Score ( $\pm$ SE)	Period	Slope	F(df)	P	R <sup>2</sup>
Sincil Dyke	ASPT	4.1 (0.04)	1987-10	2.32	5.4 (1,45)	0.025	0.11
	NTAXA	18.6 (0.62)	1987-10	3.35	11.24 (1,45)	0.002*	0.20
	BMWP	76.8 (3.23)	1987-10	3.16	9.97 (1,45)	0.003*	0.18
Bardney	ASPT	4.1 (0.04)	1987-10	2.46	6.07 (1,49)	0.017	0.11
	NTAXA	20.6 (0.50)	1987-10	3.13	9.77 (1,49)	0.003*	0.17
	BMWP	86.88 (2.78)	1987-10	3.13	9.79 (1,49)	0.003*	0.17
Tattershall Bridge	ASPT	4.2 (0.04)	1986-10	1.75	3.07 (1,47)	0.086	0.06
	NTAXA	20.8 (0.51)	1986-10	0.16	0.03 (1,47)	0.873	0.00
	BMWP	89.6 (2.76)	1986-10	0.72	0.52 (1,47)	0.476	0.01
Langrick Bridge	ASPT	4.1 (0.04)	1987-10	1.98	3.91 (1,49)	0.054	0.08
	NTAXA	20.0 (0.44)	1987-10	1.50	2.26 (1,49)	0.139	0.05
	BMWP	83.8 (2.44)	1987-10	1.77	3.12 (1,49)	0.084	0.06

\* Significant at  $< 0.004$  following Bonferroni correction.

Spatially the scores for all the indices were lower (*i.e.* poorer water quality) in Sincil Dyke than the other three main river sites. ANCOVA with site as a fixed factor and year as a covariate revealed: ASPT (site  $F_{3,192} = 3.96$ ,  $P = 0.009$ , year  $F_{1,192} = 20.26$ ,  $P < 0.0001$ ), NTAXA (site  $F_{3,192} = 3.75$ ,  $P = 0.012$ , year  $F_{1,192} = 16.01$ ,  $P < 0.0001$ ) and BMWP (site  $F_{3,192} = 4.02$ ,  $P = 0.008$ , year  $F_{1,192} = 19.50$ ,  $P < 0.0001$ ). At both Bardney and Sincil Dyke there has been a significant improvement in the NTAXA and BMWP indices during the period 1987-2010 (Table 2.2).

### 2.2.6 Flow regime and flood events

The River Witham has a largely stable low flow regime interspersed by spate conditions (Figure 2.4), but is impacted by flow regulation and inter-catchment transfer. The naturalised flow



**Figure 2.4** Mean daily flow ( $\text{m}^3 \text{s}^{-1}$ ) of the Barlings Eau at Langworth bridge, this tributary generates the largest flow input into the lower river (Environment Agency data).

regime is typical of calcareous landscapes with spates, usually during the winter time, of short duration and linked to very heavy rainfall in the catchment.

Significant flooding problems are associated with the River Witham, its tributaries and the sea. The River Witham poses a direct flood risk to the three main urban areas within the catchment area; Grantham, Lincoln and Boston (Environment Agency, 2008). Comparing the major tributaries of the River Witham, the Barlings Eau generates the largest flow input into the lower river, followed by the River Bain and upper Witham. The Barlings Eau contribution reflects its relative large overall catchment area (Figure 2.5)

Peak flows from the Barlings Eau are managed through the use of the Branston Island flood storage basin, which is flooded during high flow events, reducing the peak flow entering the River Witham (Environment Agency, 2008) and thus reducing flood risk in the catchment below. Peak flows of the River Brant, River Till, and Upper Witham are also managed in a similar fashion, through the Lincoln Washlands (Environment Agency, 2008).

The tidal cycle influences flow conditions on the lower river. Grand Sluice closes every high tide to prevent sea water passing further upstream. This stops river flow from escaping into The Wash. The influence of the tidal cycle affects the shape of the hydrograph at Woodall Spa on the River Witham during high flow events. Tidal impacts on flows are routinely seen as far upstream as Bardney Lock, and have occasionally been observed in Lincoln. Because of the relatively flat nature of the area and the effect of tide-locking, significant flooding generally occurs from prolonged storms that last several days (Environment Agency, 2008).



**Figure 2.5** Langworth bridge gauging station on the Barlings Eau, a tributary of the lower river, during normal conditions (left) and the summer flooding (right) in June 2007 (authors own photos).

### **2.2.7 Water level management**

The river level below Bardney lock is artificially manipulated on a seasonal basis to increase winter flood storage. During the summer months the river level is maintained at approximately 1.5 m Ordinance Datum Newlyn (ODN), and from 1 November – 1 April the level is reduced to and maintained at 1 m ODN, during dry conditions (Figure 2.6). Similar water level manipulation also occurs on other rivers in the area including the lower River Welland, South Forty Foot Drain, lower Hobhole Drain, Stonebridge Drain, River Ancholme and rivers in South West England. This lowering of the water level exposes the marginal areas, and particularly the vegetation that could act as refuge for fish during periods of high flows.


### **2.2.8 Trent-Witham-Ancholme water transfer scheme**

Water is abstracted from the bottom of the Barlings Eau at Short Ferry for the Trent-Witham-Ancholme water transfer scheme. Water intakes have the potential to cause huge impacts on fish populations through entrainment of adults and fry (May & Van Rossum, 1995). The intake at short ferry is fitted with rotary 10 mm screens to prevent entrainment and impingement of adult fish. However this transfer is operated during the late spring and summer to augment flows in the heavily abstracted River Ancholme to the north and therefore has the potential to entrain free drifting larvae and fry.

### **2.2.9 Morphology**

Upstream of Grantham the River Witham has a relatively narrow, steep floodplain. Downstream of Grantham the floodplain of the River Witham continues to widen. There has been extensive widening, straightening and embanking of the River Witham, especially downstream of Lincoln. This has removed many of the natural geomorphological features, and has disconnected the river from its natural floodplain. The main channel is heavily modified (Figure 2.7 & 2.8) being trapezoidal and canalised with a width of 30-40 m. The riverbanks are embanked and reinforced with rock, as a result the river has no functional floodplain and the entire flow is restrained





**Figure 2.6** Mean daily river level of the lower River Witham at Kirkstead Bridge from 1999-2010 (Environment Agency data).



**Figure 2.7** The heavily modified lower River Witham looking downstream from Dogdyke (below the confluence with the River Slea), canalised and straightened with high floodbanks preventing connection with the floodplain (authors own photo taken at winter retention level).





**Figure 2.8** The heavily modified lower River Witham looking downstream from Tattershall Bridge, canalised and straightened with high floodbanks preventing connection with the floodplain (authors own photo taken at winter retention level).

within high flood banks (Environment Agency 2008) set back 3-4 metres from the water's edge on both sides (Forbes & Wheeler 1997).

The widening of the river channels has also caused sediment to be deposited within the channel. Widening the channel can reduce river flow velocities under normal flow conditions, which reduces the ability of the river to transport sediment. The deposition of sediment reduces the channel capacity, and leads to increased maintenance, such as dredging (Environment Agency, 2008).

## **2.2.10 Habitat Assessments.**

### **River Habitat Survey**

River Habitat Survey (RHS) is a method for assessing the physical character and quality of river habitats. It has been developed to help the conservation and restoration of wildlife habitats along rivers and their floodplains. Its main purpose is to provide river managers with information needed to sustain and enhance biodiversity, using catchment management plans and environmental impact assessment as two mechanisms for realising this objective (Raven *et al.*, 1998).

RHS is carried out along a standard 500m length of river channel. Observations are made every 50m at ten equally spaced spot-checks along the channel, whilst information on valley form and land-use in the river corridor provides additional context (Environment Agency,

2003). These data can be used to derive two indices of river corridor status. These are the Habitat Quality Assessment (HQA) and the Habitat Modification (HM) score. The HQA is a measure of the structural diversity of the river corridor and the score increases with improved habitat quality. By contrast, the modification score is a measure of the extent to which the natural character of the river has been modified; a score of zero indicates no significant modification and represents natural (good) conditions. HM scores are used to assign a Habitat Modification Class; 1 (Pristine/semi-natural) - 5 (Severely modified). Results of the RHS on the lower Witham are presented in Table 2.3. All suggest the lower River Witham is highly / severely modified.

**Table 2.3** Results from River Habitat Surveys (Environment Agency data).

Upstream – downstream order	Year Surveyed	Site Name	HM (Survey Scores)	HM Class (Survey Scores)	HQA (Survey Scores)	HQA Adjusted (Survey Scores)
1	1994	Stixwold	960	4	17	10
2	2005	Tattershall Br	4254	5	24	22
3	2007	Thorpe Tilney Road End	1416	5	14	14
4	1995	Kirkstead Bridge	3217	5	14	13
5	1994	Upstream Pelhams Land	2730	5	17	11
6	1996	Pelhams Land	1426	5	20	19
7	1996	Langrick Bridge	1200	4	10	10


### Aerial Photography

Aerial photographs (source: maps.google.co.uk; Figure 2.9 & 2.10) were examined for the lower River Witham to allow a visual assessment of the nature of river habitat throughout the whole length of the lower river. Analysis aimed to determine the degree of river meander along the study reach. The total number of degrees through which the river meanders per km was calculated, and differences between the upstream (Bardney Lock – Tattershall Bridge; ~21km) and downstream (Tattershall Bridge – Boston; ~20km) halves of the study reach were analysed with Mann-Whitney U test. The upstream half of the study reach was significantly more heterogeneous (Mann Whitney U = 582.5,  $n_1 = 21$ ,  $n_2 = 20$ ,  $P = 0.0002$ : mean upstream =  $76.0 \pm 10.3$ , downstream =  $26.0 \pm 8.93$ ). Thus the downstream half of the study area was significantly less meandering than the upper reach of the river.

### Depth Profile

During April 2010 the underwater topography along the main channel of the lower river was investigated by boat (Figure 2.11). The survey was carried out from Boston to Bardney lock (~40 km), and involved a single upstream transect in the centre of the main river channel of the survey reach with the boat kept at a constant ground speed of ~5 km h<sup>-1</sup>. A GPS echosounder (Ceeducer Pro™, CEE HydroSystems, Sydney, Australia) was used, data were interpreted and





**Figure 2.9** Aerial photograph of the lower River Witham, upstream half of the study area  
Bardney – Tattershall Bridge (maps.google.co.uk).



**Figure 2.10** Aerial photograph of the lower River Witham, downstream half of the study area  
Tattershall Bridge – Boston (maps.google.co.uk).



stored on a laptop computer using Hypack Hydrographic (Hypack inc., Connecticut, USA) survey software package. Data were collected at a resolution of ten readings per metre, and then sub-sampled to resolution of one reading per 10 metres for presentation. It was hypothesised that the river bed would be variable due to natural riverine processes of erosion and deposition, creating a varied underwater environment that would influence fish distribution.

The bed level of the lower river doesn't appear to have been shaped by riverine processes and is probably much the same as when it was built in the late eighteenth century. Man-made channels would have been clay lined to prevent leakage and it appears that this material has withstood many years of weathering. Water depth was at its greatest at the downstream end of the study reach at Boston being ~4 metres slowly shelving up to ~2 metres at Bardney.




Figure 2.11 A depth profile of the lower River Witham.

### 2.2.11 Barriers to fish migration

Due to its low lying nature, the lower River Witham has relatively few barriers to fish migration. The tidal limit at Grand Sluice, Boston represents a significant barrier with no mitigating fish passage solutions currently installed. However, from here the lower river is one continuous open reach for nearly 40 km until Bardney Lock / Short Ferry sluice, which is atypical for a navigable lowland river due to the especially low lying nature of the area which negates the need for navigation locks on the lower river (Environment Agency, 2008). In late 2010, fish passage facilities were installed at the Short Ferry sluice near the confluence with the Barlings Eau allowing fish in the lower river between Boston and Bardney to access the 13 km of river upstream between Bardney and Lincoln up to Stamp End lock which again represents a barrier to fish passage into the middle reaches.

### 2.2.12 River uses

The Witham catchment area has a relatively low human population density, although there are some areas with a high population concentration *e.g.* Lincoln. The land use is predominantly agricultural, mainly producing arable crops, which has potential impacts on the water course through nutrient and pesticide run-off and silt loadings. The principle recreational uses of the river are navigation, angling and local amenity.

### 2.2.13 Conservation

The Witham drains into 'The Wash' which is internationally protected as a Ramsar site, Special Protection Area (SPA) and Special Area of Conservation (SAC). Therefore, the river could potentially impact upon the habitats and communities associated with this site. There are currently twenty two Species Action Plans within the Lincolnshire Biodiversity Action Plan (BAP), eighteen of which are considered to be of National Priority and are included on the UKBAP. Within the twenty-two species BAPs, are two general species BAPs covering bats and farmland birds in the Lincolnshire county. The following Local BAP (LBAP) species are directly linked to the aquatic environment.

- White-clawed crayfish *Austropotamobius pallipes* (Le.)
- Otter
- Water vole *Arvicola terrestris* (L.)
- Compressed river mussel *Pseudanodonta complanata* (R.)
- Witham orb mussel *Sphaerium solidum* (N.)
- Grass-wrack pondweed *Potamogeton compressus* (L.)
- Greater Water parsnip *Sium latifolium* (L.)
- Ribbon-leaved water plantain *Alisma gramineum* (Lej.)
- Spined loach *Cobitis taenia* L.

Many of the causes of degradation to LBAP habitats and the decline of these species relate to changes to the water environment (Environment Agency, 2008).

## **2.2.14 Fishery assessments**

### **Background**


Fishery surveys have been carried out by the Environment Agency and its predecessors in line with a statutory duty to 'maintain, improve and develop' freshwater fisheries under the Salmon and Freshwater Fisheries Act (1975). The monitoring programme for the lower river has undergone many changes both temporally and spatially. Survey effort has been reduced in recent years. Fisheries survey data were available for 271 sites serviced by the Manby office, Northern area, Anglian Region, Environment Agency (Penczak *et al.*, 1991) but in 2009 this was reduced to just 47 quantitative sites (Environment Agency, unpublished data). Due to the inefficiencies of netting large lowland rivers, several sites from the lower river were dropped from the programme in 2007 and sampling by hydro-acoustics was increased to twice a year, once in the spring and repeated in the autumn.

### **Historic data**


Forbes and Wheeler (1997) compared historic fisheries survey data from 1788-1796 with data gathered by the Environment Agency from 1981-1991, showing a reduction in the sizes of pike, perch and European eel. Three species; burbot, Atlantic salmon and barbel were recorded in the late-eighteenth century that had subsequently become absent from the river and there is no evidence of re-colonisation by these species in subsequent years.

### **Seine netting surveys**


The lower river has been surveyed with seine netting (Hickley, 1996); stop nets were utilised at either end of the site to produce quantitative results. The wrap around seine netting (Coles *et al.*, 1985) survey technique was employed by The Environment Agency and its predecessors, two catches were performed at each site to give quantitative population estimates. All fish caught were measured and scales were removed from a representative sub-sample of roach and common bream for subsequent age determination (Bagenal, 1973) which was performed by the Environment Agency's Fisheries Laboratory at Bampton. Eight sites (seven main river, one tributary) were sampled prior to 2007, in 2007 four netting sites (one main river, three tributary) were sampled, in 2008 three netting sites (two main river, one tributary) were sampled and 2009 two netting sites (one main river, one tributary) were sampled. Catch depletion density and biomass population estimates (Carle & Strub, 1978) were produced for fish >99mm, below this length the results become more uncertain because of varying capture efficiency of smaller fish due to the net mesh size. Data are presented to illustrate the contribution made by each species to density (Figure 2.12) and biomass (Figure 2.13) and mean density (Figure 2.14) and biomass (Figure 2.15) population estimates (Carle & Strub, 1978) during the surveys 1994-2009. A Box plot of common bream lengths caught during each survey year is presented as Figure 2.16. No statistical analysis of these data was carried out due to the large errors associated with the results and the inconsistent spatial and temporal coverage, see discussion.




**Figure 2.12** The contribution made by each species to density population estimates for years 1994-2009.



**Figure 2.13** The contribution made by each species to biomass population estimates for years 1994-2009.




**Figure 2.14** A comparison of mean ( $\pm$  95% confidence intervals) fish density estimates for fish > 99 mm recorded during the surveys 1994-2009.



**Figure 2.15** A comparison of mean ( $\pm$  95% confidence intervals) fish biomass estimates for fish > 99 mm recorded during the surveys 1994-2009.





**Figure 2.16** Common bream lengths from seine netting surveys 1994-2009. Fork length refers to the length from the tip of the snout to the end of the middle caudal fin rays.

### Hydroacoustic surveys

Hydroacoustics offer a quantitative and cost-effective method of surveying fish populations within large lowland rivers (Duncan & Kubecka, 1993; Kubecka, 1996; Kubeka & Duncan, 1998; Lucas *et al.*, 1998a). Long lengths of river can be surveyed delivering stock assessment of the fish population at the community level (Lucas *et al.*, 2002). With much greater spatial coverage, information about fish distribution and densities is delivered with much greater resolution and resilience and more cost effectively than conventional seine netting techniques. Surveys were conducted by the author as part of the Environment Agency national fisheries monitoring programme. These occurred during the night, since fish are more active and evenly distributed throughout the water column during the hours of darkness and are therefore more easily surveyed with horizontal echo-sounding when compared to the daytime (Duncan & Kubeka, 1993; Kubeka, 1996; Kubeka & Duncan, 1998). Biannual surveys were carried out (April and October) to give data on fish distribution during both the spring and autumn seasons. Surveys during both the summer and winter seasons were prevented by heavy macrophyte growth and water level manipulation, which reduces river width and therefore sampling volumes, respectively. During 2007 and in April 2008 surveys were carried out from Southery to Boston (~30 km), then the survey reach was extended and surveys were carried out from Bardney lock to Boston (~40 km) for the October 2008 survey and both 2009 surveys. The results from the spring and autumn surveys carried out in 2008 are presented in the form of fish density maps as Figure 2.17 and 2.18. Each survey involved a single downstream (true right hand bank) and upstream (true left hand bank) transect of the survey reach with the boat kept at a constant ground speed of ~5 km h<sup>-1</sup>. These transects were carried out on adjacent nights and data averaged to a single density per 10 m section. Acoustic data were collected with a split

beam portable echosounder, HTI (Hydroacoustic Technology Inc., Seattle, Washington, USA) model 241 portable split-beam echo sounder, operating at a ping rate of 5 pings s<sup>-1</sup>, a pulse duration of 0.2 ms and a bandwidth of 200 kHz were used, and with a time varied gain of 40 Log *R*. Minimum target strength thresholds were set at -50 dB, with single echo targets filtered within 0.8 and 1.2 of the echo pulse length. The split beam transducer (4.5° x 8° beam angles) was mounted at the front of the boat at ~80 cm depth, as described by Duncan & Kubecka (1993) with the sonar beam directed horizontally and perpendicular to the flow to maximise coverage of the water column and sample fish in their side aspect (Duncan & Kubecka, 1993). Acoustic data were recorded onto a laptop computer using Echoscape software (HTI, Seattle, Washington, USA) and geo-referenced with a handheld GPS (Garmin, Kansas, USA). It was hypothesised that the fish community in this homogenous riverine environment would be contagious, favouring certain areas and more highly contagious during the autumn surveys as fish aggregate for over-wintering (see Lucas *et al.*, 1998a).


To analyse how fish were dispersed during each survey an index of dispersion (Fowler *et al.*, 1998) was calculated. There are three ways in which things (in this case fish) can be dispersed: regularly, randomly and contagiously (meaning aggregated or clumped). The index of dispersion is calculated by dividing the variance by the mean. Data with small variance will have a low dispersion index suggesting regular distribution, intermediate variance suggests random dispersion and data with large variance and thus a high dispersion index is indicative of contagious dispersion (Fowler *et al.*, 1998). To compare results across surveys, mean fish densities per km were calculated for the reach between Southery and Boston (~30 km), as this was sampled during all surveys. Dispersion index values for the three spring and three autumn surveys were analysed with Mann-Whitney U test. The effects of km downstream from Southery and season were analysed with two-way ANOVA. Data were also grouped into two; upstream and downstream of Tattershall Bridge, for the upstream and downstream halves of the study area and differences analysed with two-way ANOVA. Example fish density maps from the Autumn and Spring surveys in 2008 are presented in Figure 2.16 and 2.17. A summary of the results is presented in Table 2.4.

Dispersion indices for all surveys were highly contagious (Table 2.4). Autumn surveys were more contagious than spring surveys but not significantly (t-test,  $t = -0.57$ ,  $P = 0.609$ ,  $df = 3$ ). The effects of km and season on mean fish density per Km were tested with two-way ANOVA; there was a significant effect of km ( $F_{1,29} = 2.50$ ,  $P < 0.0001$ ), but not with season ( $F_{1,29} = 2.83$ ,  $P = 0.095$ ) or the interaction between km and season ( $F_{1,29} = 0.97$ ,  $P = 0.523$ ). The effects of


**Table 2.4** Summary of hydro-acoustic survey results.

	Spring 07	Autumn 07	Spring 08	Autumn 08	Spring 09	Autumn 09
df	310	322	328	358	362	362
Mean density	36.87	15.96	58.23	52.06	78.58	103.73
SD	79.52	22.43	47.90	88.82	79.89	153.36
Dispersion Index	171.50	31.52	39.41	151.55	81.22	226.74

## Lower Witham Hydroacoustic Fishery Survey Spring 2008



**Figure 2.17** Fish densities recorded during the spring 2008 hydro-acoustic survey.



**Figure 2.18** Fish densities recorded during the autumn 2008 hydro-acoustic survey.

location (upstream or downstream river section) and season on mean fish density per km were tested with two-way ANOVA; there was a significant effect of location (upstream or downstream river section);  $F_{1,29} = 43.04$ ,  $P < 0.0001$ ), but not season ( $F_{1,29} = 2.88$ ,  $P = 0.092$ ) however there was a significant interaction between location and season ( $F_{1,29} = 5.26$ ,  $P = 0.023$ ). Thus fish are not evenly distributed along the surveyed reach, but aggregated. Fish favoured the upstream half of the study area, especially so during the autumn survey window.



### 2.2.15 Water Framework Directive classification

The Water Framework Directive (WFD/2000/60/EC) established a framework for the protection of ground and surface waters to prevent deterioration, protect and enhance aquatic ecosystems, and to achieve 'good ecological status' for all waters by 2027. Water bodies are classified under various parameters, including fish, using monitoring data. Fish Classification Scheme 2 (FCS2) is used to calculate the fishery classification, the scheme uses a combination of species diversity and biomass of observed compared with expected values (Environment Agency, unpublished data). The current (2011 & 2012) classifications for the lower River Witham (Environment Agency, unpublished data) are presented in Table 2.5.

**Table 2.5** Water Framework Directive classifications for the lower River Witham (Environment Agency, unpublished data).

Element	2011 Classification	2012 Classification
Fish	Moderate	Moderate
Macroinvertebrate	Good	Good
Diatoms	Moderate	Moderate
Macrophytes	Moderate	Moderate
Phosphate	Moderate	Moderate
Dissolved Oxygen	Good	Good
Ammonia	High	High
pH	High	High
Hydrology	Good	Good
<b>Overall status</b>	<b>Moderate</b>	<b>Moderate</b>

## 2.3 Discussion

### 2.3.1 Water Quality

Oxygen levels appear to have increased slightly over the last 30 years this may be a result of wastewater treatment investment or increased macrophyte growth and associated increase in photosynthesis. All these samples would have been taken during the day therefore night-time oxygen sags would not have been detected. Suspended solids have reduced in recent years which will promote macrophyte growth. Biological Oxygen Demand has reduced, illustrating the improvement in water quality since investment to upgrade sewage treatment works (STW) in the late 1980s. Peak Total Organic Nitrogen has also reduced in the lower river showing how recent agricultural good practice measures have reduced levels of nitrate and nitrite in run-off. However, the baseline minimum levels in this surface water have increased, possibly due to elevated levels in groundwater. Total phosphorus has also reduced, possibly as a consequence of phosphate stripping at large sewage treatment works (over 10,000 population equivalents) around 2004/5 (DEFRA, 2012). However, targets as low as 0.025 mg/L total phosphorus have been recommended by the United States Environmental Protection Agency (1986). In general these results show that organic pollution of the lower river has reduced and water clarity has increased. However the river is highly eutrophic due to nutrients from agricultural run-off from surrounding land which is dominated by large-scale arable agriculture.

### **2.3.2 Biological quality**

Temporally there is a general trend towards improvement on all indices, reflecting investment in wastewater treatment and the national trend for improved water quality brought about as a result. The Sincil Dyke scores were slightly lower than the main river; this may be due to wastewater discharges impacting on this low flowing tributary which lacks the dilution offered by the main channel. Also being low flowing the Sincil Dyke tends to have a more silt-based substrate which may favour macroinvertebrates that are associated with low water quality, such as worm (*Oligochaeta*) and non-biting midges (*Chironomidae*) (Chesters, 1980).

### **2.3.3 Habitat**

Historical accounts of how the lower Witham has been modified reveal that the lower river has been totally reshaped and transformed from a once tidal creek surrounded by transitional ecotones to a heavily managed, impounded, linear channel totally disconnected from its floodplain. The floodplain is an essential habitat component in lowland rivers (Aarts *et al.*, 2004) and the diminished hydrological connectivity between the river and its floodplain reduces the opportunity for fish, especially juveniles, to escape the force of elevated flows (Copp, 1997; Nunn *et al.*, 2007).

The Habitat Modification (HM) scores for the surveys carried out between 1994-2007 on the lower River Witham range from 960-4254, two surveys were HM class 4 (significantly modified) and the remaining five surveys were all in the highest class 5 (severely modified), illustrating the level of human intervention that has taken place on the lower river. The Habitat Quality Assessment (HQA) scores are relatively low illustrating the low diversity of natural river bank and in-channel features in the survey reach. Tributaries add some habitat diversity to this homogenous riverine environment, two side-channels were made inaccessible to fishes, Metheringham Delph in the 1990s and Branston Delph in the 1960s (Dave Fisher, Environment Agency, pers. comm.) and there is pressure to remove further tributaries (Nocton and Timberland Delphs) as they are deemed unnecessary for land drainage yet require regular investment in order to maintain them.

The depth profile results indicate that the river bed is also homogeneous with little variation. Aerial photograph analysis show how the river below Tattershall bridge is particularly homogeneous being highly straightened and lacking tributaries below the River Sleas confluence, offering fish no refuge during high flows.

### **2.4.4 Fishery assessments**

The historical data examined by Forbes and Wheeler (1997) contains reference to three species recorded in the lower Witham in the late-eighteenth century (1788-1796; prior to the river being straightened to its current course) that have subsequently become extinct from the river. All three of these species (barbel, salmon and burbot) have specialist habitat and migratory requirements (Wheeler, 1978). During the eighteenth and nineteenth centuries burbot were

described as being 'very abundant' in the Witham and other fenland rivers but numbers reduced following channelization and the species has not been reliably recorded in the UK since 1969 (Worthington *et al.*, 2011). Both salmon and barbel rely on high water quality, lotic conditions, upstream spawning migratory pathways and gravel spawning substrate. All of these requirements would have been seriously compromised by the level of river modification that has taken place since.

The lower river currently contains a typical lowland fish community, being dominated by roach and common bream but also populated by pike, perch, tench, silver bream and the European eel. Catch efficiency varied between seine netting sites and between years, with heavy weed growth producing population estimates (Carle & Strub, 1978) with large sampling errors and results with large variances. Catches of common bream in different years often varied in size and age composition as surveys captured different year classes. In some years when large numbers of juveniles were caught this could be interpreted as successful recruitment (*e.g.* 2006). However large catches of adults (*e.g.* 2002 and 2008) are likely to be an artefact of the relatively low sampling effort, as these fish were present in the river in previous years, given that adult common bream were aged between 10-15 years. A very large catch of juveniles occurred in 2007 from the entrance to the Sincil Dyke at Bardney; river flows had been very high in the main river over the summer and immediately before the survey was carried out and it appeared that very large numbers of fish were taking refuge in this slower flowing tributary. Roach x common bream hybrids were common in all survey years. Hybridisation can be a symptom of reduced lateral and longitudinal connectivity which leads to an overlap of spawning habitat (Cowx, 1983).

The hydroacoustics data shows that the fish community is highly aggregated and varies along the studied reach. Fish favour the upstream half of the study area during the cooler autumn surveys. Fish form large groups as a method of protection in clear water conditions (Jacobson *et al.*, 2004) leaving large areas periodically devoid of fish. There are consistent low densities of fish in the lower half of the study reach; this section consists of more homogenous habitat as shown by the analysis of aerial photographs. High flow rates in the lower reaches and a distinct lack of shelter habitat can cause a net annual loss through flush out (Linfield, 1985; Copp, 1997). Hydroacoustic techniques are effective at monitoring fish populations in homogenous riverine environments (Duncan & Kubecka, 1993; Kubecka, 1996; Lucas *et al.*, 1998a). However, this technique fails to differentiate to the species level and at present there is no Water Framework Directive classification scheme for data derived from this technique.

Currently the lower river is dominated by roach. Eurytopic species, like roach and perch, have less strict requirements for spawning habitats and are thus able to establish populations in a wide range of waterbodies (Schiemer & Waidbacher, 1992; Nunn *et al.*, 2007) leading to their dominance in modified habitats (Wolter, 2001a). The dominance of roach and perch is common in modified lowland rivers in Europe (Wolter & Vilcinskis, 1997; Wolter, 2001a; Wolter 2001b). The structure of the fish population reflects the overall quality of their environment, in particular the relative abundances of specifically adapted floodplain species (*e.g.* common

bream) can indicate the quality and extent of lateral connectivity (Schiemer, 2000). In the lower River Witham, common bream are present and when captured during surveys dominate the rivers biomass. Their existence may be due to available lateral spawning habitats that provide surrogates for and are functionally similar to lentic floodplain waterbodies of natural riverine ecosystems.

Recent changes in river character following water quality improvements, gained by investment in wastewater treatment, include increased water clarity and consequent heavy macrophyte growth fuelled by eutrophication. These conditions have made traditional seine netting survey techniques (Coles *et al.*, 1985) less effective as heavy weed growth upsets the level bed profile required for successful seine netting (Hickley, 1996) and leads to poor catch depletions, producing inaccurate population estimates (Carle & Strub, 1978). Increased water clarity has been shown to affect roach distribution through increased aggregation (Jacobsen *et al.*, 2004) and aggregated fish distributions mean that the variance of the total population size will be so large that estimates based on small numbers of samples will be unreliable (Kell, 1991). These factors have made temporal and spatial monitoring of fish communities in large open systems, like the lower river Witham, by traditional techniques less effective.



## **Chapter 3 The effects of surgical implantation of telemetry transmitters and translocation on the spatio-temporal behaviour of common bream.**

### **3.1 Introduction**

The development of electronic tags has been one of the most important advances for studying the temporal behaviour, movement and distribution of fishes in the freshwater environment (Lucas & Baras, 2001). Implantation into the peritoneal cavity is recommended for long-term tracking studies of fishes (Lucas & Batley, 1996) and is regularly used for cyprinid species (*e.g.* Lucas & Batley, 1996; Huber & Kirchhofer, 1998; Lyons & Lucas, 2002; Winter & Fredrich, 2003; Fredrich *et al.*, 2003; Kuliskova *et al.*, 2009). Telemetry studies are most often designed with three assumptions: (1) fish condition and mortality are not altered by transmitter presence or handling stress; (2) transmitters are retained for the duration of the observation period; and (3) tagged individuals accurately represent the population being observed (Smith *et al.*, 1998; Ramstad & Woody, 2003; Neely & Steffenson, 2010). This chapter describes the protocol used for the surgical implantation of an acoustic transmitter into the peritoneal cavity of common bream in the field. Similar techniques have been used successfully in previous studies (*e.g.* Lyons & Lucas, 2002), but a detailed description of the procedure is absent from the literature. Eight fish were recaptured at various times post-surgery and the incision site photographed to determine the state of tissue regeneration.

Meaningful data can only be collected during telemetry studies when the tagged fish are behaving naturally, thus the effects of tag attachment should not impact on the behaviour or well-being of the fish. However, the potential effects of these transmitters on fish performance is unknown (Jepsen *et al.*, 2003). Assessment of the effects of surgical implantation of transmitters on tagged fish has been carried out previously by observation for a period of days in captivity (Baras, 1997; Allouche *et al.*, 1999) and examination after recapture (Lucas & Batley, 1996) to determine the degree of healing, mortality and effects on fish weight and condition. Assessment of stress levels following implantation have also been carried out by measuring cortisol (a steroid hormone released in response to stress; Ellis *et al.*, 2004) levels released into water by individual common carp and roach tagged with dummy transmitters and held in tanks. Cortisol levels were elevated during the 12 hours post-surgery, but there was no evident long-term cortisol stress response to the presence of a tag in the body cavity in either species (Lower *et al.*, 2005). However, there is a paucity of data on behavioural effects immediately after procedures have been carried out and unusual long-distance movement have been observed during the first days of release after transmitter implantation in some species *e.g.* largemouth bass *Micropterus salmoides* (Lac.) (Mesing & Wicker, 1986) and dace (Clough & Beaumont, 1998). In my studies, long-term telemetry presented the opportunity to compare the behaviour of common bream that had just been tagged with common bream previously tagged (between 55 days and 378 days post surgery) and present in the same water-body, and thus subjected to the same environmental variables, over ten days post surgery. It was hypothesised that freshly tagged fish

would move more or less as far and occupy lengths of river longer or shorter than fish that had been tagged previously.

Studies of non-migratory fishes have revealed homing behaviour, defined as '*the return to a place formerly occupied instead of going to equally probable places*' (Gerking, 1959), following translocation in common bream (Langford *et al.*, 1979) and displacement by high flow events in barbel (Lucas *et al.*, 1998a). This indicates that these fish exhibit site fidelity. Site fidelity has been observed in common bream; Vostradovsky (1975) marked 3460 common bream between 181-260 mm in Lipno reservoir (4650 hectares) Czech Republic, over a period of five years and these were recaptured by recreational and commercial fisherman. Eighty eight % of recaptures (although the total recapture rate was only 10%) were from within 1 km of the original capture site. Homing behaviour following translocation has also been observed; Langford *et al.* (1979) found that displaced acoustic tagged indigenous common bream in the River Witham returned to their capture point, although this was only observed during the winter. Also, Goldspink (1978) translocated 4619 opercular tagged (marked) common bream from Zwartemeer to Tjeukemeer, two large lakes connected by canals in The Netherlands. The fish dispersed and were recaptured up to 60 km from the release site. It seemed that the fish were homing back towards Zwartemeer via the canal system. In this study three common bream that were being tracked were recaptured and translocated ~35 km downstream and their behaviour compared with other individuals that were being tracked but were not recaptured or translocated. My intention was to ascertain whether there was any homing behaviour and if there were other behavioural differences between the two groups that were present in the same water-body and so subject to the same environmental variables. It was hypothesised that translocated fish would move longer distances and over longer lengths of river than non-translocated fish as they homed back to the area where they were captured.

Here, the spatio-temporal movements of fish measured in terms of both Linear Range (LR) and Total Distance Moved (TDM) were compared. Behaviour of fish that had been translocated and freshly tagged were compared with fish that had not been translocated or freshly tagged, but were present in the same waterbody during the same period to determine the behavioural effects of surgery and translocation.

## **3.2 Material and methods**

### **3.2.1 Study area**


The study area (Figure 3.1) is a continuous open reach of approximately 40 km of the non-tidal lower River Witham and associated tributaries starting upstream near Short Ferry (53°13'38"N; 0°21'23"W) where the Barlings Eau tributary enters the old river loop and ending downstream at the tidal limit, Grand Sluice in Boston (52°58'53"N; 0°1'46"W).

### **3.2.2 Sampling procedures**


In total, eighty-three indigenous adult common bream (mass: mean 2.37 kg  $\pm$  SD 0.25, range 1.93-2.95 kg; fork length: mean 485.6 mm  $\pm$  SD 16.8, range 440-522 mm; age determined by

scale analysis: 13.62 mean  $\pm$  SD 1.97, range 10-19 years; gender: 48% male  $n = 40$ , 52% female  $n = 43$ ; details of all tagged fish are given in Table 3.1) were caught from the River Witham, Lincolnshire, UK, and tagged in seven batches between November 2006 and February 2009. A frequency histogram for the length of the tagged common bream is presented as Figure 3.2, the distribution illustrates that the sample were all large adults.

Common bream were caught in seven groups by electro-fishing (240V, 4A, pulsed DC), wrap around seine netting (Coles *et al.*, 1985; Figure 3.3) and by rod and line from the main river at Bardney (53°12'27"N; 0°20'12"W) and Tattershall Bridge (53°5'26"N; 0°12'54"W) and from the tributaries Billingham Skirth at Tattershall Bridge and Sincil Dyke at Bardney between November 2006 and February 2009. Details of all fish tagged are presented in Table 3.1. All fish were retained overnight; on one occasion in net cages placed in the river (site was secure), or more commonly in holding tanks between capture and tagging. After surgery fish were released at the site of capture. No tracking took place during July-September 2008, as existing tags had expired and tagging new fish was delayed until water temperatures reduced during the autumn / winter months due to the benefits to fish recovery at higher dissolved oxygen levels.



**Figure 3.1** Map of the lower River Witham between Lincoln and Boston, the study area runs from the old river loop upstream of Bardney to the tidal limit at Boston (source: Environment Agency).



**Figure 3.2** Frequency distribution of the length of the tagged common bream.



**Figure 3.3** Seine net catch of 990 common bream caught in a single 100 m haul on 3<sup>rd</sup> December 2007 downstream of Bardney Bridge. Mobile tracking revealed the presence of several previously tagged fish in the vicinity which enabled the area to be targeted with seine netting (authors own photo).

**Table 3.1** Details of common bream tagged and tracked during the study. Individual tag number, tag type, capture method, date tagged, capture site, mass (kg), fork length (mm), age determined by scale analysis, gender, nominal tag delays, tag life, length of dataset collected (days), projected tag expiration date and whether the tag automatically switched itself off (y) or ran until batteries ran flat (n).

No.	Tag No.	Tag Type	Capture Method	Date Tagged	Capture / Release Site	Mass (kg)	Fork Length (mm)	Age (years) <sup>5</sup>	Male / Female	Average Tag Delay (s)	Stated Tag Life (days)	Length of Dataset (days)	Projected tag end point	Auto Shut-off
1	2	V9	Electro fishing <sup>1</sup>	21/11/2006	Bardney Lock <sup>3</sup>	2.268	473	16	F	20	135	141	05/04/2007	n
2	241	V9	Electro fishing <sup>1</sup>	21/11/2006	Bardney Lock <sup>3</sup>	2.835	510	10	M	60	330	501	17/10/2007	n
3	242	V9	Electro fishing <sup>1</sup>	21/11/2006	Bardney Lock <sup>3</sup>	2.155	465	11	F	60	330	130	17/10/2007	n
4	243	V9	Electro fishing <sup>1</sup>	21/11/2006	Bardney Lock <sup>3</sup>	1.928	440	10	M	60	330	108	17/10/2007	n
5	247	V9	Electro fishing <sup>1</sup>	21/11/2006	Bardney Lock <sup>3</sup>	2.948	512	-	M	40	240	255	19/07/2007	n
6	248	V9	Electro fishing <sup>1</sup>	21/11/2006	Bardney Lock <sup>3</sup>	2.608	485	10	M	40	240	233	19/07/2007	n
7	249	V9	Electro fishing <sup>1</sup>	21/11/2006	Bardney Lock <sup>3</sup>	2.495	475	12	F	40	240	163	19/07/2007	n
8	1	V9	Rod and Line	28/02/2007	Tattershall Bridge <sup>4</sup>	2.098	460	10	F	10	80	111	19/05/2007	n
9	244	V9	Rod and Line	28/02/2007	Tattershall Bridge <sup>4</sup>	2.381	484	13	M	60	330	495	24/01/2008	n
10	245	V9	Rod and Line	28/02/2007	Tattershall Bridge <sup>4</sup>	2.438	510	13	F	60	330	69	24/01/2008	n
11	246	V9	Rod and Line	28/02/2007	Tattershall Bridge <sup>4</sup>	2.098	480	10	M	60	330	489	24/01/2008	n
12	250	V9	Rod and Line	28/02/2007	Tattershall Bridge <sup>4</sup>	2.381	492	10	F	40	240	400	26/10/2007	n
13	251	V9	Rod and Line	28/02/2007	Tattershall Bridge <sup>4</sup>	2.495	505	-	F	40	240	399	26/10/2007	n
14	252	V9	Rod and Line	28/02/2007	Tattershall Bridge <sup>4</sup>	2.268	465	15	F	40	240	400	26/10/2007	n
15	6066	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	2.041	476	13	F	40	210	208	07/05/2008	y
16	6067	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	2.069	461	12	M	40	210	208	07/05/2008	y
17	6068	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	2.523	508	14	M	40	210	210	07/05/2008	y
18	6069	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	2.608	496	13	F	40	210	210	07/05/2008	y
19	6070	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	2.155	472	12	F	40	210	209	07/05/2008	y
20	6071	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	2.495	508	15	F	40	210	210	07/05/2008	y

No.	Tag No.	Tag Type	Capture Method	Date Tagged	Capture / Release Site	Mass (kg)	Fork Length (mm)	Age (years)	Male / Female	Average Tag Delay (s)	Stated Tag Life (day's)	Length of Dataset (day's)	Projected tag end point	Auto Shut-off
21	6072	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	2.325	496	13	F	40	210	210	07/05/2008	y
22	6073	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	2.665	520	15	F	40	210	208	07/05/2008	y
23	6074	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	2.381	500	16	F	40	210	208	07/05/2008	y
24	6075	V9	Rod and Line	10/10/2007	Bardney Bridge <sup>3</sup>	1.928	468	14	M	40	210	210	07/05/2008	y
25	6076	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.863	487	15	F	40	210	184	01/07/2008	y
26	6077	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.183	472	13	M	40	210	203	01/07/2008	y
27	6078	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.722	506	13	F	40	210	209	01/07/2008	y
28	6079	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	1.956	458	15	M	40	210	210	01/07/2008	y
29	6080	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.183	471	12	M	40	210	208	01/07/2008	y
30	6081	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.296	492	12	F	40	210	179	01/07/2008	y
31	6082	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.636	480	12	F	40	210	183	01/07/2008	y
32	6084	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.608	506	13	F	40	210	210	01/07/2008	y
33	6085	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.353	495	11	M	40	210	202	01/07/2008	y
34	6086	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.495	496	15	M	40	210	160	01/07/2008	y
35	6087	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.381	490	13	F	40	210	210	01/07/2008	y
36	6088	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.410	495	13	F	40	210	210	01/07/2008	y
37	6089	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.098	474	12	M	40	210	209	01/07/2008	y
38	6090	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.920	511	14	M	40	210	176	01/07/2008	y
39	6091	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.155	472	13	F	40	210	191	01/07/2008	y
40	6092	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.438	502	14	M	40	210	153	01/07/2008	y
41	6093	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.523	475	14	M	40	210	203	01/07/2008	y
42	6094	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.069	475	11	M	40	210	210	01/07/2008	y
43	6095	V9	Seine netting <sup>2</sup>	4/12/2007	Bardney Bridge <sup>3</sup>	2.438	468	11	F	40	210	202	01/07/2008	y

No.	Tag No.	Tag Type	Capture Method	Date Tagged	Capture / Release Site	Mass (kg)	Fork Length (mm)	Age (years)	Male / Female	Average Tag Delay (s)	Stated Tag Life (day's)	Length of Dataset (day's)	Projected tag end point	Auto Shut-off
44	12247	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.835	480	14	F	50	526	224	11/03/2010	y
45	12248	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.296	487	14	M	50	526	360	11/03/2010	y
46	12249	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.438	505	16	M	50	526	299	11/03/2010	y
47	12250	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.268	470	17	M	50	526	535	11/03/2010	y
48	12251	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.665	498	12	F	50	526	534	11/03/2010	y
49	12252	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.608	482	12	F	50	526	259	11/03/2010	y
50	12253	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.551	486	14	M	50	526	531	11/03/2010	y
51	12267	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.551	488	14	M	60	621	571	14/06/2010	y
52	12268	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.438	478	15	F	60	621	278	14/06/2010	y
53	12269	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	1.984	483	13	F	60	621	629	14/06/2010	y
54	12270	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.665	468	13	F	60	621	244	14/06/2010	y
55	12271	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.325	486	15	M	60	621	306	14/06/2010	y
56	12272	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.778	495	13	F	60	621	202	14/06/2010	y
57	12273	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.325	494	15	M	60	621	589	14/06/2010	y
58	12274	V13	Electro fishing <sup>1</sup>	1/10/2008	Billinghay Skirth <sup>4</sup>	2.551	495	11	M	60	621	599	14/06/2010	y
59	12254	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.410	478	13	M	50	526	158	20/05/2010	y
60	12255	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.041	476	12	M	50	526	528	20/05/2010	y
61	12256	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.325	478	18	F	50	526	165	20/05/2010	y
62	12257	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.495	488	17	F	50	526	535	20/05/2010	y
63	12258	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.325	496	14	F	50	526	235	20/05/2010	y
64	12259	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.041	489	14	F	50	526	184	20/05/2010	y
65	12275	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.155	468	18	M	60	621	132	23/08/2010	y
66	12276	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.155	479	14	M	60	621	187	23/08/2010	y

No.	Tag No.	Tag Type	Capture Method	Date Tagged	Capture / Release Site	Mass (kg)	Fork Length (mm)	Age (years)	Male / Female	Average Tag Delay (s)	Stated Tag Life (day's)	Length of Dataset (day's)	Projected tag end point	Auto Shut-off
67	12277	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.835	522	14	F	60	621	253	23/08/2010	y
68	12278	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.155	478	14	F	60	621	167	23/08/2010	y
69	12279	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	2.098	472	15	M	60	621	174	23/08/2010	y
70	12280	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	1.984	460	15	M	60	621	171	23/08/2010	y
71	12281	V13	Electro fishing <sup>1</sup>	10/12/2008	Sincil Dyke <sup>3</sup>	1.928	454	14	M	60	621	543	23/08/2010	y
72	12260	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.098	488	15	M	50	526	410	05/08/2010	y
73	12261	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.211	482	13	M	50	526	143	05/08/2010	y
74	12262	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.438	510	14	M	50	526	130	05/08/2010	y
75	12263	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.580	518	16	F	50	526	136	05/08/2010	y
76	12264	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.551	501	19	F	50	526	40	05/08/2010	y
77	12265	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.098	466	15	M	50	526	134	05/08/2010	y
78	12266	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.722	506	18	M	50	526	534	05/08/2010	y
79	12282	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.069	492	14	M	60	621	338	08/11/2010	y
80	12283	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.495	490	14	F	60	621	169	08/11/2010	y
81	12284	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.041	452	13	F	60	621	521	08/11/2010	y
82	12285	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.325	487	14	F	60	621	82	08/11/2010	y
83	12286	V13	Electro fishing <sup>1</sup>	25/02/2009	Sincil Dyke <sup>3</sup>	2.551	502	16	F	60	621	155	08/11/2010	y

<sup>1</sup> Carried out using 240V, 4A, pulsed DC.

<sup>4</sup> 53°5'26"N; 0°12'54"W.

<sup>2</sup> Using the wrap around technique (Coles *et al.*, 1985).

<sup>5</sup> Determined by scale analysis (Bagenal, 1973).

<sup>3</sup> 53°12'27"N; 0°20'12"W.



Two types of tags were used as manufacturing specification and tag life expectancy changed as the study progressed. For the first part of the study (November 2006 - June 2008) Vemco (Nova Scotia, Canada) V9-2L coded pingers which were cylindrical with dimensions of 29 mm by 9 mm diameter, weight in air of 4.7 g, weight in water 2.9 g and with operational life of 80-330 days were used. For the second part of the study Vemco V13-1L coded pingers with dimensions of 36 mm by 13 mm diameter, weight in air of 11.0 g, weight in water 6.0 g and with operational life of 526-621 days were implanted into the body cavity. The tag weight in air would represent 0.16-0.57% of the fish's weight out of water, well below the 2% recommended by Winter (1983). The first fourteen V9 tags used in the winter 2006/2007 had no automatic shutdown, meaning they transmitted until their internal batteries ran out. From then on all tags had an automatic shut down, meaning they ran for a specific period then switched off. This was due to manufacturer specification changing during the study.

### **3.2.3 Tag implantation protocol**

The procedure described is regulated and licensed in the UK by the Home Office under the Animals (Scientific Procedures) Act 1986, project licence number PPL 80/2016. The surgery itself took place within the shelter of a mobile laboratory under conditions that were as aseptic as possible in-the-field. A specially constructed operating table was used to secure and restrain the fish in an upside-down position with the incision site clearly accessible. The mobile laboratory and operating table were washed and disinfected before any procedures were started. Surgical equipment was sterilised in surgical spirit and then rinsed with sterile water or saline solution. Prior to the procedure the activated tag was tested in water with a hydrophone. Tags were then sterilised with a 25% Dettol™ solution, rinsed with sterile water or saline solution and stored in a sterile swab.

A 60 litre anaesthetic bath of Tricaine methanesulphonate MS-222 at 50 mg/L of river water (Carter *et al.*, 2011), and a similar sized recovery tank of river water, both with aeration were clearly labelled, and the pH and dissolved oxygen levels monitored (Figure 3.4). The anaesthetic bath was tested with a single fish (which was not used again for subsequent anaesthesia and tagging) until anaesthesia was reached as indicated by loss of the righting reflex and a slowed operculum rate, which did not stop. The test fish was allowed to fully recover before any fish to be tagged were anaesthetised.

All surgeries were conducted by myself (Home Office personal licence number PIL 80/9908) with an assistant. Hands were disinfected with an alcohol based hand wash and maintained as aseptic a procedure as was possible under field conditions. I wore sterile surgical gloves, changing them between fish, or after coming into contact with anything away from the surgical area.

Fish were firstly weighed, measured, their sex determined, primarily by the presence of tubercles and also by vent & body morphology (Kennedy & Fitzmaurice, 1968), and a scale sampled, with a scalpel, from the mid-flank above the lateral line, for subsequent age determination (Bagenal, 1973). The fish was then placed into the aerated anaesthetic bath. When

anaesthesia was reached the fish was removed from the bath, inverted and secured in the operating table between wet smooth foam padding with Velcro™ straps and transferred to the mobile laboratory. During the surgery an assistant monitored the operculum movement throughout. If a problem with the fish's health or well being was encountered during the procedure then the team had the option to administer an overdose of anaesthetic in-line with schedule one of the Animals (Scientific Procedures) Act 1986.

The incision site was on the ventral surface of the fish, anterior of the pelvic fins and associated muscle blocks. To allow access, a line of 4 to 6 scales were removed, with tweezers, from the incision site (Figure 3.5), another scale was removed midway and perpendicular to the incision site to allow suture entry. The site was then swabbed with an iodine based antiseptic (Betadine™). An incision was made with a sterile scalpel and was kept to the minimum size required, approximately 20-30 mm in length (Figure 3.6). The incision was begun just in front of the pelvic muscle blocks, where the body wall thins, and extended towards the pectoral fins.

The incision was made slowly by dragging the scalpel lightly. An assistant used sterile tweezers to hold the incision open while it was being cut, this allowed the surgeon to see exactly when the incision breached the body cavity and thus avoided making an incision that was too deep resulting in damage to vulnerable internal organs. Once the incision was of sufficient size, the sterile tag was inserted using a sterile gloved hand and/or sterile tweezers (Figure 3.7). The incision was closed with a single suture (Ethicon PDS\*II Polydioxanone violet monofilament absorbable W9125; Ethicon, Piscataway, NJ, U.S.A.; Figure 3.8). One scale had already been removed to allow suture entry and another 1-2 scales were removed with tweezers to achieve suture exit. The surgeon and an assistant worked together to achieve this. The suture was secured with a surgeon's knot and excess suture material was trimmed with sterile scissors 5-10 mm from the knot.

The site was swabbed again and G7 wound sealer (Lincolnshire Fish Health, UK) was applied and allowed a few seconds to, at least partly, dry. The site was then liberally covered with Orabase™ (Squibb & Sons, Uxbridge, UK) protective paste to provide a temporary barrier (Figure 3.9), G7 wound sealer was reapplied on top of the Orabase™, to help prevent the Orabase™ barrier dissolving when fish were returned to the water. The whole procedure took three to four minutes. The fish was then removed from the operating table and immediately placed in the aerated recovery tank (Figure 3.10) and supported by hand in an upright position. An antibiotic injection of 1 ml Baytril at 2.5% directly behind the dorsal fin was given while the fish was recovering, although antibiotic use post surgery has subsequently been questioned by Mulcahy (2011). The injection was made on the same line as the erect last dorsal ray between the two main muscle blocks (Figure 3.11). This site reduces leakage of antibiotic post injection (Ian Wellby, pers. comm.).

Once each fish was deemed recovered, which took no more than 5 minutes, it was removed to a separate retention net in the river for further observation (Figure 3.12). Fish were retained in this way for an hour after the last fish was tagged, to ensure they had regained balance and were actively swimming, then collectively released as a group (Figure 3.13).



**Figure 3.4** Anaesthetic and recovery tanks (authors own photo).



**Figure 3.5** Removing scales from the incision site (authors own photo).





**Figure 3.6** Making the incision (authors own photo).



**Figure 3.7** Tag insertion (V9 model; authors own photo).



**Figure 3.8** Closing the incision with a single suture (authors own photo).



**Figure 3.9** Using Orabase™ protective paste to provide a temporary barrier (authors own photo).





**Figure 3.10** Post-surgery fish were placed in the recovery tank until they had recovered their righting reflex and were actively swimming (authors own photo).



**Figure 3.11** Antibiotic injection of 1 ml Baytril at 2.5% directly behind the dorsal fin was given while the fish was recovering. The injection was made on the same line as the erect last dorsal ray between the two main muscle blocks (authors own photo).




**Figure 3.12** Fish were retained in-river in holding nets for an hour after the last fish was tagged, before collective release as a group (authors own photo).



**Figure 3.13** Releasing a tagged common bream on 10<sup>th</sup> October 2010 at Bardney Bridge (authors own photo).

### 3.2.4 Tracking

After release, the movements of the common bream were monitored in the lower River Witham, Lincolnshire between Short Ferry (53°13'38"N; 0°21'23"W) and Boston (52°58'53"N; 0°1'46"W). Tracking results were obtained from up to twenty-six fixed VR2 and VR2W acoustic receivers (Vemco, Nova Scotia, Canada). These were fixed to wooden posts positioned in the river margins so receivers were maintained at approximately mid-water depth, a metre below the surface. Figure 3.14 illustrates how the number of fish being tracked and the number of receivers changed during the study period. Receiver coverage increased during the first eighteen months of the study and then remained constant. The number of fish tracked varied throughout the study, increasing when new fish were tagged (seven occasions) and decreasing as tag batteries expired and when tagged fish permanently left the study area.



**Figure 3.14** Receiver coverage and sample size of tracked fish. Grey bars represent the number of VR2 and VR2W acoustic receivers used to track the fish and, solid black line with black diamond markers, the number of fish tracked during the study period. No tracking took place during July-September 2008, as existing tags had expired and tagging new fish was delayed until water temperatures reduced during the autumn / winter months due to the benefits to fish recovery at higher dissolved oxygen levels.

### 3.2.5 Recapture and translocation of tagged fish

Fish were recaptured both intentionally, for translocation, and unintentionally, during fishing operations to capture fresh fish for tagging. When new fish were required for tagging, mobile tracking with a VR100 mobile receiver (Vemco, Nova Scotia, Canada) was undertaken to find areas where tagged fish were present and fishing for fresh fish was concentrated in these areas. Mobile tracking involved the use of an Omni-directional hydrophone which was placed into the



water to detect any tags within range of the receiver. Common bream are a shoaling species (Backiel & Zawiska, 1968) and it was reasoned that tagged fish would be associated with many untagged fish.

Recaptured tagged fish were isolated and identified with the VR100 mobile receiver (Figure 3.15) in separate bankside tanks, identification was confirmed by the presence of surgery scars. The incision site was inspected and photographed to show the extent of tissue regeneration. All fish were returned to the river alive.

On 16 March 2010 three fish (tag numbers: 12266, 12257 & 12255) were caught by wrap-around seine netting: 35 m by 3 m pull down and 50 m by 3 m wrap (Coles *et al.*, 1985) in the Sincil Dyke when three separate single haul nettings covered a length of drain of ~1 km. The fish were placed in individual large plastic bags (60 cm by 100 cm) containing approximately 20 L of water, bags were sealed at the top and the air space filled with oxygen from a bottle supply. Fish were then translocated ~35 km downstream (Figure 3.16) by road and released at Langrick Bridge (53°0'36"N; 0°6'52"W) so their spatio-temporal behaviour could be compared with control tagged fish that were present in the river, but not recaptured or translocated.



**Figure 3.15** Identifying and isolating recaptured tagged fish on the banks of the Sincil Dyke.

Fish were placed in tanks and tagged fish identified with a VR100 (Vemco, Nova Scotia, Canada) mobile tracking unit (authors own photo).



**Figure 3.16** Releasing a translocated fish at Langrick Bridge 35km downstream from capture site (authors own photo).

### **3.2.6 Data handling and statistical analysis**

Data were downloaded to a laptop computer using VR2PC and VUE software packages (Vemco, Nova Scotia, Canada). Data were collected in this manner at first on a monthly basis but later, as confidence in data security grew and the dataset became very large, every two-three months. Fish were also manually tracked with a VR100 (Vemco, Nova Scotia, Canada) mobile unit although no mobile tracking results were integrated into the dataset for subsequent analysis. If fish left the study area but later returned, they were still considered part of the study while outside the range of the study, but when fish left the study area and did not return they were no longer considered part of the study. Allocation of a km value to the nearest 0.1 km (measured using ArcMap v9.1 Geographic Information System, ESRI Ltd, Redlands, CA, USA) upstream of the tidal limits at Boston for each receiver allowed the movements for individual fish to be plotted.

There were five events when new fish were tagged while other tagged fish were being tracked in the river, allowing the behavioural effects of capture, handling and tagging to be compared between these two groups; freshly tagged (tagged) and previously tagged (control). The effects of the single translocation event of three fish were also compared. Two methods of quantifying the spatio-temporal behaviour of tracked fish were used. The cumulative distance moved between the receivers visited by an individual fish gave the Total Distance Moved (TDM) in km. The longitudinal distance between the most upstream and downstream detections (Young, 1999; Ovidio *et al.*, 2000) gave the linear range (LR) in km with positive values indicating ranges upstream while negative values indicating ranges downstream. Both TDM and LR were calculated for every fish tracked over the ten day period post operation for the effects

of tagging procedure analysis and over concurrent five day periods up to twenty five days post translocation, as all fish had homed back to the place of capture during this time period.


The effects of tagging and translocation on TDM and LR during each period following tagging and translocation were analysed by Mann-Whitney U tests, as data were not normally distributed. All analyses were performed with Minitab® v15.1.1 (Minitab Inc., PA, USA).

### 3.3 Results

All eighty-three common bream made a full recovery post-surgery and swam away strongly on release. One fish implanted was euthanased in-line with schedule one of the Animals Scientific Procedures Act (1986) after it failed to recover from anaesthesia, post-mortem revealed no surgical complications. Individual fish were tracked from 40-629 days (mean  $266.0 \pm \text{SD } 146.7$ ) producing a dataset of over 3.1 million detections.

Figure 4.17 illustrates the seasonal pattern of data collection with more detections being registered during the winter and spring periods. Tag battery life exceeded the manufacturer's specification where no automatic stop was placed on the tag, usually substantially. For example tag 250 (stated life 240 days) appeared intermittently in the study area for 1035 days, although only 400 days of data (from tagging in February 2007 to June 2008) were incorporated into the dataset for analysis as detections past this point appeared to be discontinuous suggesting that the transmitter was operating intermittently. Data from other tags which appeared to operate intermittent once post the stated tag life were also deemed unreliable due to its discontinuous nature and was thus omitted from the analysis. Tags with automatic stops all shut down in line with specifications. Subsequent data analysis showed no evidence of tag expulsion and survival rates of 100% were experienced for fish that stayed within the study area. All fish released showed upstream movements for at least two months, usually substantially more, post surgery. Continuous datasets were collected for all fish that stayed within the study area for up to 20 months post surgery and displayed several upstream and downstream movements.

A number of tagged fish left the study area and did not return (38%), they were last detected at possible exits from the study area; fish leaving the study area and returning after a period was a common occurrence. Figure 3.18 illustrates how the sample size of the study changed over time compared with what was projected, assuming no fish left the study area. In the first half of the study the sample size was close to the projected sample size, although at times were greater than expected; when tags ran longer than expected *e.g.* at the beginning of 2008. During the second part of the study sample size was lower than projected; as tracked fish left the study area. During the first half of the study four tagged fish (9%) left the study area, during the second half of the study twenty-eight tagged fish (70%) moved out of the study area; predominantly through the Billingham Skirth and Sincil Dyke.




**Figure 3.17** Data collected each month. Solid black line and black square markers is the total number of detections collected, dark grey line with triangular markers is the number of detections per tagged fish present and the light grey line with diamond markers is the number of detections per receiver in each month of the study.


### 3.3.1 Recaptured fish

A single haul seine netting on 3<sup>rd</sup> December 2007 at Bardney Bridge (53°12'27"N; 0°20'12"W) resulted in 990 adult common bream with masses between ~2-3 kg. The netting recaptured four tagged common bream, three (Tag numbers: 6070, 6073 & 6074) 51 days post-surgery and one (Tag number: 252) 275 days post-surgery. One fish (Tag number: 12275) was recaptured 76 days post-surgery by electro-fishing in the Sincil Dyke on 24<sup>th</sup> February 2009. One fish (Tag number: 12266) 384 days post-surgery and two fish (Tag number's: 12257 & 12255) 461 days post-surgery were recaptured during the nettings on 16<sup>th</sup> March 2010, which resulted in a catch of 1,270 adult common bream weighing between ~1-3 kg.

Recaptured fish appeared to be behaving naturally; they were associated with numbers of untagged fish, sometimes in very high densities. Physically they exhibited clean healing and tissue regeneration (see Figure 3.19) although in one case there was some haemorrhaging. There was no evidence of the tag expulsion process that has been described by Jepsen *et al.* (2008). All recaptured fish appeared to be in no worse a physical condition than when they were tagged. All three fish recaptured 51 day post-surgery and the single fish recaptured 76 days post-surgery still had sutures present. All four fish recaptured after 275-461 days post surgery displayed clean healing and suture absorption, with a small scar being the only evidence of the surgical procedure. This suggests that the sutures were absorbed in line with the manufacturers' specifications, which state that absorption is minimal until about the 90<sup>th</sup> post-surgery day and is essentially complete between 180 and 210 days when assayed in laboratory rats *Rattus norvegicus* (B.).




**Figure 3.18** Projected sample size compared with the actual sample size. Solid black line with black square markers represents the projected sample size assuming no fish left the study area, grey line with grey diamond markers is the observed number of tagged fish within the study area.




**Figure 3.19** Incision site of fish 6073 - 51 days post surgery. Note the very clean advanced healing and incision closure, suture still present (authors own photo).

### 3.3.2 Effects of tagging procedure

The mean TDM and the mean LR during the ten day period post-surgery and release of freshly tagged fish and control tracked fish for all five tagging events are presented as Figure 3.20 and Figure 3.21 respectively. There was no consistent pattern in the TDM by each group during the



**Figure 3.20** The mean total distance moved ( $\pm$  SE) for all five tagging events over the ten day period after tagging and release of freshly tagged fish (n=61) and control tracked fish (n=55).



**Figure 3.21** The mean linear range ( $\pm$  SE) for all five tagging events over the ten day period after tagging and release of freshly tagged fish (n=61) and control tracked fish (n=55). Positive values denote ranges upstream whilst negative values denote ranges downstream.

five tagging events, sometimes the control fish had higher TDM, sometimes the freshly tagged fish had higher TDM and sometimes they were equivalent. Similarly there was no consistent pattern in the LR of the two groups during the ten day period post surgery. Mann-Whitney U




tests revealed no significant differences, following Bonferroni correction, between the two groups in terms of TDM and LR during the ten day period post surgery for all five tagging events; results are presented in Table 3.2.

### 3.3.3 Effects of translocation

Following translocation all three fish returned to the Sincil Dyke, between 6-24 days after being released ~35 km downstream at Langrick Bridge. Fish 12255 (a male; Figure 3.22A) returned to the Sincil Dyke at 9am on 27<sup>th</sup> March, 11 days after release at Langrick Bridge. On release this fish moved directly upstream to Tattershall Bridge (~15km) within 24 hours where it stayed for 8 days then moved again directly upstream back to the Sincil Dyke. By contrast fish 12257 (a female; Figure 3.22B) returned to Sincil Dyke at 7am on 22<sup>nd</sup> March, 6 days after release. On release this fish moved downstream and upstream for a day, before moving directly back to the Sincil Dyke, with a slight pause of ~1 day at Tattershall Bridge. Finally, fish 12266 (a male; Figure 3.22C) returned to Sincil Dyke at 3am on 9<sup>th</sup> April, 24 days after release. This fish had moved directly upstream to Tattershall Bridge after release where it stayed wandering upstream and downstream over ~5 km of river for ~2 weeks. It then moved upstream to Bardney where it stayed for ~1 week before moving back ~7 km downstream, then moved upstream once more into the Sincil Dyke and remained highly mobile in the following months.

**Table 3.2** Mann-Whitney U tests results for comparisons of TDM and LR during ten day period post-surgery between freshly tagged (T) and control (C; previously tagged) fish during all five tagging events. Significant at < 0.01 following Bonferroni correction.

Event	Total Distance Moved (TDM)			Linear Range (LR)		
	Median T	Median C	Mann-Whitney U Test results	Median T	Median C	Mann-Whitney U Test results
1	18.30	1.45	U=16, n <sub>1</sub> =4, n <sub>2</sub> =7, P = 0.154	5.70	0.00	U=12, n <sub>1</sub> =4, n <sub>2</sub> =7, P = 0.027
2	6.10	14.45	U=70, n <sub>1</sub> =10, n <sub>2</sub> =6, P = 0.044	-0.95	-2.80	U=70, n <sub>1</sub> =10, n <sub>2</sub> =6, P = 1.000
3	6.70	3.40	U=224, n <sub>1</sub> =15, n <sub>2</sub> =19, P = 0.192	0.90	0.90	U=292, n <sub>1</sub> =15, n <sub>2</sub> =19, P = 0.283
4	3.60	3.30	U=127, n <sub>1</sub> =11, n <sub>2</sub> =13, P = 0.580	1.00	0.00	U=117, n <sub>1</sub> =11, n <sub>2</sub> =13, P = 0.236
5	4.00	1.00	U=242, n <sub>1</sub> =19, n <sub>2</sub> =12, P = 0.011	0.00	0.00	U=319, n <sub>1</sub> =19, n <sub>2</sub> =12, P = 0.511



**Figure 3.22** Fish plots for A; tag ID 12255 (male), B; 12257 (female) and C; ID 12266 (male) showing behaviour before and after translocation. Return journey to Sincil Dyke shown in red, which took 11, 6 and 24 days after release at Langrick Br respectively.




The mean TDM over concurrent five day periods up to twenty five days after release for the translocated group and non-translocated control group is presented as Figure 3.23. The translocated fish moved substantially more than the control fish during the first, third and fourth five day periods post translocation, otherwise values were comparable. Mann-Whitney U tests (Table 3.2) revealed no significant effect in any five days period following translocation.


The mean LR over concurrent five day periods up to twenty five days after release of the translocated group and control group is presented as Figure 3.24. The translocated fish moved long distances upstream as they returned to the Sincil Dyke, while the control fish slowly ranged downstream. Mann-Whitney U tests revealed significant differences in every concurrent five day period; results are presented in Table 3.3.

**Table 3.3** Mann-Whitney U tests results for comparisons of TDM and LR between fish captured and translocated ~35 km downstream, and control fish not captured or translocated.

Concurrent five day period	Total Distance Moved	Linear Range
1 (1-5 days)	U=25, $n_1=3$ , $n_2=7$ , $P=0.0674$	U=27, $n_1=3$ , $n_2=7$ , $P=0.0227$
2 (6-10 days)	U=19, $n_1=3$ , $n_2=7$ , $P=0.6475$	U=27, $n_1=3$ , $n_2=7$ , $P=0.0227$
3 (11-15 days)	U=19, $n_1=3$ , $n_2=7$ , $P=0.6485$	U=27, $n_1=3$ , $n_2=7$ , $P=0.0227$
4 (16-20 days)	U=24.5, $n_1=3$ , $n_2=7$ , $P=0.0855$	U=27, $n_1=3$ , $n_2=7$ , $P=0.0227$
5 (21-25 days)	U=16, $n_1=3$ , $n_2=7$ , $P=1.0000$	U=27, $n_1=3$ , $n_2=7$ , $P=0.0227$



**Figure 3.23** The mean total distance moved ( $\pm$  SE) over concurrent five day periods up to twenty five days (0-5 days open bars to 20-25 days dark grey bars) after release for the translocated group ( $n=3$ ) and non-translocated control group ( $n=7$ ).



**Figure 3.24** The mean linear range ( $\pm$  SE) over concurrent five day periods up to twenty five days (0-5 days open bars to 20-25 days dark grey bars) after release for the translocated group (n=3) and non-translocated control group (n=7). Positive values denote ranges upstream whilst negative values denote ranges downstream.

### 3.4 Discussion

#### 3.4.1 Tagging protocol

The tagging protocol described above was conceived after a review of the literature and discussion with practitioners. Implanting transmitters and transponders into the body cavity of fishes (see Baras & Philipart, 1989; Moore *et al.*, 1990; Wagner *et al.*, 2011) dates back to at least Johnson & Hasler (1977). Implantation into the peritoneal cavity is recommended for long-term tracking studies of fishes (Lucas & Batley, 1996) and is regularly used for cyprinid studies (Lucas & Batley, 1996; Huber & Kirchhofer, 1998; Lyons & Lucas, 2002; Winter & Fredrich, 2003; Fredrich *et al.*, 2003; Kuliskova *et al.*, 2009). Surgery is usually quick with the procedure taking approximately 5 minutes (*e.g.* Johnson & Hasler, 1977; Lucas & Batley, 1996).

In earlier studies transmitters were often attached externally (*e.g.* Langford, 1974, 1979 & 1981; Schulz & Berg, 1987; Baade & Fredrich, 1998) with, for example, the tag harnessed dorsally by means of degradable gut immediately to the rear of the dorsal fin (Langford, 1974), or attached with silver wire in front of the dorsal fin (Schulz & Berg, 1987). Many of these early studies were with relatively primitive technologies and tags were often only operational for a matter of days. However, technological advances have allowed the production of tags with a battery life of many years and surgical implantation eliminated the problems of fouling and infection associated with external attachment (Cooke *et al.*, 2012). Tags can also be placed into the fishes stomach via the mouth (*e.g.* Lucas & Johnstone, 1990), either inserted while under anesthesia or voluntarily by feeding within bait. This technique is ideal for short term study because it is less invasive than surgery, and the latter can even be carried out without the need to capture the target fish. Migratory salmonids tend not to feed once they enter the freshwater environment; as a consequence stomach tagging is widespread for studying this aspect of their

life-cycle (*e.g.* Laughton, 1989; English *et al.*, 2005). However, long-term retention is often poor in actively feeding fish as transmitter presence interferes with digestion and as a consequence are often regurgitated (Lucas & Johnstone, 1990; Neely & Steffensen, 2010) and can cause mortality (Neely & Steffensen, 2010). This technique is also considered inappropriate for cyprinids as they possess pharyngeal teeth (Lucas & Baras, 2001).

There are many types of anaesthetic available for use with fish (see Ross & Ross, 2008; Bridger & Booth, 2003). Cyprinid telemetry studies involving surgery have used 2-phenoxy ethanol (Winter & Fredrich, 2003; Kuliskova *et al.*, 2009), Tricaine methane-sulphonate (MS-222; Huber & Kirchhofer, 1998; Baade & Fredrich, 1998; Fredrich *et al.*, 2003; Winter & Fredrich, 2003; Bolland *et al.*, 2008) and Benzocaine (ethyl *p*-aminobenzoate; Lucas & Batley, 1996). However, in the UK only Tricaine methane-sulphonate (MS-222) has been licensed for use with fish by the Veterinary Medicines Regulations (VMR) 2011. Other products may be used if specified in licenses under Animals (Scientific Procedures) Act 1986.

A mid-ventral incision, as recommended by Lucas & Baras (2001), was used on the belly of the fish where the two muscle blocks meet. With this technique damage to the striated muscles is kept to a minimum, thus quickening recovery. Lateral incisions as used by Knights & Lasee (1996), cause greater tissue damage increasing recovery time. Incisions were kept as small as possible and were never greater than ~30 mm. The body wall of common bream is very tough, with layers of rigid overlapping scales that has made surgery difficult in previous studies (Jepsen *et al.*, 2002), but this may aid tag retention.

The size of incision necessary to implant a transmitter should be kept as small as possible to aid quick healing and recovery, although this will be dependent on the size of the transmitter itself. The incision site is often inconsistent between studies, for example; acoustic tags 8 cm by 2 cm were implanted through 3-4 cm lateral to the midline incisions cut with a scalpel and opened with scissors (Johnson & Hasler, 1977). Radio tags 10.9 mm diameter and 37.5 – 55.5 mm in length were implanted through a 2-3 cm incision 3 cm behind the base of the left pectoral fin (Fredrich *et al.*, 2003). Ventral incisions involve the least tissue damage and therefore should aid quick recovery and healing (Wagner *et al.*, 2011). The removal of scales to access the incision site makes surgery less complicated and is frequently practiced for intraperitoneal implantation (Cooke *et al.*, 2012). However, there is some evidence that this practice leads to higher levels of tissue necrosis compared with those fish implanted without removing scales (Bauer, 2005). Scales are embedded in the dermis so their removal can cause additional trauma (Harms, 2005). Thick scales, like those found on many large cyprinids blunt scalpel blades extremely quickly, severely reducing the quality of the underlying incision, and this is the most common and acceptable reason for removing scales (Cooke *et al.*, 2012).

Transmitters are usually inserted along the longitudinal axis of fish's body (*e.g.* Johnson & Hasler, 1977) and positioned above the pelvic girdle to reduce pressure on the body wall and the incision site itself, which is usually sutured with absorbable materials (*e.g.* Winter & Fredrich, 2003) either with a single suture or multiple separate sutures (*e.g.* Fredrich *et al.*, 2003; Kuliskova *et al.*, 2009) dependant on the size of the incision. Suturing is not a requirement with

small incisions (< 5 mm) to implant PIT tags (Skov, 2005). Accounts of post surgery tag retention vary; Baras (1997) retained radio tagged barbel in-river for six days, feeding chironomid larvae to allow recovery from the procedure and allow an evaluation of the success of surgery to be made. Short periods of retention are usually practiced once fish regain equilibrium and normal swimming behaviour this is usually in the order of minutes (*e.g.* Lucas & Batley, 1996; Fredrich *et al.*, 2003) rather than hours.

### **3.4.2 Recaptured fish**

Whenever surgery is involved fish will be subjected to disturbance and post-surgical healing rates vary according to species, age, the size of the incision and associated trauma and water temperature (Lucas & Baras, 2001). Although the survival of tagged fish is often not the best measure of the impact of the surgical procedure on fish in tagging studies (Jepsen *et al.*, 2008), it remains the simplest to measure without invasive or destructive techniques. All recaptured fish in this study showed advanced healing and no suture loss after 51 days with water temperatures between 4-14 °C. The four fish recaptured 275-461 days post tagging showed complete external recovery with the incision site hard to identify and the suture had disappeared.

Temperature is an important factor in healing (Knights & Lasee, 1996). Lucas & Batley (1996) found no evidence of mortality or morbidity in tagged barbel in the Yorkshire Ouse, UK, following the implantation of radio transmitters. They found recaptured tagged fish that had incision closure and advanced healing within 21 days and by 42 days recognition of the incision site was difficult. The observations of Lucas & Batley (1996) were made between late May and early June when UK water temperatures, and thus fish metabolism and healing rates, would have been higher. Lyons & Lucas (2002) reported no evidence of mortality of common bream in the river Trent, Nottinghamshire UK, with intraperitoneal implanted acoustic transmitters; surgery was also carried out during the summer months, post spawning.

Water temperatures during the tagging described here were between 5-13 °C and were carried out during the autumn and winter to utilise low water temperatures and associated benefits to fish recovery due to increased oxygen levels. Increased mortality, lower tag retention rates and higher risks of infection have been shown at higher temperatures, *i.e.* 20 °C (Knights & Lasee, 1996). However, whilst cooler conditions should aid fish recovery, healing rates and suture absorption appear to be slowed down.

The specification of sutures used here states that absorption is minimal until about the 90<sup>th</sup> post-implantation day and is essentially complete between 180 and 210 days when assayed in laboratory rats. The three fish recaptured 51 days post-surgery still had sutures evident, whilst the fish recaptured 275 days post-surgery had no sutures present, in line with the manufacturers' specifications. Low water temperatures may affect absorption rates and will certainly affect healing speed. Retention of sutures until healing is advanced is preferable to premature loss which may increase tag loss and mortality risk (Jepsen *et al.*, 2002). A single suture was seen as sufficient for incision closure due to the splinting effect of adjacent scales and body morphology; it also had the benefit of being fast and simple. Constant irrigation of the gills with

oxygenated weak anaesthetic solution throughout the procedure is often recommended (Bridger & Booth, 2003). However with the whole procedure taking as little as three minutes, this was deemed unnecessary and would further prolong the operation. Simplicity of operation as recommended by Jepsen *et al.* (2002) was adopted.

### **3.4.3 Shoaling behaviour**

Efforts to recapture fish often resulted in the capture of large numbers of individuals indicating tight shoaling behaviour (as described by Backiel & Zawiska, 1968; Phillips & Rix, 1985), possibly exaggerated by the time of year as common bream have been known to gather in high densities during the winter months (Borcherding *et al.*, 2002.). Observations, of tagged fish with other untagged fish (*e.g.* Clough & Ladle, 1997; Clough & Beaumont, 1998), and of fish engaged in activities such as migration (*e.g.* Baras, 1997) and spawning (*e.g.* Lucas & Batley, 1996) soon after tagging have been used to suggest ‘normal’ behaviour by the tagged fish and that they are not placed at a disadvantage by tag presence.

### **3.4.4 Fish that left the study area**

Incidence of fish leaving the study area was much more prevalent in the second half of the study. Twenty-eight of the forty tagged fish (70%) compared with four out of forty-three (9%) in the first half, appeared to leave the study area while they were being tracked. The spring and summer months saw the highest rate of fish leaving the study area (Figure 3.7), the reduction in sample size during spring/summer 2009 was due to fish moving out of the study area, the reduction in spring/summer 2010 was due to tag life being reached as well as fish moving out of the study area.

Where each group of fish were caught may provide a possible explanation for this high emigration rate in the latter half of the study. The fish tracked in the second half of the study were all caught in tributaries (Billinghay Skirth and Sincil Dyke), which are natural exits to the study area, whereas fish caught in the first half of the study were all caught in the main river channel several km away from any such possible exits. Fish displaying an affinity to the site of capture, site fidelity or occupying a home range in the areas close to exits from the study area are more likely to leave by these exits. The tags used in the second half of the study also had much longer operational lives which would have increased the likelihood of fish leaving the study area while tags were operational.

Common bream are classified as semi-migratory by Backiel & Zawiska (1968), floy tagging of 2,763 adult common bream re-released into the river Suck, Ireland, resulted in 110 recaptures, 35 were recorded at 2-59 km from the original tagging site while 75 were recaptured on site (Whelan, 1983). Goldspink (1978) marked 4,619 adult common bream in Tjeukemeer, a large shallow lake in the Netherlands, the majority of fish recaptured were from the lake, although recapture rates were low and this only represented 351 fish (7.5%). Common bream also travelled through connecting channels to adjoining waterbodies, with 101 fish (2.1%) caught outside the lake with individuals recorded up to 60 km from the release point. The

present study took place in a very large lowland riverine ecosystem and fish should be expected to move outside of the monitored area and into adjacent connected waterbodies. The seemingly high incidence of fish emigration from the study area is consistent with previous studies and there would appear to be a high level of interchange of individuals between adjacent connected drain networks.

#### **3.4.5 Effects of tagging procedure**

There were no consistent effects of the tagging procedure on the distances the newly tagged fish moved or their linear range immediately after being tagged, when compared with fish tagged previously. It is important to know if tagging disturbs the behaviour observed (Fredrich *et al.*, 2003) to validate the data collected and conclusions drawn. Previous studies have excluded data from the period immediately after tagging (*e.g.* one week by Allouche *et al.*, 1999) or excluded it from analysis to mitigate for the effects of tagging and handling on fish behaviour (Winter, 1996). Behaviour after release can vary with species. 88% of tagged dace in the River Frome, UK, moved upstream immediately after release, making large excursions on the day of release and three fish were lost outside of the study area (Clough & Beaumont, 1998). Unusual long-distance movement may occur during the first days of release after transmitter implantation (Mesing & Wicker, 1986). However, in contrast Lyons & Lucas (2002) observed no large movements (> 100m) of tagged common bream in the River Trent in the first hour after release and within 24 hours all nine fish had taken up a 'daytime position' which was heavily used throughout the rest of the tracking period, displaying 'normal' behaviour in the context of the whole study. The difference between these two reactions to surgery could reflect their vulnerability to predators, dace being a common prey item for pike, in the River Frome (Masters *et al.*, 2002) whilst the larger common bream had no real predators in the River Trent at this time. The low spatial resolution of the present study, when compared to say radio tracking, may have been unable to detect subtle behavioural changes following tagging. However in the context of this study there was no justification to exclude data collected immediately after fish were tagged.

#### **3.4.6 Effects of translocation**

Although the sample size for this experiment was low, as only 3 fish could be captured for translocation, some interesting results were achieved. Translocation did not significantly affect the activity levels of the fish translocated (although this could be a consequence of limited statistical power), but their linear ranges were significantly affected, as fish appeared to display 'homing behaviour' and moved back upstream towards the original capture site whilst control fish tended to move downstream at a slower rate over the same period (in line with the seasonal trend in movements / behaviours; see Chapter 4). It would appear that the translocated fish displayed a homing ability even over ~35 km of apparently homogenous river. Interestingly, all the translocated fish paused their upstream movements for varying periods of time in the Tattershall Bridge area.

How fish achieve navigation in complex habitats has been the subject of much speculation (see Walker & Hasler, 1949; Hasler & Wisby, 1958; Malinin, 1971; Carlson & Haight, 1972; Smith, 1985; Mesing & Wicker, 1986; Hert, 1992; Baras, 1997, Odling-Smee, 2003) but no firm conclusions have been made and it is likely that many mechanisms are used. However, olfactory cues are thought to be particularly important and are possibly the primary mechanism used by adult Atlantic salmon to locate their natal river, according to the home-stream odour hypothesis (Smith, 1985). Experiments have suggested this mechanism is also important for fish navigating 'home' following translocation. For example, fourteen radio tagged brown trout were displaced over 800-3600 m upstream and downstream in the River Eden, Scotland. Twelve of these fish subsequently returned to the areas from which they were taken, seemingly to follow specific orientation cues (*e.g.* olfactory) rather than searching at random (Armstrong & Herbert, 1997).

Here, data lend support for the notion that fish are capable of relocating by directed movements using specific orientation cues, rather than by accident or a random search pattern. Prior to translocation all fish were relatively resident in the Sincil Dyke, and had been for some time. Following translocation, once fish had returned 'home', they tended to remain in relatively localized areas of the channel for a subsequent period (Figure 3.22). The preceding tracking data of the fish allows some assessment of each fish's 'familiar area' (the zone through which the fish could remember having moved; Bovet 1992). Interestingly the one fish that took the longest to return 'home' had prior knowledge of the lower river where it was translocated. Therefore, this fish could be using olfactory beacons (the presence or absence of familiar odours emanating from the familiar area; Halvorsen & Stabell 1990), or any of a range of other potential cues, such as visual and olfactory landmarks, or areas of distinct waterflow patterns (Armstrong & Herbert 1997). Whereas the other two translocated fish had not experienced the lower river in the preceding ~15 months and may have no experience of its visual and olfactory landmarks to use as navigational aids, and were the quickest to return 'home'.

#### **3.4.7 Possible experimental bias**

During the first half of the study fish were sampled by a variety of methods which had the potential of introducing bias as they may be more or less selective against certain behavioural types *e.g.* rod and line may select for bolder fish than electro-fishing (see Biro & Post, 2008; Klefoth *et al.*, 2012). Ten fish tagged in October 2007 were caught by rod and line, and although generally healthy some fish showed signs of physical damage, exhibiting scale loss, associated inflammation and slight bacterial infection. Subsequently when three of these fish were recaptured in November 2007, they were in similar physical condition relative to the pre-surgery condition, although healing at the incision site was evident. They were also conspicuous in the shoal of 990 captured individuals in that they were in relatively poor physical condition. It is possible that out-of-condition fish may be more susceptible to capture by rod and line, making the method unsuitable for fish capture for tracking studies as the technique is best used with fish as fit as possible. Fish were sampled from just the main river channel in the first two years of the study and from just inside tributaries during the third year, which also has the potential to

influence results as fish captured in a certain channel type may have an affinity with that habitat type.

In conclusion, this study detected no impacts of tagging on the spatial temporal behaviour of common bream in respect to both the distances they moved and their linear range. Recaptured fish had clean healing/scars and appeared to be exhibiting 'normal' behaviour in that they were part of large shoals of common bream. Translocated fish appeared to exhibit homing behaviour, moving quickly back to the site of capture.



## **Chapter 4 Seasonal movements with shifts in lateral and longitudinal habitat use.**

### **4.1 Introduction**

Large lowland rivers are often heavily influenced by human activity, resulting in uniform riverine ecosystems with regard to both physical form and water flow (Brookes, 1988) that present many challenges to fish populations (Linfield, 1985). Direct interventions on lowland riverine ecosystems for the purposes of navigation (impoundment and barriers), flood control (artificial levee construction), land drainage (artificially high peak flows) and flow regulation (modification of hydrograph) are prevalent. Changes in river morphology generally shift riverine ecosystems from lentic towards lotic environments (Welcomme, 1994) through habitat degradation and reduced longitudinal and lateral connectivity. Negative impacts on fish community structure centre on the reduced ability of fish to exploit specialised habitats required for the completion of their life-cycles (Pinder, 1997), specifically impacting those fish *e.g.* common bream adapted to utilise lentic habitats for spawning, nursery and flow refuge. Therefore, lowland rivers often require rehabilitation, which is now being driven by legislation (*e.g.* EU Water Framework Directive 2000/60/EC). However, long-term field studies documenting the influence of environmental variables on longitudinal and lateral movements of adult fish are required to provide empirical evidence to support management actions aimed at conserving populations and rehabilitating habitats. In Europe, large lowland riverine ecosystems are dominated by cyprinid fishes but studies of their seasonal movements and habitat use in such environments are relatively rare.


Acoustic telemetry is an established technique for long-term study of the movements and habitat use of large fish (Lucas & Baras, 2001; Jellyman, 2009), including common bream (Langford *et al.*, 1979; Lyons & Lucas, 2002). In this study acoustic telemetry is used to determine seasonal patterns of activity, distribution and habitat use of common bream within the lower River Witham between November 2006 and June 2008, and October 2008 and August 2010. Three habitat types were identified; a deep tributary, shallow tributaries and the main river channel, it was hypothesised that these different habitats would be used at different times of the year and in relation to changing environmental conditions, such as flow rate and temperature.

### **4.2 Materials and methods**

#### **4.2.1 Study area**

The River Witham, in eastern England, rises near South Witham (52°45'44"N; 0°37'38"W), Lincolnshire, UK, and flows north to Lincoln and then south east to Boston (52°58'53"N; 0°1'46"W), where it discharges into The Wash. The study area (Figure 4.1) is a continuous open reach of approximately 40 km of the non-tidal lower river and associated tributaries starting upstream near Short Ferry (53°13'38"N; 0°21'23"W) where the Barlings Eau tributary enters the Witham and ending downstream at the tidal limit, Grand Sluice in Boston (52°58'53"N; 0°1'46"W). This uniform man-made channel is managed for the purposes of navigation and land drainage, and has been straightened, widened and deepened (Wheeler, 1990) with high levees

constructed (Environment Agency, 2008) 3-4 metres from the water's edge on both banks (Forbes & Wheeler, 1997). This environment presents fish populations with a variety challenges such as the absence of a functional floodplain, large floodwater discharge, high velocity flows in the lower reaches (Environment Agency, 2008) leading to flush out (Linfield 1985) and poor in-river and marginal habitat. Intensive use of adjacent agricultural land results in high nutrient and silt levels from run-off leading to eutrophication and the potential to reduce water quality. River levels are artificially lowered during the winter months to increase the rivers flood storage capacity (Figure 4.2). During April - October the normal river level is maintained at approximately 1.5 m above Ordinance Datum Newlyn (ODN), and from November - March the level is reduced to and maintained at 1 m ODN, during dry conditions. The main channel is trapezoidal and canalised with a depth of 2-4 m at normal



**Figure 4.1** Map of study area; the lower River Witham, showing the river zone number (1-7) to which receivers were allocated for the ‘monthly longitudinal distribution’ analysis. Place names are also included, tributaries are labelled in italics and capture and release sites are labelled by capital letters (see Table 1). Insert (top right) is an expanded view of the study area showing Bardney Lock ‘z’ which is generally impassable to fish, the old river loop that is fed from the main navigation channel at an impassable weir ‘w’, and the Sincil Dyke which contains more than one receiver (‘x’ and ‘y’).

summer level with a width of 30-40 m, and usually hosts substantial macrophyte growth during the summer months.

The study area contains a number of tributaries, which group into two types according to depth. Firstly, the Sincil Dyke is a linear channel ~2 m deep at summer level, approximately 10-15 m wide and flows from its inception at Lincoln to the west, parallel with the main river to Bardney where it enters the lower river (Wheeler, 1990). Secondly, Nocton Delph, Timberland Delph, Duckpool Drain, Billingham Skirth, River Bain and River Sleas (Figure 4.1) are shallow (0.6-1 m depth at summer level) linear channels approximately 10-25 m wide. These channels carry water drained from the surrounding agricultural land, but are heavily silted and are generally slow flowing to static in normal dry conditions. Marginal habitat of Sweet Reed Grass *Glyceria* spp. and Common Reed *Phragmites australis* (Cav.) is prevalent in all of these tributaries and there is substantial macrophyte growth during the summer months, interspersed with heavy filamentous algae *Cladophora* spp. growth on the bed.

The layout of the upper end of the study area (Figure 4.1, insert) warrants explanation. The Sincil Dyke joins the main river downstream of Bardney Lock (Figure 4.1, insert 'z') which is a barrier to fish passage but allows boat navigation to Lincoln. Upstream from Bardney lock the 'old river' loops round past the Barlings Eau confluence back towards the navigation channel, where the river flows from Lincoln entering the old river at an impassable weir (Figure 4.1, insert 'w').


#### **4.2.2 Sampling and tagging procedures**

Eighty-three adult common bream with a mean fork length of 485.6 mm ( $\pm$  SD 16.8; range 440-522 mm) and a mean mass of 2.37 kg ( $\pm$  SD 0.25; range 1.92-2.94 kg), were caught from the study area in seven groups by electro-fishing (240 V, 4/5 A, pulsed DC), rod and line or seine netting. Details of the tagging and tracking are described in Chapter 3. Tributary receivers were positioned inside each channel to ensure detections on these receivers were from fish inside the tributary and not in the adjacent main river. The Sincil Dyke was monitored with more than one receiver (Figure 4.1, insert receivers 'x' and 'y').

Independent range tests carried out *in-situ* showed V9 tags were detected up to ~200 m and V13 tags up to ~400 m in macrophytes and macroalgae free conditions (author's observations). Receivers were visited on a regular basis (weekly during summer months) to remove macrophytes and macroalgae from the surrounding margins (~4 m<sup>2</sup>) so they were accessible to signals in the main river channel, which remained relatively weed free due to boat traffic. Data from the receivers were regularly downloaded onto a laptop computer using VR2PC and VUE software packages (Vemco, Nova Scotia, Canada). Occasionally receivers would record detections for tags that were not present. The reason for these false detections was not clear, but could result when tag signals clashed. False detections were more prevalent on receivers near underwater structures, such as bridge supports, and could possibly be caused by reflected signals and echoes. These obvious false detections were removed prior to analysis.

#### 4.2.3 Environmental data

Flows were gauged by flat V broad crest weir ( $< 2 \text{ m}^3\text{s}^{-1}$ ) and ultrasonics ( $> 2 \text{ m}^3\text{s}^{-1}$ ) on the Barlings Eau at Langworth bridge, ~6 km upstream of the study area which was used as a proxy for flow rate in the main river in the study area. River level was gauged by stilling well at Kirkstead Bridge, near the Timberland Delph confluence, approximately midpoint of the study area (Figure 4.2). Average daily air temperatures were from the weather archive recorded at RAF Coningsby, Lincolnshire. Water temperatures ( $^{\circ}\text{C}$ ) were measured at two points (Bardney Lock and Tattershall Bridge) in the main river from 20 February 2007 on an hourly basis with Aquatic and Aquatic 2 dataloggers (TinyTag, Gemini Data Loggers UK Ltd., Chichester, UK).



**Figure 4.2** Flow ( $\text{m}^3 \text{s}^{-1}$ ; black line) gauged on the Barlings Eau at Langworth Bridge (~6 km upstream of the study area) and river level (m ODN; grey line) recorded at Kirkstead Bridge, near the Timberland Delph confluence (see Figure 4.1) approximate midpoint of study area.

#### 4.2.4 Data handling

Allocation of a km value (measured using ArcMap v9.1 Geographic Information System, ESRI Ltd, Redlands, CA, USA) upstream of the tidal limits at Boston for each receiver allowed the movements for individual fish to be plotted. Tributary receivers were allocated the km value measured to the mouth of the tributary. ‘Monthly distance’ (km) was calculated as the cumulative distance moved between receivers by an individual fish during each calendar month and was used as a measure of activity.

New receivers were added during the study and to overcome this inconsistent temporal receiver coverage, receivers were grouped into seven river ‘zones’ ~5 km in length (except zone seven which was ~10 km; Figure 4.1). Each zone had at least one receiver present throughout the whole study period. In order to provide a measure of the longitudinal use of the study area, ‘Monthly longitudinal distribution’ was calculated as the number of detections per tagged fish, per zone, per month (corrected for the number of receivers per zone).

The study area consists of three distinct habitat types; the main river channel, the shallow tributaries and a deeper tributary, the Sincil Dyke. To analyse seasonal patterns in lateral habitat use the number of detections per fish, per month in each habitat type was compared. The

occupancy of the tributaries in relation to water temperature and flow, was analysed using daily detections per tagged fish, as opposed to monthly detections, in order to capture daily fluctuations in flow and temperature. When analysing Sincil Dyke occupancy and flow rate, only data from the single receiver positioned immediately inside this tributary (Figure 5.1, insert receiver 'y') were used. Air temperatures were used to analyse monthly trends, as this dataset was complete. Water temperatures were used to analyse daily tributary occupancy, because this dataset, although partially incomplete, offered greater representation of in-river conditions. Seasons were defined as follows; spring (March 21 – June 20), summer (June 21 – September 20), autumn (September 21 – December 20) and winter (December 21 – March 20).

#### **4.2.5 Statistical analysis**

The effects of gender, month and temperature on 'monthly distance' were analysed with GLM ANOVA and Pearson correlation respectively. Data for 'Monthly longitudinal distribution' and occupation of the three different habitat types are counts and thus were analysed using a generalised linear model, with a Poisson error distribution and a log-link function (Crawley, 2002). Maximal models were constructed with the significance of the independent variables assessed following stepwise deletion (starting with the higher order interaction terms) and changes in residual deviance assessed for significance using F-tests, which were used as the data were found to be over dispersed (Crawley, 2002). The same analysis was applied to the number of daily detections (also count data) in the tributaries in relation to the environmental variables of flow and temperature. Simple parametric analyses were performed using Minitab 15® (Minitab Inc., USA), whilst the generalized linear models were performed in S-plus (TIBCO Software Inc., California, USA).


### **4.3 Results**

A dataset of over three million fish detections was collected. Individual fish were tracked from 40-629 days (mean  $266.0 \pm \text{SD } 146.7$ ). Some tagged fish left the study area, being last detected at possible exits from the study area. Not all fish returned to the study area. Data for those fish that did not return were included in the analysis up to the point at which they were last detected within the study area.


#### **4.3.1 Monthly distance and monthly longitudinal distribution**

The largest 'monthly distance' recorded by a single fish was 118.5 km, as it moved upstream and downstream during April 2007. Month had a significant effect on 'monthly distance', *i.e.* common bream activity (ANOVA;  $F_{1,41} = 6.42$ ,  $P < 0.0001$ ) with the tagged common bream moving relatively small distances during the cooler months of November through to March in comparison to the warmer months of April through to October (Figure 4.3 and 4.4). There was no effect of gender ( $F_{1,41} = 1.76$ ,  $P = 0.185$ ) nor the interaction between month and gender ( $F_{1,41} = 0.67$ ,  $P = 0.941$ ) on 'monthly distance'. Mean 'monthly distance' was positively correlated with mean monthly air temperature (Pearson correlation  $r = 0.698$ ,  $df = 42$ ,  $P < 0.001$ ).


The ‘monthly longitudinal distribution’ varied over the course of the study (Figure 4.5) and the twelve month cycle (Figure 4.6). Log linear analysis of the number of detections per zone (corrected for the number of receivers present per zone) revealed a significant effect of zone ( $\Delta$  deviance = 679695,  $df = 6$ ,  $F = 96$ ,  $P < 0.0001$ ), month ( $\Delta$  deviance = 71221,  $df = 11$ ,  $F = 5.5$ ,  $P < 0.0001$ ) and the interaction between zone and month ( $\Delta$  deviance = 203960,  $df = 66$ ,  $F = 2.6$ ,  $P < 0.0001$ ). Thus, fish tended to occupy (or avoid) certain longitudinal zones over others and this pattern of longitudinal distribution varied according to month. For example fish were more likely to be detected in zone 1 during the cooler winter months (November through to March). The number of longitudinal zones occupied by the tagged fish increased during the spring and summer months which coincided with an increase in activity (as determined by mean ‘monthly distance’).




**Figure 4.3** Mean ( $\pm$  SE) ‘Monthly distance’ (km; grey columns) in relation to mean monthly air temperature ( $^{\circ}\text{C}$ ; black line). Gap between July and September 2008 represents period of no tagged fish in the study.



**Figure 4.4** Mean ( $\pm$  SE) 'Monthly distance' (km; grey columns) in relation to mean monthly air temperature ( $^{\circ}$ C; black line), averaged over the twelve month cycle.



**Figure 4.5** 'Monthly longitudinal distribution': zone 1 (upstream) to zone 7 (downstream).



**Figure 4.6** ‘Monthly longitudinal distribution’ averaged over the twelve month cycle for each zone of the river: zone 1 (upstream) to zone 7 (downstream).

#### 4.3.2 Use of lateral habitats

Month had a significant effect on the number of detections per fish in the shallow side tributaries ( $\Delta$  deviance = 24703,  $df = 11$ ,  $F = 94.6$ ,  $P < 0.0001$ ) and in the deeper Sincil Dyke ( $\Delta$  deviance = 10410,  $df = 11$ ,  $F = 39.8$ ,  $P < 0.0001$ ). The shallow tributaries were visited by the fish in the spring and apart from a few detections, the fish rarely ventured into the shallow tributaries outside of this time. By contrast, the deeper Sincil Dyke was used more frequently by the common bream during the winter months (Figure 4.7 and 4.8).

Log-linear analysis revealed that the corrected number of daily detections in the shallow tributaries was not related to daily water temperature ( $\Delta$  deviance = 79.1,  $df = 1$ ,  $F = 1.55$ ,  $P = 0.2$ ) nor the interaction between water temperature and flow ( $\Delta$  deviance = 24.8,  $df = 1$ ,  $F = 0.47$ ,  $P = 0.49$ ), but was negatively related to flow ( $\Delta$  deviance = 1604,  $df = 1$ ,  $F = 31.2$ ,  $P < 0.0001$ ). However, the non-significant effect of water temperature is not surprising given spring and autumn temperatures are roughly equivalent. Rather it appears that the common bream enter the shallow tributaries during the spring as water temperatures are increasing (Figure 4.9). To examine this relationship, the number of corrected detections per month were analysed in relation to the monthly change in water temperature and monthly flow rates. There was no significant interaction between monthly change in water temperature and monthly flow rate ( $\Delta$  deviance = 19.9,  $df = 1$ ,  $F = 1.6$ ,  $P = 0.2$ ). However, there was a significant positive relationship between corrected monthly detections in the shallow tributaries and change in water temperature ( $\Delta$  deviance = 406,  $df = 1$ ,  $F = 34.1$ ,  $P < 0.0001$ ) such that fish tended to move into shallow tributaries as water temperature increased (Figure 4.9). There was a negative relationship between monthly flow rate and the corrected number of fish detections ( $\Delta$  deviance = 130,  $df = 1$ ,  $F = 10.9$ ,  $P = 0.02$ ). It is worth noting that the effect of flow is likely to be confounded by



seasonal differences in flow rate, as few fish enter the shallow side tributaries during the autumn and winter, when flow rates tend to be high.

The occupancy of the deeper Sincil Dyke tributary appears to follow a different pattern. Log-linear analysis of data collected at the entrance to the Sincil Dyke tributary revealed a significant effect of water temperature on the corrected number of daily detections ( $\Delta$  deviance = 4133,  $df = 1$ ,  $F = 128.3$ ,  $P < 0.0001$ ), with more fish detections occurring at lower water temperatures. The analysis also revealed a significant positive relationship between occupancy of the entrance to the Sincil Dyke and flow rate (Figure 4.10, insert receiver 'y';  $\Delta$  deviance = 257.0,  $df = 1$ ,  $F = 8.04$ ,  $P = 0.004$ ), but no interaction between water temperature and flow rate on the corrected number of daily detections ( $\Delta$  deviance = 57,  $df = 1$ ,  $F = 1.78$ ,  $P = 0.18$ ). Examination of the corrected number of daily detections in relation to flow rate on a seasonal basis revealed the common bream to be more likely to be found in the slow-flowing Sincil Dyke during periods of high main-channel flow in the Autumn months only ( $\Delta$  deviance = 1179,  $df = 1$ ,  $F = 40.9$ ,  $P < 0.0001$ ), with no relationship being observed in the spring ( $\Delta$  deviance = 27.6,  $df = 1$ ,  $F = 0.73$ ,  $P = 0.39$ ), summer ( $\Delta$  deviance = 76,  $df = 1$ ,  $F = 2.79$ ,  $P = 0.09$ ) and winter ( $\Delta$  deviance = 116,  $df = 1$ ,  $F = 3.25$ ,  $P = 0.07$ ).

Collectively, these data show that adult common bream used shallow tributary habitat during the spring as temperatures increased. By contrast, the deeper Sincil Dyke was used throughout the winter and during high main channel flows in the autumn.

## **4.4 Discussion**

### **4.4.1 Monthly distance and monthly longitudinal distribution**

In general fish were more evenly distributed, occupying greater lengths of the river channel, in the spring and summer in comparison to the cooler autumn and winter months, during which the common bream were generally aggregated in one or two locations. Activity levels (monthly distance) peaked in the spring, with one individual moving nearly 120 km in a month. Baade & Fredrich (1998) also recorded a peak in activity in the spring when studying roach in the River Spree, Germany, and these observed seasonal patterns are typical of poikilotherms in temperate parts of the world. Part of these circannual patterns is likely to be related to reproductive activity, with fish entering the shallower tributaries primarily during the spring and early summer to spawn when water temperatures increase. Langford (1974, 1979, 1981) found that common bream in the River Witham moved from 400 m to 8 km over periods of 2-5 days. Whelan (1983) observed common bream in the River Suck, Ireland moving up to 56 km over a 4 year period.


Lucas and Baras (2001) suggest that common bream would tend to be more nomadic than rheophilic species, such as barbel, particularly in homogeneous river environments. The behaviour patterns observed during this study partly fit this description with some common bream covering considerable distances. However, others moved over relatively small distances; this variation in behaviour is consistent with Lyons and Lucas (2002). Langford (1974, 1979) tracked both indigenous and introduced common bream in both the River Witham and River

Thames, and observed considerable variation in the directions and distances moved by individual fish. The Witham fish appeared to be more mobile than the Thames fish, although this may have been an artefact of the size of the study area; 13.5 km on the Witham and 5 km on the Thames, as size of available environment can influence home range size (Woolnough *et al.*, 2009).


Kell (1991) found aggregations of common bream in the Sixteen Foot Drain (Cambridgeshire, UK), which has a similar cross-sectional profile to the River Witham. Here, the tagged common bream were most frequently recorded in zones 1 and 5 (Figure. 4.1); sites where tributaries join the main channel. Confluences are known to be associated with the deposition of organic sediments (Carling, 1995), which harbour benthic food organisms (Rice *et al.*, 2001; Sevensen *et al.*, 2009), preyed upon by common bream (Backiel & Zawisza, 1968), thus it is likely that the common bream congregate in these areas to forage (see also Lucas *et al.*, 1998a).

The lower section of the study area (zones 6 and 7; ~15 km upstream of Boston) was infrequently utilised by the tagged fish and was only visited during the spring/summer. Hydroacoustic surveys carried out in 2006, 2007 and 2008, during April and October each year, revealed relatively low fish densities in this area (see Chapter 2). Due to the low lying nature of the study area, the lower river is often run-off to achieve water level differences further up the catchment that allow certain flood prone areas to be drained by manual penstock sluice. This lower section contains no tributary habitat, is particularly uniform and straight in nature and can experience high flows upto  $160 \text{ m}^3\text{s}^{-1}$  (Environment Agency, 2008) during rainfall events. This makes this section of river particularly inhospitable to fish and any fish taking up residence in this lower section risk being flushed out to tidal waters (Linfield, 1985).


The long-term analysis of 'monthly distance' and 'monthly longitudinal distribution' of common bream revealed seasonal patterns in their behaviour, with fish being more active during the relatively warmer months of the year and occupying a greater range of the available main-channel habitat. In studying the movements of common bream over several years it has been possible to identify the use of three broad habitat types at different times of the year and in relation to different environmental conditions. During the summer, the common bream tended to occupy the main river channel, whilst in the cooler autumn months and especially when main channel flow rates are high, the common bream took refuge in the slower flowing, yet relatively deep Sincil Dyke tributary. As temperatures increased during the spring, the fish were more likely to be found in the shallow tributaries, using these as spawning habitat.




**Figure 4.7** ‘Lateral habitat use’ mean proportion of total detections recorded in the shallow side-channels (black), the deeper Sincil Dyke (light grey) and the main river channel (dark grey).



**Figure 4.8** ‘Lateral habitat use’ mean proportion of total detections recorded in the shallow side-channels (black), the deeper Sincil Dyke (grey) and the main river channel (dark grey) averaged over the twelve month cycle.



**Figure 4.9** Daily detections per tagged fish present, recorded in the shallow tributaries (grey columns) and mean daily water temperature (°C; black line).



**Figure 4.10** Daily detections per tagged fish present, recorded in the Sincil Dyke tributary (grey columns) and mean daily flow recorded at Langworth Bridge on the Barlings Eau ( $\text{m}^3 \text{s}^{-1}$ ; black line).

#### 4.4.2 Use of lateral habitats

Heavy modification of the River Witham's channel has reduced longitudinal and especially lateral connectivity making the available tributaries the only off-line habitat available to the fish population. Lateral habitats are often essential to the life cycle of riverine fish populations, providing spawning and nursery habitat (Staas & Neumann, 1994; Grift *et al.*, 2001; Pollux *et al.*, 2006; Nunn *et al.*, 2010), shallow, warm water for enhanced growth (see Garner *et al.*, 1998), refuge during flood events (Copp, 1997; Pinder, 1997; Nunn *et al.*, 2010), foraging sites (Nunn *et al.*, 2007) and overwintering habitats (Borcherding *et al.*, 2002; Nunn *et al.*, 2010). During May 2008 pre-spawning behaviour by male fish was observed (Giles, 1994; Figure 4.11) in which aggressive guarding of marginal territories (Figure 4.12) and spawning were observed in the shallow tributaries (author's observation). Successful spawning was confirmed by the presence of common bream larvae in subsequent days. Common bream are known to spawn on macrophyte substrates in areas of low current velocity (Pinder, 1997) and marginal reedbeds (Giles, 1994). Both are habitat features of the shallow tributaries joining the lower River Witham but are lacking in the canalised main channel, with its steep, stony embankments, regulated flow dynamics and reduced hydrological connectivity to the floodplain. Such modification appears to be characteristic of many large European rivers and tributaries are likely to be particularly important for the recruitment of fish (Pollux *et al.*, 2006). Where rivers are allowed to meander, common bream have been shown to spawn in the slack-water of oxbows (Molls, 1999; Borcherding *et al.*, 2002). Thus, in the absence of natural meanders, shallow tributaries with low current velocities are likely to be important spawning habitats, functionally similar to lentic floodplain waterbodies of natural riverine ecosystems.



**Figure 4.11** Pre-spawning behaviour, males guarding marginal territories, observed in Nocton Delph May 2007 (authors own photo).



**Figure 4.12** Aggressive interaction between two males occupying adjacent marginal territories, Nocton Delph May 2008 (authors own photo).

The entrance to the Sincil Dyke was used more heavily during times of high flow especially during the autumn, when water temperatures were falling. Under such conditions, fish metabolism will be low (Wieser, 1991) and common bream may find it physically challenging to maintain their position during high flow conditions. Stream dwelling giant kokopu *Galaxias argenteus* (G.) in New Zealand have been shown to utilise low-velocity habitat during periods of high main channel flow (Bruno & Gerard, 2002), whilst adult chub have been observed to move into backwaters during peak flows in the French Upper Rhône (Allouche *et al.*, 1999).

Drainage ditches of floodplain lakes in the lower Rhine, Germany, were used for over-wintering by aggregations of fishes, including common bream, when water temperatures dropped below 10°C (Borcherding *et al.*, 2002) and radio tracked roach were found more often in the oxbows of the River Spree during the colder winter months (Baade & Fredrich, 1998). Similarly here, receivers positioned further inside the Sincil Dyke revealed this as a significant over-wintering area away from the main channel.

During the spring and summer months, adult common bream travelled on average 20-30 km per month, with one individual travelling just under 120 km in a month (April). During three of the four years of study activity peaked in April. This peak corresponded with an increase in the longitudinal habitat distribution and movements into the shallow tributary habitats. Baade & Fredrich (1998) reported a similar spring peak in the activity of roach in the River Spree, Germany. In 2007 and 2009 a second summertime peak in activity occurred later in the summer, this is likely to be temperature driven, the midsummer drop in activity is possibly linked to the cessation of spawning activities with associated energetic costs (Wieser, 1991). The study area here is atypical of many heavily modified lowland rivers, offering a long 40+ km reach without

barriers to fish passage. The findings of this study demonstrate the potential common bream have for moving over large distances in relatively short periods of time, illustrating the implications of reduced hydrological connectivity that leads to habitat fragmentation in modified rivers. Common bream are described by Backiel & Zawisza (1968) as 'semi-migratory' (when occupying unmodified habitats) making long migrations in the spring and autumn from the Black, Caspian and Azov seas of up to 100 km from brackish feeding areas in the lower reaches and deltas of large rivers. Langford (1974, 1979, 1981) used telemetry to observe common bream in a discrete 13.5 km reach of the River Witham (between Lincoln and Bardney Lock, upstream of the area of study here) and found that they moved 400 m to 8 km over periods of 2-5 days, whilst Whelan (1983) found common bream in the relatively unmodified River Suck, Ireland, to move up to 56 km during a five year mark-release-recapture study, but the timescales for such movements are unclear.

During the first half of this study fish were sampled by a variety of methods which had the possibility of introducing bias, as they may be more or less selective against certain behavioural types *e.g.* rod and line may select for bolder fish than electro-fishing. However, the broad trends in fish activity and habitat use observed in Figure 4.3 to 4.9 are consistent across 43 months of study, thus the results are unlikely to be an artefact of the different sampling techniques. Fish were sampled from just the main river channel in the first two years of the study and from just inside tributaries during the third year, although this does not seem to have influenced the consistent trends in the results. The tags used at the start and end of the study had different detection ranges, although given the relatively coarse resolution of the tracking, with fixed receivers positioned ~2-3 km apart; it is not thought that this influenced the patterns observed.



## **Chapter 5 Seasonal water level manipulation for flood risk management influences home-range size.**

### **5.1 Introduction**

The increased threat of flooding as a result of climate change via its effects on the level (Groisman *et al.*, 1999 & 2005) and intensity (Huntington, 2006) of rainfall requires those authorities responsible for alleviating flood risk to increase the management and modification of rivers (Wilby *et al.*, 2008). Riverine ecosystems drain water falling on the earth's surface and conduct it to the sea (Welcomme, 1994). These systems are amongst the most human-degraded ecosystems worldwide (Malmquist & Rundle, 2002; Huckstorf *et al.*, 2008) with biodiversity threatened by water extraction, flow regulation, channelisation and habitat degradation (Welcomme, 1994; Pinder, 1997; Huckstorf *et al.*, 2008). Although large lowland rivers support a significant proportion of the world's fish diversity (Huckstorf *et al.*, 2008) the majority of these environments, especially in Europe, have been modified through direct interventions that alter river morphology and reduce longitudinal and lateral connectivity (Welcomme, 1994; Cowx & Welcomme, 1998).

European lowland rivers have been particularly susceptible to modification due to their locality to sea ports, which leads to increased industrialisation, whilst fertile floodplains have been converted to agricultural land (Huckstorf *et al.*, 2008) via increased drainage and land clearance, with associated effects on habitat quality. Lowland rivers are also impacted by actions upstream; silt, pollutants and nutrients introduced or disturbed in the upper reaches are transported and can accumulate in the lower reaches (Welcomme, 1994). Run-off from agricultural land and sewage discharges have enriched lowland rivers leading to elevated levels of nitrogen and phosphorus (*e.g.* Pinder *et al.*, 1997) leading to eutrophication (Edmondson, 1991; O'Sullivan, 1995). The majority of large lowland rivers in Europe are managed with respect to navigation, power generation, water storage, development, flood risk management and abstraction all of which have the potential to impact on fish populations (see Mann, 1988; Junk *et al.*, 1989; Welcomme, 1994; Copp, 1997; Hadderingh & Bakker, 1998; Turnpenny, 1998; Buijse *et al.*, 2002; Huckstorf *et al.*, 2008) with many lowland rivers in Europe subject to flow and water level regulation (Buijse *et al.*, 2002). The River Witham in Lincolnshire is one such river, which has its levels altered twice yearly as part of a flood risk management strategy. Other rivers in Eastern England that are manipulated in a similar manner include the South Forty Foot drain, the Boston drains network, lower Hobhole drain, River Ancholme and River Welland. Rivers in South West England on the Somerset levels, such as the Huntspill River (see Langler & Smith, 2001) also have their water levels managed on a seasonal basis.

The effects of river management on fish populations centre around the impacts brought about by changes in river morphology, such as the lack of functional floodplains and associated lateral habitats that are required by fish to complete important stages in their life-cycles (Pinder, 1997). However, less emphasis has been placed on day-to-day or seasonal management

activities, such as the artificial manipulation of river levels. For example, weed cutting to reduce flood risk through improved conveyance, can reduce the density of zooplankton and spawning and nursery habitats (Mann, 1988) which can negatively impact fish growth and abundances (Garner *et al.*, 1996). However, there is a general paucity of evidence relating to the impact of routine river management actions on fish behaviour and ecology.

Home-range size (HRS) is used to describe and quantify the spatial behaviour of freshwater fish (see reviews by Northcote, 1978; Lucas & Baras, 2001). Hayne (1949) described the home-range as '*the area over which an animal normally travels*' and this was first applied to fish by Gerking (1950, 1953, 1959) following mark, release and recapture experiments. Linfield (1985) proposed that cyprinid communities are totally mobile, with their movements predominantly influenced by environmental conditions. However, site fidelity and home-ranges have been observed in many fishes including cyprinids (Lucas & Baras, 2001) for example; nase (Huber & Kirchhofer, 1998), dace (Clough & Ladle, 1997; Clough & Beaumont, 1998), roach (Baade & Fredrich, 1998), chub (Allouche *et al.*, 1999; Fredrich *et al.*, 2003), common bream (Lyons & Lucas, 2002), pike (Knight *et al.*, 2008 ; Knight *et al.*, 2009) and barbel (Baras & Cherry, 1990; Lucas & Batley, 1996). Seasonal differences in HRS have been observed (*e.g.* Huber & Kirchhofer 1998; Allouche *et al.*, 1999; Knight *et al.*, 2009) and other environmental factors, such as turbidity (*e.g.* Kuliskova *et al.*, 2009) and available habitat (Woolnough *et al.*, 2009) are also known to influence HRS.

The home-range concept has been applied to the ecological impact of habitat management (Kavanagh *et al.*, 2007; Knight *et al.*, 2009), including river corridor fragmentation (Woolnough *et al.*, 2009) on populations. Here, the home-range concept is used to study the impact of water level management on the behaviour of common bream in the lowland River Witham. The need for increased in-river flood storage capacity dictates that the level of the river is artificially dropped by 0.5 m in the autumn and raised again in the spring. The effect of this manipulation on common bream was assessed via an acoustic telemetry study that covered four episodes of artificial water level elevation and three episodes of artificial water level reduction. To my knowledge this is the first *in-situ* study of this type.

## **5.2 Materials and methods**

### **5.2.1 Study area**

The lower River Witham is described in sections 2.2.1, 2.2.9, 3.2.1 & 4.2.1. River levels are artificially lowered during the winter months to increase the river's flood storage capacity. During April to October the normal river level is maintained at approximately 1.5 m above Ordinance Datum Newlyn (ODN), and from November to March the level is reduced to and maintained at 1 m above ODN, during dry conditions. These manipulations are effected on the 1<sup>st</sup> April for spring manipulations, from winter level (1 m ODN) to summer level (1.5 m ODN), and on the 1<sup>st</sup> November for autumn manipulations, from summer level (1.5 m ODN) to winter level (1 m ODN). The timing of these transitions from one stable water level to another is

dependent on rainfall events and river flows, and as such this change is rarely achieved within 24 hours.


### **5.2.2 Sampling and tagging procedures**

Eighty-three adult common bream were tracked with acoustic telemetry, see above. River levels were gauged by stilling well at Kirkstead bridge (53°8'35"N; 0°14'36"W), approximately midpoint of the study area, by The Environment Agency hydrometry and telemetry systems.

### **5.2.3 Data handling and statistical analysis**

The calculation of HRS from fixed receiver data followed the approach of Crook (2004). Total linear ranges were estimated by manually determining the distance along the river channel between the outermost locations coordinates for each individual fish (Young, 1999; Ovidio *et al.*, 2000) to give the 100% HRS. In addition, 90% HRS which is suggested to provide a better indication of the core habitat size regularly used by each fish as rare excursive movements are excluded (Hodder *et al.*, 1998; Knight *et al.*, 2008) were estimated for each fish by calculating the minimum linear distance containing 90% of the observed locations of each individual fish (Crook, 2004). The spacing of acoustic receivers can also affect the spatial resolution of data. As receivers were spaced by 2-3 km, the spatial resolution might be compromised as some fish might travel past a receiver (and be detected), but not sufficiently far up- or downstream to be detected by the next receiver. Others may also venture beyond the outermost receivers. However, temporal resolution will be complete as receivers operate continuously. HRS was calculated on seasonal and monthly basis and for ten day periods before and after water level manipulations (see below). Only fish that had 100 or more locations (detections) in the sampling period (*e.g.* season, month, ten day pre- and post-level manipulations) were included in the analysis. Where data were combined on a seasonal basis, seasons were defined as follows; spring (March 21 – June 20), summer (June 21 – September 20), autumn (September 21 – December 20) and winter (December 21 – March 20).

To investigate the effects of water level manipulations on fish behaviour, HRSs were calculated for individual fish during a ten day period pre- and post-artificial water level manipulations. To assess if fish activity levels differed during these same ten day periods, total distance moved, calculated as the cumulative distance moved between receivers by an individual fish was determined. Seven water level manipulations occurred during the study, four elevations and three reductions. Manipulation from one stable water level to the resultant stable level often took a number of days (*e.g.* autumn 2007 took 5 days and spring 2008 took 18 days due to a rainfall event, Figure 5.1.). In these cases ten day sampling periods were selected as near to the point of level change as possible to capture periods of stable water level pre- and post-level manipulation (*i.e.* excluding the period of transition, Figure 5.1.).



**Figure 5.1** Water level (metres above Ordinance Datum Newlyn, m ODN) recorded at Kirkstead Bridge (midpoint of study area) during two of the seven artificial water level manipulations, illustrating the periods of flux in levels when changes are made and the selection of ten day sampling periods pre- and post-level manipulations. A; level manipulation in autumn 2007 (summer level to winter level), 5 days between sampling periods due to level flux, (i) = ten day sampling period pre-manipulation and (ii) = ten day sampling period post-manipulation. B; level manipulation in spring 2008 (winter level to summer level), 18 days between sampling periods due rainfall events, (iii) = ten day sampling period pre-manipulation and (iv) = ten day sampling period post-manipulation. Shaded areas represent periods of transition.

During three water level change events significant rainfall obscured changes, thus transition periods of 18, 24 and 8 days respectively were imposed (Table 5.1). The number of tagged fish experiencing the spring manipulations was greater than those experiencing the autumn manipulations, as firstly there were more spring events during the study and secondly because fish were tagged during the winter season and tags had a limited life, therefore many tags had expired or fish had left the study area by the time autumn manipulations occurred. Details of the numbers of fish used to generate HRS calculations and the number of days between ten day home-range calculation periods are given in Table 5.1.

Differences in HRS between all seasons and months, and gender differences were analysed with GLM ANOVA with *post-hoc* Tukey tests used to identify differences between seasons. Differences between adjacent months were analysed with two sample t-tests (significance level of 0.004 following Bonferroni correction) as datasets were made up from months in different years during the study when different fish were being tracked. Differences between the home-ranges and total recorded distance moved in ten day periods before and after artificial water level changes were possible using data from the same individual fish tracked pre- and post-level manipulations and were therefore analysed with paired t-tests (significance level of 0.008 following Bonferroni correction). For these analyses using paired data individual fish that were tracked during spring and autumn level manipulations in different years were identified and an average of results for each individual was used to avoid pseudo-replication. This reduced the sample size at spring manipulations from 60 to 56 and during autumn manipulations from 26 to 24. The relationship between mean monthly HRS and mean ‘monthly distance’ moved (see Chapter 4) was analysed with Pearson’s correlation. All means are given  $\pm$  one standard error.

**Table 5.1** The number of individual fish and the number of days between ten day home-range calculation periods used to determine the effects of seven independent artificial water level manipulations.


Level manipulation	Number of fish used in ten day home-range calculations	Number of days between ten day home-range calculation periods	Reason for gap between ten day sampling periods
Spring 2007	3	7	Natural run off
Autumn 2007	10	5	Natural run off
Spring 2008	19	18	Rainfall
Autumn 2008	12	24	Rainfall
Spring 2009	33	0	-
Autumn 2009	4	8	Rainfall
Spring 2010	5	0	-

### 5.3 Results


A two-way ANOVA on square root transformed data (to normalise the data) with season and gender as fixed factors revealed a significant effect of season on both the 100% HRS (GLM ANOVA,  $F_{3,307} = 18.00$ ,  $P < 0.0001$ ) and 90% HRS (GLM ANOVA,  $F_{3,307} = 10.74$ ,  $P < 0.0001$ ) with fish tending to have smaller HRSs in the autumn and winter months (Figure 5.2). A *post-hoc* Tukey test ( $P < 0.05$ ) revealed spring and summer home-ranges to be equivalent as were autumn and winter home-ranges, for both 100% and 90% estimations. Male fish had marginally larger 100% seasonal HRSs than females (mean HRS male =  $10.59 \pm 0.65$ , female =  $8.91 \pm 0.72$ ;  $F_{1,307} = 5.10$ ,  $p = 0.024$ ) although this was marginally non-significant when 90% HRSs were analysed (mean HRS male =  $6.51 \pm 0.50$ , female =  $5.49 \pm 0.54$ ;  $F_{1,307} = 3.13$ ,  $P = 0.078$ ). There was no interaction between month and sex for either the 100% or 90% HRSs ( $F_{3,304} = 0.39$ ,  $p = 0.76$  and  $F_{3,304} = 0.18$ ,  $P = 0.93$ , respectively). A similar analysis of monthly HRS (square root transformed) data revealed both 100% and 90% HRS to vary with month (GLM ANOVA,  $F_{11,682} = 17.08$ ,  $P < 0.0001$  and  $F_{11,682} = 11.09$ ,  $P < 0.0001$  respectively; Figure 5.3), with HRS consistently larger from April through to October than during the winter months. There was a significant effect of gender on both the 100% monthly HRS (male =  $6.67 \pm 0.32$ , female =  $5.87 \pm 0.37$ ,  $F_{1,682} = 7.04$ ,  $P = 0.008$ ) and the 90% monthly HRS (male =  $4.38 \pm 0.26$ , female =  $3.58 \pm 0.27$ ,  $F_{1,682} = 6.44$ ,  $P = 0.01$ ) but no interactions between month and sex for either HRS estimate ( $F_{11,671} = 0.51$ ,  $P = 0.89$  and  $F_{11,671} = 1.07$ ,  $P = 0.38$  respectively). Monthly HRS and 'Monthly distance' (common breem activity; see Chapter 4) were found to correlate for both 90% (Pearson correlation,  $r = 0.611$ ,  $df = 695$ ,  $P < 0.0001$ ) and 100% (Pearson correlation,  $r = 0.693$ ,  $df = 695$ ,  $P < 0.0001$ ), in effect fish had larger HRSs when activity levels were high.

#### 5.3.1 Effect of artificial water level manipulations

During spring manipulations (winter level increased to summer level) both 100% and 90% ten day home-ranges for individual fish increased significantly (paired t-test,  $t = 5.45$ ,  $P < 0.0001$ ;  $t = 4.93$ ,  $P < 0.0001$ ,  $df = 55$ , respectively); with home-ranges being significantly larger following the artificial increase in water levels. During autumn manipulations (summer level decreased to winter level); 100% and 90% ten day home-ranges for individual fish decreased significantly (Figure 5.4.; paired t-test,  $t = 5.00$ ,  $P < 0.0001$ ;  $t = 3.85$ ,  $P = 0.001$ ,  $df = 23$  respectively).




**Figure 5.2** Mean seasonal 100 % (dark grey) and 90% (light grey) home-ranges ( $\pm$ SE) for common bream in spring (n=92), summer (n=32), autumn (n=86) and winter (n=98). Common lettering indicate statistical equivalence following *post-hoc* Tukey test ( $P < 0.05$ ).



**Figure 5.3** Mean 90% home-ranges ( $\pm$ SE) for common bream in January (n=75), February (n=79), March (n=98), April (n=89), May (n=83), June (n=47), July (n=29), August (n=21), September (n=19), October (n=44), November (n=40), December (n=72). Months during winter level retention period are coloured light grey and months during summer level retention period are dark grey. Common lettering indicate statistical difference following two sample t-test (significance at  $< 0.004$  following Bonferroni correction) of adjacent month pairings around artificial water level manipulations (Table 5.2).

These ten day differences were also reflected in monthly differences (Table 5.2). Thus, between March and April, when water levels were increased, HRS increased significantly and between October and November when levels decreased, HRS decreased significantly. The increase and decrease in HRS following water level manipulation could reflect seasonal patterns in common bream activity. For example, Figure 5.3 shows that between February and April HRS increased (presumably due to increased common bream activity; see Chapter 4) and between October and December HRS decreased. Thus, it is possible the differences in HRS detected either side of the water level manipulations might be compounded by seasonal patterns of behaviour. To explore this, HRS in adjacent months prior to and post-water level manipulation were also compared (Table 5.2). For spring level manipulations there was a significant increase in HRS between February and March but not between April and May (Table 5.2). Also, in the autumn, when water levels were reduced, there was a significant decrease in HRS between October and November. No such decrease in HRS was detected between the months preceding this, September and October, nor the months following, November to December.

During spring manipulations (winter level to summer level) the total recorded distance moved by individual fish during the ten days pre- and post-artificial water level manipulations was marginally non-significant (paired t-test,  $t = 1.97$ ,  $df = 55$ ,  $P = 0.054$ ), whilst the total recorded distance moved pre- and post-autumn manipulation (summer level to winter level) were statistically equivalent (paired t-test,  $t = -0.09$ ,  $df = 23$ ,  $P = 0.929$ ; Figure 5.5). Thus, changes in water level significantly affected HRSs, but did not statistically affect activity.




**Figure 5.4** Mean 100% (solid symbols, solid line) and 90% (open symbols, broken line) home-ranges ( $\pm$ SE) for individual common bream tracked during the ten days pre- and post-artificial water level manipulations. Level manipulations occurred bi-annually in spring ( $n=56$ ; winter level to summer level) and autumn ( $n=24$ ; summer level to winter level).



**Table 5.2** Two sample t-test results for adjacent month pairings around artificial water level manipulations. Levels are manipulated biannually in spring (1<sup>st</sup> April) and autumn (1<sup>st</sup> November). Asterisks denote significance at < 0.004 following Bonferroni correction.

Home range estimate	Spring water level manipulation			Autumn water level manipulation		
	February – March	March – April	April – May	September – October	October – November	November – December
100%	Mean home ranges	Feb = 2.55 Mar = 5.32	Mar = 5.32 Apr = 9.55	Apr = 9.55 May = 9.55	Sept = 10.26 Oct = 10.41	Oct = 10.41 Nov = 3.74 Dec = 3.10
	t-test results	t = -3.28, df = 162, P = 0.001*	t = 4.00, df = 172, P < 0.0001*	t = 1.70, df = 164, P = 0.091	t = 0.11, df = 47, P = 0.914	t = 6.11, df = 71, P < 0.0001*
90%	Mean home ranges	Feb = 1.17 Mar = 3.75	Mar = 3.75 Apr = 6.07	Apr = 6.07 May = 5.10	Sept = 6.54 Oct = 6.60	Oct = 6.60 Nov = 2.71 Dec = 1.82
	t-test results	t = -3.95, df = 134, P < 0.0001*	t = 4.00, df = 172, P < 0.0001*	t = 1.10, df = 167, P = 0.271	t = 0.05, df = 50, P = 0.960	t = 3.94, df = 71, P < 0.0001*



**Figure 5.5** Mean total recorded distance moved ( $\pm$ SE) for individual common bream tracked during the ten days pre- and post-artificial water level manipulations. Level manipulations occurred bi-annually in spring (n=56; winter level to summer level) and autumn (n=24; summer level to winter level).

## 5.4 Discussion

### 5.4.1 Home-range size

The spatio-temporal scales at which fish utilise their habitats are of importance for meeting their resource requirements, in terms of both habitat and foraging and consequently are important to individual and population fitness (Hansen & Closs, 2005). Habitat quality will influence day-day and seasonal home range sizes as fish require certain habitat types on both of these scales. Bruylants *et al.* (1986) individually marked and recaptured perch from the Kleine Nete, Belgium, in two distinct habitat types; homogeneous sections with regard to depth, bottom type and current speed; and heterogeneous sections with a riffle pool sequence. Perch in the homogeneous sections were mobile, while fish in the heterogeneous sections were more likely to occupy restricted home ranges. A tendency was also observed for older larger perch to occupy home ranges as opposed to smaller younger fish which were more mobile. Prey abundance / availability will also influence home range size. Letourneur (2000) demonstrated that the foraging area and territory size of benthic feeding damselfish *Stegastes nigricans* (Lac.) were smaller when food supplies were high in summer, but increased in the winter when food supplies were lower.

HRSs varied with season being larger during the spring and summer months in comparison to the autumn and winter. Mean monthly home-range estimates also correlated with mean 'monthly distance' moved, as a measure of fish activity (see Chapter 4) indicating that during the warmer summer months when common bream were more active they also occupied larger home-ranges. This is consistent with other studies of cyprinids in temperate riverine ecosystems. Allouche *et al.*, (1999) radio tracked eight chub with activity sensitive tags, six in the summer and two in the winter, in the French River Rhône, and found fish to use smaller home-ranges, be less mobile and less active during winter. Kuliskova *et al.* (2009) also used radio telemetry to track seventeen ide for one year in the River Elbe, Czech Republic, and found home-ranges to be equivalent in spring, summer and autumn but significantly lower during the winter. Huber & Kirchhofer (1998) reported smaller HRSs in autumn and winter after radio tracking 21 nase in the River Aare, Switzerland, for between 15-448 days. These activity patterns are most likely the result of reduced metabolism by poikilothermic animals during the winter (Wieser, 1991; Huber & Kirchhofer, 1998). However, in the present study water level manipulations were also shown to influence HRS, confounding the observed seasonal effects.

The HRSs observed here were larger than those reported elsewhere for common bream. The 100% HRS over the duration of the study of nine common bream implanted with acoustic transmitters was reported as 0.35 to 5.4 km over a 55 day summer study on a 7.6 km stretch of the River Trent, UK (Lyons & Lucas, 2002). However, their study area was bounded by physical barriers, limiting the maximum home-range possible (see also Woolnough *et al.*, 2009). The duration of study may also result in differences in HRS; the longer the study, the further fish are likely to travel. Langford (1981) used acoustic telemetry to manually track both introduced and indigenous common bream in a 5 km heterogeneous reach of the River Thames and a 13.5 km homogeneous reach of the lower River Witham (upstream of the study area here),

UK. In the River Witham 100% HRS over the duration of the study varied from 3.5 km to 11.7 km, with mean home-ranges of 8.6 km and 8.8 km for introduced and indigenous common bream respectively. In the River Thames home-ranges were smaller, averaging 2.33 km, reflecting the smaller study area, although the Witham common bream used an average of 65% of the available reach length, while in the Thames common bream used only 46%, possibly reflecting differences in the quality of the two habitats.

The data also suggest a significant effect of gender on HRS with males tending to have larger home ranges than females. This is the first study to my knowledge which demonstrates a gender differences in the HRS of fish; difficulties in external (non-destructive) sex determination of most fish species have limited the study of gender differences in HRS (*e.g.* Gilroy *et al.*, 2010). Where gender has been determined, HRS has seldom been the focus of the study. However, Knight *et al.* (2008) found no significant difference between male and female pike for either seasonal HRS or mean annual excursion distance (excursions were defined as any significant movement outside the individual's home range), although sample sizes were low. This reveals another problem in studying gender differences in HRS of fish, that of sample size. Most telemetry studies tag and track relatively few individuals, therefore sample sizes are further reduced when splitting data by gender. For example, Lyons & Lucas (2002) tracked nine common bream in the River Trent, England, but only one of these fish was male, hindering efforts at making gender based comparisons. Lucas & Batley (1996) were able to determine the gender of radio tracked barbel (River Nidd, England) during the tagging procedure; a dental mirror was used to examine the gonads through the incision made for tag insertion. No gender differences in activity levels were observed, but some behavioural differences were observed, males tended to linger in the vicinity of spawning sites after upstream spawning migrations, whereas females soon return downstream once spawning was complete.

#### **5.4.2 Effects of artificial water level management**

Water level affected HRS; when levels were reduced in the autumn HRS decreased, whilst in the spring when levels were increased HRS increased. Water level manipulations occurred during the spring and autumn seasons when fish metabolism is increasing and decreasing respectively (Wieser, 1991). Thus, changes in HRS could reflect seasonal differences in common bream activity and not necessarily represent a consequence of changes in water level. However, water levels changed over a relatively short period of time so the differences observed are unlikely to be related to seasonal changes in photoperiod or temperature. Activity levels of common bream pre- and post-water level manipulations measured as total recorded distance moved were statistically equivalent, although this was marginal in the spring which may reflect an increase in common bream activity in April (see Chapter 4). This suggests the differences in HRS pre- and post-water level manipulation were not an artefact of differences in activity, although with this protocol of extensive observations during periods of environmental change it is possible that differences in HRS could reflect seasonal changes in biotic factors such as prey

availability. Thus, to ascertain cause and effect of water level manipulation on HRS requires experimental manipulation of water levels throughout the year.

Although the impacts of river channel modification are well documented (see Gregory, 2006), this is the first account of a flood risk management practice to affect fish behaviour. However, changes in behaviour do not necessarily equate to a negative impact on fitness, although reduced HRS does have implications for habitat availability and the number of competitive, predatory and parasitic interactions encountered (Woolnough *et al.*, 2009). Woolnough *et al.* (2009) performed a meta-analysis of 71 studies from 66 species of fish from around the world, showing that home-range estimates increased with body size and available water environment size. Water level manipulation of the River Witham reduced the volume of the water body; river width is reduced by ~10 m (~25%) and depth by 0.5 m, equating to ~12% reduction at the downstream limit of the study area and ~25% reduction at the upstream limit. Our study lends support to the notion that HRS is constrained by the extent of the available habitat (Woolnough *et al.*, 2009). However, within our study the length of the linear habitat remained unchanged, thus differences in HRS appear to arise as a result of volumetric changes in the amount of habitat.

Habitat availability and foraging opportunities are likely to be the most important implications of the behavioural changes brought about by these water level manipulations. Also, winter water levels restrict the availability of lateral habitat, as some tributaries are too shallow at winter levels to be accessible by common bream and other fishes, thus reducing the quantity of habitat available for refuge and over-wintering.

## Chapter 6 Social Networks

### 6.1 Introduction

Animals exist in populations and while members of a population may all appear to be similar, genetic and phenotypic differences exist that isolate them as individuals. These genetic, phenotypic and genotype-by-phenotype interactions can produce consistency in behavioural style. Behavioural style may be considered the animal's personality, although definition of personality, as related to animals, is problematic because of its anthropomorphic nature. Thus, some authors prefer temperament and/or behavioural syndrome (Gosling, 2008). Five main aspects of animal personality have been identified as behavioural syndromes in fishes (personality and behavioural syndrome are used interchangeably and appear to be the same thing); these are shyness-boldness, exploration-avoidance, activity, aggressiveness and sociality (Conrad *et al.*, 2011). Boldness is a measure of an individual's reaction to perceived danger and has been measured in many ways, such as the latency to approach a novel object (Frost *et al.*, 2007) or the time taken to resume feeding following disturbance (Pike *et al.*, 2008). Boldness may reflect an individual's risk taking behaviour such as foraging under the threat of predation. Shyness-boldness is likely to represent a continuum rather than a simple dichotomy and an individual's boldness may be plastic, changing in relation to the boldness of other fish that it associates with (Magnhagen & Saffan, 2005). Exploration-avoidance behaviours relate to an individual's willingness to investigate novel environments, food items or objects (Conrad *et al.*, 2011), whilst activity relates to the amount of movement measured in a safe and familiar environment and may depend on dominance rank and feeding strategy (Conrad *et al.*, 2011). Aggression occurs in contexts ranging from competition to territory defence and is likely to be plastic, being affected by various factors such as season and also the costs/benefits of being territorial which could change in relation to population density (Conrad *et al.*, 2011). Finally sociality is measured by quantifying an individual's response to conspecifics or their cues, with asocial individuals demonstrating avoidance and social individuals displaying an attraction to conspecifics (Reale *et al.*, 2007; Conrad *et al.*, 2011).

Analysis of when and how often individuals within a population interact with one another provides a method to study the social organisation of animals (Krause *et al.*, 2009; Makagon *et al.*, 2012) with the potential to reveal ecologically significant aspects that would otherwise have remained hidden. Network analysis and social network theory offer valuable insights into how members of a population interact and how this may impact on fitness (Krause *et al.*, 2009; Sih *et al.*, 2009). For example, Canon Jones *et al.* (2010) used network analysis to study aggressive interactions and associated fin damage of farmed Atlantic salmon parr in a hatchery environment. Higher and more intense aggressive interactions were observed in fish restricted of feed and fish initiating aggressive interactions had less fin damage, gained more weight and attained more central positions within schools, whilst fish receiving aggression had more fin damage and gained less weight. In most populations of animals, individuals interact preferentially with some other individuals and rarely interact with others. Thus, social

interactions can be highly structured reflecting differences between individuals in their social behaviour and thus the degree to which individuals are central or on the periphery of the population network, and the formation of substructures within the network (Krause & Ruxton, 2002; Krause *et al.*, 2007; Croft *et al.*, 2008; Krause *et al.*, 2009; Makagon *et al.*, 2012). Social network structure can be governed by the age and gender of animals (Lusseau & Newman, 2004) and how the different genders within a population interact can have reproductive implications. For example, males may interact to establish dominance relations which later govern access to females during the breeding season, thus male-male interactions may govern male-female interactions (Sih *et al.*, 2009). Similarly males may interact to establish breeding territories which once established attract breeding females.

While some animal behaviours are innate, some are learnt. Those behaviours which are more habitat/location specific are more likely to be the ones that have to be learned, either by an individual (*e.g.* trial and error) or learnt from another con-specific. With regard to the latter, social interactions represent an important pathway for such information exchange, which can impact on mate choice, alternative mating tactics, competition, cooperation, reciprocal altruism, dominance hierarchies, disease / parasite transmission and cooperative anti-predator behaviour (Croft *et al.*, 2004; Sih *et al.*, 2009). Long lived animals such as gorillas *Gorilla gorilla* (S.), red deer *Cervus elaphus* (L.), elephants *Loxodonta africana* (B.) and bottlenose dolphins *Tursiops truncatus* (M.) rely on information transfer to exploit their habitat (Janik 2000; Conradt & Roper 2003; Lusseau 2003), as do short lived animals, including fish, for example three-spined stickleback *Gasterosteus aculeatus* L. (Pike *et al.*, 2008) and have a capacity to learn behaviours from conspecifics.

Fish use learning and memory to orientate within their environments, being predisposed to learn specific associations at specific times or places for particular navigational problems they are likely to encounter (Odling-smee & Braithwaite, 2003). Social transmission of information plays a major role in determining how animals obtain information about the environment and can influence movement, resource use and patterns of interaction (Danchin *et al.*, 2004; Webster *et al.*, 2013). Recent research in Canada, tracking Atlantic salmon smolts (juveniles migrating to sea for the first time) originating from five rivers through the Gulf of St Lawrence as far as the Strait of Belle Isle (between the northern tip of Newfoundland and mainland Canada), found that the timing of smolt migrations from the different rivers were synchronised with each other and synchronised with that of kelts (adults which had spawned in fresh water and were returning to the sea). This has led to an interesting suggestion that smolts may learn migration routes from kelts (NASCO, 2012).

Some of the earliest research into animal social networks was carried out on aquatic animals such as dolphins and guppies *Poecilia reticulata* P. (Krause *et al.*, 2009). Lusseau (2003) studied a population of bottlenose dolphins (64 individuals) in the Doubtful Sound, Fjordland, New Zealand (between 1995 and 2001). Every time a school of dolphins was encountered, each adult member was photographed; individuals were identified from natural markings on the dorsal fin, resulting in data on the composition of 1292 schools. This information was used to

construct a social network for the population which was characterized by the presence of ‘centres’ of associations, showing that not all individuals play an equal role in this society. These hubs were mainly adult females, with the exception of one adult male and seemed to comprise older individuals, that had many scars and that tended to be larger in size. These well-connected individuals are known as keystone individuals and have a disproportionately large effect on the group dynamics or function (Sih *et al.*, 2009). Studying the same dolphin population, Lusseau & Newman (2004), identified a number of sub-communities within the population that appeared to be the result of associations of individuals of similar age and gender. Sub-communities were linked by ‘broker’ individuals which appeared to play a crucial role in maintain cohesiveness of the dolphin community.

Croft *et al.* (2004) marked 123 adult guppies in the Arima River, Trinidad, with individually identifiable visible implant elastomer and released simultaneously into the pool in which they were captured. Individual shoals of guppies were then re-sampled by seine net every day for seven days and the individuals making up the shoal noted, after which a total of 101 marked fish had been recaptured at least once. This network was found to be highly structured with well connected keystone individuals and less connected individuals on the periphery. Persistent pairwise associations were also observed but only between females. These interactions are of particular interest because repeated associations are thought to be important in the development of familiarity, which can confer a number of important advantages such as reduced food competition and increased predator evasion (see Dugatkin, 1997; Griffiths, 2003). Furthermore, repeated associations between pairs are the basic requirement for the evolution of reciprocal altruism (Milinski, 1987). Further study of guppies and three-spined sticklebacks by Croft *et al.* (2005a & 2005b) revealed social network structure was influenced by body length and shoaling tendency, with individuals interacting more frequently with conspecifics of a similar body length and shoaling tendency. Well-connected individuals were also more likely to interact with other well-connected individuals.

Pike *et al.* (2008) individually marked three-spined sticklebacks that had been assessed for bold and shy behavioural types. These were held in aquaria of either all shy, all bold or mixed individuals, and fish interactions recorded every five minutes, by still photography over a 24 hour period. Each image was then assessed for interactions; if fish were within one body length (< 45 mm) they were deemed to be interacting. Behavioural differences such as how bold or shy an individual was were shown to influence network structure with bold individuals having fewer overall interactions than shy fish, and bold fish tending to interact evenly across all group members and being more likely to move their location within the tank, thus being more active overall. By contrast, shy fish tended to interact with a smaller number of preferential group members and had a reduced tendency to move their location within the tank (thus less active overall). Learning ability and hence information exchange has been shown to vary between bold and shy individuals. Sneddon *et al.*, (2003) investigated whether hatchery reared rainbow trout *Oncorhynchus mykiss* (W.) could be classified as either bold or shy individuals based on their behaviour. Since boldness and shyness affect the willingness to take risks, the amount of time

spent in the open area of the tanks was assessed. The two groups were then tested for their speed at acquiring a basic learning task. Bold fish demonstrated a quicker learning ability than shy fish.

Social network structure can be assessed with simple descriptive statistics that can be used to characterise structural components of the network and the position of chosen individuals in relation to others (Krause *et al.*, 2009; Makagon *et al.*, 2012). For example, an individual's 'degree' is the number of direct connections a node has with other network nodes (*i.e.* number of direct social partners an individual has; Wey *et al.*, 2008; Sih *et al.*, 2009; Makagon *et al.*, 2012) or the number of immediate neighbours a node (individual) has when visualising the network. Degree can be binary or weighted; binary degree (see Makagon *et al.*, 2012) simply tells how many nodes a node is connected with, thus if all individuals interact with all other individuals there would be no variation in binary degree. However, weighted-degree (sometimes referred to as strength) takes account of the number of interactions between nodes (Makagon *et al.*, 2012), thus producing a more variable metric in the situation described above. A node's 'clustering co-efficient' is a measure of the degree to which nodes tend to cluster together (Makagon *et al.*, 2012) or the tendency for the individual's social partners to also be socially connected with each other (Sih *et al.*, 2009). For example; if there are three individuals ('x', 'y' and 'z') and individual 'y' interacts with both 'x' and 'z', but 'x' and 'z' do not interact with each other, then 'y' has a binary-degree of 2 but no clustering co-efficient. However, if 'x' and 'z' interact once, then 'y' will still have a binary-degree of 2, but it will now have a low clustering co-efficient, and if 'x' and 'z' interact many times then their clustering co-efficient's will increase as would their weighted-degrees. Networks can also be presented in diagrammatic form with nodes (points) representing individuals (or groups, subgroups, behaviour, location etc.) and lines (or edges) representing the relationships (*e.g.* social interactions) between them (Makagon *et al.*, 2012).

Common bream are true social animals, which according to Pitcher (1979) behave normally only when in full sensory contact, employing both lateral line and vision simultaneously to maintain individual distance and shoal position. Visual, chemical (pheromone), sound and touch stimuli are used to maintain social groups. Pitcher (1979, 1983, 1986) defined the term 'shoal' to refer to any group of fish formed by social attraction and 'school' to refer to synchronised and polarised swimming groups. Social interaction amongst cyprinids is thought to be important with respect to predator-prey interactions, fright reaction, foraging/feeding, reproduction *e.g.* spawning territoriality; which has been observed in common bream (Pocin *et al.*, 1996) and migration/homing (Smith, 1991). There is also evidence of within shoal dominance hierarchies in cyprinids (Smith, 1991) including common bream (Pitcher, 1979). Pitcher (1979) observed behaviours of common bream in a tank arrangement that allowed fish to be observed when in full sensory contact (visual, lateral line & olfactory), visual and olfactory contact and visual contact only. He observed that both lateral line and vision were employed simultaneously to maintain individual distance and therefore shoal position. Lack of lateral line input resulted in greater synchrony with a neighbour and common bream in full sensory contact were more likely



to make forays away from each other. Pitcher (1979) concluded that lateral line, vision and olfaction are used in shoals (a social group of fish), but probably only lateral line and vision in the maintenance of schools (polarised synchronised groups). Krause (1993) studied the feeding rates of roach and chub in a shallow stream near Cambridge, UK and found roach at the front of the shoal had significantly higher feeding rates and also had a higher tendency to palagic feeding than roach at the back of the shoal which fed more on the bottom, linking shoal position and potentially social status with increased fitness potential. These fish were caught and half were deprived of food and the other half well-fed for 3 days in captivity. After release 36% of them joined their old shoal again. Individuals from the starved group occupied front positions significantly more often than well-fed fish, but after 2 days this difference disappeared. Indicating that shoal position may be associated with resource needs.

The study of social interactions amongst fish has been largely confined to relatively small species inhabiting small environments over small periods of time, due to the ease that these can be studied in both the laboratory and field settings (Hoare & Krause, 2003). Here, social networks are constructed for each of the seven groups of fish tracked using acoustic telemetry. To my knowledge this is the first study to construct and analyse social networks from telemetry data of wild fish in their natural environment. Temporal trends in individual weighted-degree are analysed as a measure of sociality and social network structure is examined in the context of individuals' gender, age, size and spatio-temporal behaviour. It was hypothesised that because males and females face different selection pressures, which are likely to differ at different times of the year (Andersson, 1994), male and female weighted-degree would differ and that this would depend on the time of year. Additionally, because the activity levels and home-range size (HRS) varied in relation to season and because common bream are known to become more aggregated in the autumn and winter months (see Chapters 4 & 5), weighted-degree was predicted to vary in relation to month. In light of the results of Croft *et al.* (2005a & 2005b) it was also predicted that sociality would vary with individual body size.

## **6.2 Materials and Methods**

Study area, sampling techniques and tagging procedure are described previously. In summary eighty-three adult common bream were tracked with acoustic telemetry in seven groups. These seven groups were treated as discreet units when investigating their social networks because fish from the different groups were often unable to interact with one another during the study due to the temporal spread of tagging and the lifespan of the acoustic transmitter.

## **6.3 Data handling and statistical analysis**

To assess social interactions between fish from each group, the number of days in each month that fish were recorded interacting was calculated. The definition of an interaction has usually been study-specific, used to estimate relationships by observed interactions or associations between individuals, ranging from behavioural events (such as grooming) to co-occurrence (Lusseau *et al.*, 2008). Thus here, like studies by Lusseau (2003), Croft *et al.* (2004) and Pike *et*

*al.* (2008), interactions were based on co-occurrence with an interaction defined as two fish detected at the same receiver within one minute of each other. Given the range of the receivers (up to 400 m in good conditions) an interaction might be recorded between two fish when they are several hundred metres apart, thus fish may be classed as interacting when in fact they may not be. However, given the constraints of the tracking technology used, this was deemed a suitable criterion in that the method should give an indication of interaction opportunities, if not actual interactions themselves. Data were analysed for each group until 80% of the group had either left the study area or their tags had expired, due to the bias of including fish/tags that did not have the opportunity to interact for these reasons. Details of the size and sampling period for each group are presented in Table 6.1.

Social networks were constructed for each group using the mean number of interaction days per month with ‘Netdraw’ as part of the Ucinet© software (Borgatti *et al.*, 2002). These diagrams illustrate social network metrics; symbols or nodes represent individuals and lines or edges represent interactions. Group social network structure was measured by the individual’s monthly weighted-degree (which takes account of the number of interactions between nodes; Makagon *et al.*, 2012). This was calculated for each individual on both a monthly basis (where possible) and across the entire study period (Table 6.1), using the Ucinet© software and was used as measures of each individual’s sociality during that period. During some months an individual’s weighted-degree could not be calculated *i.e.* when an individual did not interact with any other fish. Each individual’s spatio-temporal behaviour was quantified by calculating mean ‘monthly distance’ and mean monthly 90% & 100% HRS (Chapters 4 & 5) during the sampling period (Table 6.1). The effects of the time of year (*i.e.* month), group (*i.e.* tagging event, Table 6.1), time since tagging (number of months), gender, fork length, and ‘monthly distance’ on an individual’s monthly weighted-degree were analysed with GLM-ANOVA (with group as a random factor). Monthly HRS was not used in the model because ‘monthly distance’ and HRS co-vary (Chapter 5). Maximal models were constructed with the significance of the independent


**Table 6.1** Sample size and sampling periods for the analysis of social networks for each group.




Group (see Figure 6.1)	Number of individuals in group	Start of sampling period	End of sampling period	Length of sampling period (months)
A	7	November 06	April 07	6
B	7	February 07	April 08	15
C	10	October 07	May 08	8
D	19	December 07	June 08	7
E	15	October 08	September 09	12
F	13	December 08	June 09	7
G	12	February 09	July 09	6

variables assessed following stepwise deletion (starting with the higher order interaction terms). Minimal regression models were then used to confirm results on significant covariates. In addition, the effects of gender and fork length, and mean ‘monthly distance’, mean monthly 90% & 100% HRS on an individual’s overall (across the entire study period, Table 6.1) weighted-degree were analysed with GLM-ANOVA (with group as a random factor) and Pearson correlation respectively. The frequency distribution of individual overall (entire study period; Table 6.1) weighted-degree was tested for normality using the Anderson-Darling method.


## 6.4 Results

Social networks for each group (A-G Table 6.1) were constructed using the mean number of interaction days per month with Netdraw as part of the Ucinet© software (Borgatti *et al.*, 2002) and are presented in Figure 6.1. The mean monthly weighted-degree for each group is presented in Figure 6.2. Group had a non-significant effect on monthly weighted-degree (GLM-ANOVA;  $F_{6,587} = 1.30$ ,  $P = 0.25$ ) as did fork length (GLM-ANOVA;  $F_{1,587} = 1.04$ ,  $P = 0.31$ ), gender (GLM-ANOVA;  $F_{1,587} = 0.14$ ,  $P = 0.71$ ), time of year *i.e.* month (GLM-ANOVA;  $F_{11,587} = 1.38$ ,  $P = 0.18$ ) nor the interaction between gender and month (GLM-ANOVA;  $F_{11,587} = 0.82$ ,  $P = 0.62$ ). However, significant effects of both ‘monthly distance’ (GLM-ANOVA;  $F_{1,587} = 69.45$ ,  $P < 0.0001$ ) and time since tagging (GLM-ANOVA;  $F_{1,587} = 5.77$ ,  $P = 0.017$ ) were revealed. Minimal regression models also revealed significant effects of ‘monthly distance’ ( $r = 0.013$ ,  $t = 8.58$ ,  $P < 0.0001$ ) and time since tagging ( $r = -0.05$ ,  $t = 5.3$ ,  $P < 0.0001$ ). Thus, the sociality of the fish varied according to how long they had been tagged; fish were more sociable soon after tagging. Also there was an effect of ‘monthly distance’ moved, with more active individuals (those that moved over greater distances) being more social. The mean monthly weighted-degree over the 12-month cycle is presented in Figure 6.3 and the mean monthly weighted-degree since tagging event (months) is presented in Figure 6.4.











**Figure 6.1** Social network structure (mean number of interaction days per month) for the 7 groups (A-G Table 6.1) constructed with Netdraw (Ucinet©; Borgatti *et al.*, 2002). Connecting lines are weighted to indicate the number of interaction days. Females are indicated by open nodes, males by shaded nodes. Numbers next to nodes are the individual fish (tag) ID numbers.




**Figure 6.2** Mean monthly weighted-degree per group ( $\pm$  SE). Weighted-degree represents how many other individuals each individual is connected to, taking account of the number of interactions between individuals.


Analysis of individual fish's weighted-degree across the entire study revealed no effect of gender (ANOVA;  $F_{1,82} = 1.73$ ,  $P = 0.199$ ) nor fork length (ANOVA;  $F_{1,82} = 0.67$ ,  $P = 0.778$ ). Overall weighted-degree correlated with mean 'monthly distance' moved (Pearson correlation  $r = 0.449$ ,  $df = 83$ ,  $P < 0.0001$ ) but did not correlate with either mean monthly 90% HRS (Pearson correlation  $r = 0.000$ ,  $df = 83$ ,  $P = 0.999$ ) nor mean monthly 100% HRS (Pearson correlation  $r = 0.039$ ,  $df = 83$ ,  $P = 0.724$ ). Thus gender, fish size nor HRS appeared to affect sociality, but activity level did. To investigate the individual variation in the sociality of the tracked fish a frequency histogram of weighted-degree for all 83 common bream is presented as Figure 6.5. There was no evidence to suspect data were not normally distributed ( $n = 83$ ,  $AD = 0.487$ ,  $P = 0.218$ ). The distribution of the weighted-degree suggests that some fish are more social than others. However, because the frequency histogram does not depart from normal this simply represents either end of a continuum. Thus, there is no evidence for a dichotomy in sociality.



**Figure 6.3** Mean monthly weighted-degree over the 12-month cycle ( $\pm$  SE) for all groups.



**Figure 6.4** The influence of time from tagging event on the mean monthly weighted-degree for all groups ( $\pm$  SE).



**Figure 6.5** Frequency histogram of the overall weighted-degree for all fish.

## 6.5 Discussion

The networks presented in Figure 6.1 illustrate how some individuals are more social than others and that some fish pairings interact more often than others. Also, in every group every individual is directly connected to every other individual, this is likely to be an artefact of all fish in each group being caught, tagged and released together. The extent of sociality appears to follow a normal distribution, such that there was no evidence of discrete classes of social/non-social fish. Overall weighted-degree peaked in October; this is possibly an artefact of the results from groups C and E, which were both tagged in October and had high numbers of interactions in the first month after tagging. There was only one other group which had results for October, leading to this unusually high mean for this month. This is consistent with the time after tagging analysis. Sociality was positively related to the activity levels of the fish, with more active individuals being more socially connected. More active fish would have more opportunity to interact with conspecifics so that they are more social would be expected. However, HRS did not affect individual sociality, thus it appears that activity level, is more important in sociality as opposed to HRS. Therefore an active animal with a small HRS may be more social than a less active individual which occupies a larger HRS.

That sociality peaks in the autumn / winter months is expected as common bream are known to aggregate around these seasons offering more opportunity for social interaction. However, the observation that gender nor size influenced sociality might reflect a true lack of effect, or it might be obscured by other factors, including time of year and time since tagging event. Fish interacted more often shortly after tagging which could have simply been due to fish being in the same location and therefore having more opportunity to interact. This in itself demonstrates



a lack of strict shoal faithfulness, and while common bream are a recognised shoaling species, the composition of these shoals may be plastic given the overall size of the population present.

However, little evidence could be found to support the hypothesis that these groups of fish remained in each others' company for extended periods. This may also be due to the small sample size of tracked individuals within a large population; which could be hiding other interactions with fish not tracked. Hanneman & Riddle (2005) state that social network analysis tends to study whole populations by means of census rather than drawing on samples. Within the current study sampling occurs at two levels; firstly only a small sample of the population was tagged / tracked and secondly sampling only occurred when tagged fish were in range of the receivers. Thus, the number of tagged fish is a relatively small proportion of the overall population, which may be insufficient to draw conclusions on how these fish interact socially and by sub-sampling interactions at receivers it is possible that a substantial number of interactions outside the range of the receivers were missed. Lusseau *et al.* (2008) draw attention to this source of uncertainty and its consequences on results and analyses. However, in this respect my sampling strategy is similar to that used by Lusseau (2003) and Lusseau & Newman (2004) who sample interactions when dolphins are encountered within Doubtful Sound on boat trips, and as such the temporal and spatial coverage of my study probably cause less uncertainty than these previously mentioned studies in this particular regard.

Given the resolution of the study (a small sub-sample of the population, sampled at a relatively small number of locations) and the fact that there are many potential confounding variables, it is perhaps unsurprising that few factors or co-variants explained the observed variation in individual sociality.

## **Chapter 7 Conclusions, management implications and recommendations for further research.**

### **7.1 Rehabilitation of lowland rivers**

Studies show that fish preferring both flowing water (rheophilic) and still water (limnophilic) are both at risk in large rivers worldwide (Buijse *et al.*, 2002). The underlying cause for this continued environmental degradation and overexploitation of aquatic resources is the ever growing human population that places increased demands for agriculture, industry and power generation (Collares-Pereira & Cowx, 2004). However, legislation such as the EU Habitats Directive (92/43/EEC) and Water Framework Directive (2000/60/EC) are giving new impetus to the restoration of freshwater habitats. Although restoration to pristine conditions is a utopian view, improvement is possible and if carried out appropriately, fish conservation management will deliver wider environmental benefits and protect biodiversity for future generations (Collares-Pereira & Cowx, 2004). Therefore, restoration should focus on the dynamics and condition of the main channel, floodplain and tributaries as a whole to increase the diversity of ecotones that would have been present before human intervention. Without ecologically sound restoration of degraded systems, the remnants of endangered species rich riverine/floodplain environments will continue to decline (Ward *et al.*, 1999).

In-channel river restoration (see [www.therrc.co.uk](http://www.therrc.co.uk)) is a relatively common rehabilitation technique suited to the upper reaches of modified riverine ecosystems. By contrast, restoration schemes at the floodplain scale along lowland rivers are few (Buijse *et al.*, 2002) although such ambitious schemes can deliver increased benefit to fish populations (*e.g.* Grift, 2001). However, recommendations made here (see sections 7.2.4 & 7.2.5 below) are meant to represent realistic and achievable management actions aimed at increasing the value of existing habitat types to fish populations.

Restoration of riparian habitats is an effective rehabilitation tool in lowland rivers where embankments have disconnected river channels from their floodplains (Langer & Smith, 2001). Moulton *et al.* (2012) used electro-fishing to survey juvenile fish assemblages in four riparian mesohabitat types (spawning ground, concrete embankment, foreshore & semi-natural) in the heavily modified lowland River Yser, Belgium. Juvenile fishes preferred natural habitat types, whereas artificial embankments showed the lowest species richness, abundance and functional organisation of juvenile fish species. Restored riparian habitats were preferred to artificial embankments, with juvenile fish avoiding bare microhabitats favouring grassy, woody and reedy habitat types equally, emphasizing the importance of microhabitat diversity. While the reconnection of existing floodplain waterbodies such as gravel pits (Nunn *et al.*, 2007) and borrow pits (Rysava-Novakova *et al.*, 2009) have been studied as surrogates for lost backwaters and floodplain waterbodies, there is little scope for such schemes along the lower Witham due to the lack of such waterbodies adjacent to the river channel and the requirement to move levees to achieve such reconnections. However, if such areas were to become available their reconnection would also be a valuable management action. These environments are usually far

from ideal, lacking shallow water and cover, but can act as sub- habitats and improvements on the status quo, although they could be improved to contain such features through intervention or by design.

## **7.2 Conclusions**

### **7.2.1 Review of rivers, fish, human interventions, telemetry and common bream.**

- Migration is an essential aspect in the lives of freshwater fishes. The modification of rivers during recent history has restricted habitat availability, leading to reductions in fish numbers and biodiversity. Understanding the temporal - spatial behaviour of fishes is vital to the management and rehabilitation of modified rivers. Migration by coarse fish in lowland rivers is an important component of their life-cycle and to sustain stocks in a moderately natural state, river management must be sensitive to these requirements (Lucas, 1998).
- Modification of lowland riverine ecosystems for the purposes of navigation (impoundment and barriers), flood control (artificial levee construction), land drainage (artificially high peak flows) and flow regulation (modification of hydrograph) are prevalent. Changes in river morphology generally shift riverine ecosystems from lentic towards lotic environments (Welcomme, 1994) through habitat degradation and reduced longitudinal and lateral connectivity. The reduced ability of fish to exploit specialised habitats required for the completion of their life-cycles negatively impacts on fish populations and community structure (Pinder, 1997).
- Common bream are adapted to utilise lentic habitats for spawning, nursery and flow refuge, therefore river modification will impact this species which is common to lowland rivers in Europe.
- Telemetry offers unrivalled insights into the behaviour and habitat use of fishes with the potential to direct rehabilitation effort. Given the extent of river modification and its prevalence in the lower reaches of rivers throughout the world (Welcomme, 1994) and the information that can be gained from the use of telemetry techniques, there is a paucity of such research in lowland river environments.

### **7.2.2 The ecology of the lower River Witham**

- The lower River Witham is a heavily modified and managed riverine environment lacking meanders and a free flooding floodplain; both having importance for fish communities and ecology. Lateral habitat is available, although limited and this has been reduced in the past by the disconnection of tributary waterbodies. Understanding the functions of such habitats is essential to management.
- Currently the lower river is dominated by roach. Eurytopic species, like roach and perch, have less strict requirements for spawning habitats and are thus able to establish populations

in a wide range of waterbodies (Schiemer & Waidbacher, 1992; Nunn *et al.*, 2007) leading to their dominance in modified habitats (Wolter, 2001a). The dominance of roach and perch is common in modified lowland rivers in Europe (Wolter & Vilcinskas, 1997; Wolter, 2001a; Wolter, 2001b). However, specifically adapted floodplain species like common bream are present, but their behaviour prevents consistent representation in fishery/ecological assessments.

### **7.2.3 Effects of tagging and translocation**

- The tagging protocol used did not significantly affect the fish's behaviour immediately after tagging and therefore all telemetry data collected should be used in subsequent analyses.
- Translocation did not significantly affect the level of the activity (distance moved) of the translocated fish; however it did affect their linear range as fish display 'homing behaviour' back towards the original capture site. The most likely scenario is that fish follow olfactory cues.
- Capturing fish on the fringes of the study area may increase the likelihood of those fish leaving the study area and depleting experimental sample sizes.

### **7.2.4 Seasonal movements and shifts in habitat use**

- The results presented have implications locally and for wider lowland river management and rehabilitation schemes, as well as emphasising the importance of connectivity within riverine ecosystems. Thus the addition of fish passes to all barriers to migration, will allow fish, eels and elvers to access habitats upstream.
- Shallow tributaries provide valuable spawning habitats in lowland riverine ecosystems, which should be protected and possibly enhanced in terms of both quality and quantity. Two such tributaries, Branston and Metherringham delphs (Figure 7.1), of the lower River Witham have been made inaccessible to fishes. The reconnection of these channels to the main river channel would seem an obvious rehabilitation action to boost recruitment by providing additional spawning and nursery habitat to enhance the common bream population, and other species of fish.
- Deeper tributaries appear to provide both refuge habitat in the autumn and over-wintering habitat. Thus the management of water levels could be used to increase the value of lateral habitat; retaining the main river depth at summer levels throughout the whole year would increase the winter depth of the shallower tributaries, making them more accessible to common bream and other fishes during periods of cold weather and high main-channel flow.



**Figure 7.1** Metherringham delph which was disconnected from the main river in the 1990s. The delph could be reconnected to the lower river to provide additional spawning, refuge and nursery habitat. In this photograph the main river runs perpendicular to the delph, behind the position of view (authors own photo).

A more consistent depth (during normal flow conditions) would also improve the value of marginal habitats (see Riis & Hawes, 2002) throughout the whole reach and tributaries.

- More homogeneous riverine environments (*i.e.* the lower reach of the study area; zone 7) were seldom used by the tracked common bream. The Huntspill River, Somerset UK, which has suffered similar levee construction as the lower River Witham, has benefited from a reduction of bank gradients and the construction of off-channel bays. These measures increased the abundance and diversity of 0-group fishes by producing spawning, nursery and refuge sites for coarse fish assemblages (Langler & Smith, 2001). Potential sites for such habitat creation should be identified along this length of lower river, to provide shelter and spawning sites that could boost recruitment and survival of juvenile fishes (see also Mouton *et al.*, 2012).
- Increasing lateral migration opportunities for adult fish by reconnecting floodplain waterbodies (see Staas & Newman, 1994; Grift *et al.*, 2001; Nunn *et al.*, 2007) has benefited recruitment of riverine fish populations. Existing tributary habitat should be protected and the feasibility of reconnecting Branston and Metherringham delphs to the main river channel should be investigated to provide additional spawning and nursery habitat to enhance fish populations.

- Such targeted improvements will be necessary if riverine habitats are to meet ‘good ecological status/potential’ by 2027, as stipulated by the EU Water Framework Directive 2000/60/EC.

#### **7.2.5 Home range size and effects of artificial water level manipulations**

- Home-range size varied with season being larger during the spring and summer months in comparison to the autumn and winter months. Mean monthly home-range estimates also correlated with mean ‘monthly distance’ moved, as a measure of fish activity indicating that during the warmer summer months when common bream were more active they also occupied larger home-ranges.
- Water level change appeared to affect home-range size; when levels were reduced in the autumn home-range size decreased, whilst in the spring when levels were increased home-range size increased.
- The seasonal manipulation of water levels to increase in-river flood storage capacity appears to be a historic practice. Thus, river managers should carry out hydrological modelling to investigate if such actions really do benefit flood risk mitigation. If the practice is ineffective against flood risk mitigation there is reason to discontinue such practice as retaining the main river at summer levels throughout the whole year would increase the depths of tributaries during the winter, making them more accessible to common bream and other fishes during periods of cold weather and high main channel flow for over-wintering and refuge (see Chapter 5). Alternative strategies should be researched; rather than historical seasonal manipulation, levels could be lowered on an *ad-hoc* basis before forecast rain events, thus minimising behavioural impacts on resident fishes. On the River Witham some preliminary modelling has taken place which suggests that although there are implications for flood risk should the river be retained at summer level all year round, mitigation against this increased risk may be straight forward, however more detailed investigation is required (J. Brown, Environment Agency, pers. comm.). While there are a number of conflicting interests surrounding the water level management of the lower River Witham, the principle of a stable water level has gained support and is being investigated within the context of the Lower Witham Catchment Management Plan, which aims to meet Good Ecological Status/Potential as stipulated by the EU Water Framework Directive 2000/60/EC (Dr. H. Barber, Environment Agency, pers. comm.). Other affected catchments are recommended to follow a similar strategy.
- The current water level management regime prohibits the establishment of permanent marginal habitat vegetation, which is of great value to fish and aquatic wildlife. Un-manipulated water levels would aid the establishment of permanent marginal habitats which would benefit fishes and other aquatic wildlife. River management should strike a balance between allowing the most natural discharge pattern possible, thus helping to maintain



**Figure 7.2** The lower river at Chapel Hill at winter retention level with marginal habitat high, dry and exposed to frost. Varying water levels prohibit the establishment of permanent marginal habitat vegetation, which is of great value to fish and aquatic wildlife (authors own photo).

natural biological communities, whilst safe-guarding local conurbation and valuable agricultural land from damaging flooding events.

#### **7.2.6 Social networks**

- The study of social networks of the tracked common bream revealed that more active individual fish were found to be more socially connected, but home-range size did not significantly affect sociality. However, there was no evidence of long-term shoal faithfulness recorded amongst any of the seven groups tracked and the extent of sociality appears to follow a normal distribution, such that there was no evidence of discrete classes of social/non-social fish, but sociality represented a continuum rather than a dichotomy. Results may have proved inconclusive due to the relatively small proportion of the rivers' common bream population that were tagged and therefore had their interactions recorded.

## 7.2 Recommendations for further research

- Investigation of the habitat use of roach to provide data on areas for spawning and shelter during floods, similar to the data collected here, would provide an interesting comparison. Anecdotal reports and observations made during this study suggest roach have different behaviours from those of the common bream population. Roach are dominant in many lowland rivers and thus may have less specific habitat requirements than common bream. However, there is concern over roach stocks in many rivers which have led to artificial propagation programmes such as the Avon Roach Project (see [www.avonroachproject.co.uk](http://www.avonroachproject.co.uk)). Technological developments will see acoustic tags coming on the market in the near future of a much smaller size than previously available, to overcome the problem of tag size to battery life ratio that can be used with smaller species and / or age classes.
- The study here examined the movements and habitat requirements of adult common bream. When technology allows, similar studies should be performed on juvenile common bream to identify their requirements.
- The effects of water level manipulation on home-range size should be investigated more closely by the random manipulation of river levels throughout the year and home-range sizes measured accordingly. The principle of reduced water levels influencing home-range size could also be tested more simply in laboratory based artificial tanks.
- The study of social networks amongst common bream in lowland rivers should be carried out where all individuals in the population can be captured and tagged. This therefore lends itself to a small impounded study area, which would aid the study as fish would be in relatively close proximity therefore reducing the chance of fish being spatially despite and therefore unable to interact for long periods of time.
- A more in-depth study of how and when common bream use the tributaries for spawning should be carried out. With emphasis on what micro-habitat features of the tributaries and environmental variables are required for the successful spawning of common bream.



## References

- Aarestrup, K., Thorstad, E.B., Koed, A., Jepsen, N., Svendsen, J.C., Pedersen, M.I., Skov C. & Okland, F. (2008) Survival and behaviour of European silver eel in late freshwater and early marine phase during spring migration. *Fisheries Management and Ecology* **15**, 435-440.
- Aarts, B.G.W., Van den Brink, F.W.B. & Nienhuis, P.H. (2004) Habitat loss as the main cause of the slow recovery of fish faunas of regulated large rivers in Europe: The transversal floodplain gradient. *River Research and Applications* **20**, 3-23.
- Alabaster, J.S. & Robertson, K.G. (1961) The effects of diurnal changes in temperature, dissolved oxygen and illumination on the behaviour of roach (*Rutilus rutilus* L.), bream (*Abramis brama* L.) and perch (*Perca fluviatilis* L.). *Animal Behaviour* **9**, 187-192.
- Allouche, S., Thevenet, A. & Gaudin, P. (1999) Habitat use by chub (*Leuciscus cephalus* L.) in a large river, the French Upper Rhone, as determined by radio telemetry. *Archiv fuer Hydrobiologie* **145** (2), 219-236.
- Andersson, M (1994) *Sexual Selection*. Printon University Press, New Jersey, USA. 601 pp.
- Armstrong, J.D. & Herbert, N. A. (1997) Homing movements of displaced stream-dwelling brown trout. *Journal of Fish Biology* **50**, 445-449.
- Arnold, G. & Dewar, H. (2001) Electronic tags in marine fisheries research: a 30-year perspective. *Methods and Techniques in Fish Biology and Fisheries* **1**, 7-64.
- Baade, U. & Fredrich, F. (1998) Movement and pattern of activity of the roach in the river Spree, Germany. *Journal of Fish Biology* **52**, 1165-1174.
- Backiel, T. & Zawisza, J. (1968) *Synopsis of biological data on the bream: Abramis brama* (L.) Food and Agriculture Organization of the United Nations, Fish Biology Synopsis **No. 36**, 122 pp.
- Bagenal, T.B. (1973) *Aging of Fish*, Unwin Brothers, London, 240 pp.
- Balon, E.K (1975) Reproductive guilds of fishes: a proposal and definition. *Journal of the Fisheries Research Board of Canada* **32**, 821-864.
- Balon, E.K (1981) Additions and amendments to the classification of the reproductive styles in fishes. *Environmental Biology of Fishes* **6**, 377-389.

- Baras, E. & Philippart, J.C. (1989). Application du radio-pistage a l'etude ecoethologique du barbeau fluviatile (*barbus barbus*); problemes, strategies et premiers resultats. *Cahiers d'Ethologies appliquee* **9**, 467-494
- Baras, E. & Cherry, B. (1990). Seasonal activities of female barbel *Barbus barbus* (L.) in the River Ourthe (Southern Belgium), as revealed by radio tracking. *Aquatic Living Resources* **3**, 283–294.
- Baras, E. (1992) Etude des strategies d'occupation du temps et de l'espace chez le barbeau fluviatile, *Barbus barbus* (L.). *Cahiers d'Ethologie* **12**, 125-412.
- Baras, E. (1995) Seasonal activities of *Barbus barbus*: effect of temperature on time budgeting. *Journal of Fish Biology* **46**, 806-818.
- Baras, E. (1997) Environmental determinants of residence area selection by *Barbus barbus* in the River Ourthe. *Aquatic Living Resource* **10**, 195-206.
- Baras, E. (1998) Selection of optimal positioning intervals in fish tracking: an experimental study on *Barbus barbus*. *Hydrobiologia* **371/372**, 19-28.
- Bauer, C. (2005) Potential problems with removing scales before surgical transmitter implantation. *Journal of Fish Biology* **66**, 847-850.
- Bendall, B. & Moore, A. (2008) Temperature sensing telemetry - possibilities for accessing the feeding ecology of marine mammals and their potential impacts on returning salmonid populations. *Fisheries Management and Ecology* **15**, 339–345
- Biro, P.A. & Post, J.R. (2008) Rapid depletion of genotypes with fast growth and bold personality traits from harvested fish populations. *Proceedings of the National Academy of Sciences of the United States of America* **105(8)**, 2919-22.
- Bolland, J.D., Cowx, I.G. & Lucas, M.C. (2008) Movements and habitat use of wild and stocked juvenile chub, *Leuciscus cephalus* (L.), in a small lowland river. *Fisheries Management and Ecology* **15**, 401-407.
- Bolland, J.D., Nunn, A.D., Lucas, M.C. & Cowx, I.G. (2012) The importance of variable lateral connectivity between artificial floodplain waterbodies and river channels. *Rivers Research and Applications* **28**, 1189-1199.

- Borcherding, J., Bauerfeld, M., Hintzen, D. & Neumann, D. (2002) Lateral migrations of fishes between floodplain lakes and their drainage channels at the Lower Rhine: diel and seasonal aspects. *Journal of Fish Biology* **61**, 1154-1170.
- Borgatti, S.P., Everett, M.G. & Freeman, L.C. (2002) Ucinet for Windows: software for social network analysis. Harvard: Analytic Technologies. See <http://www.analytictech.com>
- Borja, A. & Elliott, M. (2007) What does 'good ecological potential' mean, within the European Water Framework Directive? *Marine Pollution Bulletin* **54**, 1559-1564.
- Bovet, J. (1992). Mammals. In: *Animal Homing* (ed: Papi, F.), Chapman & Hall, London, 321–362 pp.
- Breukelaar, A.W., Lammens, E.H.R.R., Klein Breteler, J.G.P. & I. Tatrai (1994) Effects of benthivorous bream (*Abramis brama*) and carp (*Cyprinus carpio*) on sediment resuspension and concentrations of nutrients and chlorophyll a. *Freshwater Biology* **32**, 113-121.
- Bridger, C.J. & Booth R.K. (2003). The effects of biotelemetry transmitter presence and attachment procedures on fish physiology and behaviour. *Reviews in Fisheries Science* **11(1)**, 13-34.
- Britton, J.R. & Shepherd, J.S. (2005) Biometric data to facilitate the diet reconstruction of piscivorous fauna. *Folia Zoologica-Praha* **54**, 193–200.
- Brookes, A. (1988) *Channelized Rivers: Perspectives for Environmental Management*. Wiley-Blackwell, Chichester, 342 pp.
- Bruno D.O. & Gerard C.P. (2002) Behaviour of a stream-dwelling fish before, during, and after high discharge events. *Transactions of the American Fisheries Society* **131**, 762-771.
- Buijse, A.D., Coops, H., Staras, M., Jans, L.H., Van Geest, G.J., Grifts, R.E., Ibelings, B.W., Oosterberg, W. & Roozen, F.C.J.M. (2002) Restoration strategies for river floodplains along large lowland rivers in Europe. *Freshwater Biology*, **47**, 889-907.
- Cain, M.L., Bowman, W.D. & Hacker, S.D. (2011) *Ecology*. Sinauer Associates, Inc., Sunderland USA, 648 pp.
- Canon Jones, H.A., Hansen, L.A., Noble, C., Damsgard, B., Broom, D.M. & Pearce, G.P. (2010) Social network analysis of behavioural interactions influencing fin damage

- development in Atlantic salmon (*Salmo salar*) during feed-restriction. *Applied Animal Behaviour Science* **127**, 139-151.
- Carle, F.L. & Strub, M.R. (1978) A new method for estimating population size from removal data. *Biometrics* **34**, 621-630.
- Carling, P. (1995) Implications of sediment transport for instream flow modelling of aquatic habitat. In: *The Ecological Basis for River Management* (eds: Harper, D. M. & Ferguson, A.J.D.) John Wiley & Sons, Chichester, 17-49 pp.
- Carlson, H.R. & Haight, R.E. (1972). Evidence for a homesite and homing of adult yellowtail rockfish, *Sebastes flavidus*. *Journal of Fisheries Research Board Canada* **29**, 1011-1014.
- Carter, K.M., Woodley, C.M. & Brown, R.S. (2011) A review of tricaine methanesulfonate for anaesthesia of fish. *Reviews in Fish Biology and Fisheries* **21**, 51-59.
- Chadd, R.P. (2010). Assessment of Aquatic Invertebrates. In: *Conservation Monitoring in Freshwater Habitats* (eds: Hurford, C., Schneider, M. & Cowx, I.) Springer, Netherlands 63–72 pp.
- Chapman, B.B., Hulthen, K., Brodersen, J., Nilsson, P.A., Skov, C., Hansson, L.-A. & Bronmark, C. (2012a) Partial migration in fishes: causes and consequences. *Journal of Fish Biology* **81**, 456-478.
- Chapman, B.B., Skov, C., Hulthen, K., Brodersen, J., Nilsson, P.A., Hansson, L.-A. & Bronmark, C. (2012b) Partial migration in fishes: definitions, methodologies and taxonomic distribution. *Journal of Fish Biology* **81**, 479-499.
- Chesters, R.K. (1980) Biological monitoring working party. The 1978 National Testing Exercise, *Department of the Environment, Water Data Unit, Technical Memorandum* **19**, Reading, UK.
- Clough, S. & Ladle, M. (1997) Diel migration and site fidelity in a stream-dwelling cyprinid, *Leuciscus leuciscus*. *Journal of Fish Biology* **50**, 1117-1119.
- Clough, S. & Beaumont, W.R.C (1998) Use of miniature radio-transmitters to track the movements of dace, *Leuciscus leuciscus* (L.) in the River Frome, Dorset. *Hydrobiologia* **371/372**, 89-97.

- Cochran, W.W. (1980) Wildlife telemetry. *Wildlife Management Techniques Manual* **4**, 507-520.
- Coles T.F., Wortley, J.S. & Noble, P. (1985) Survey methodology for fish population assessment within Anglian Water. *Journal of Fish Biology* **27**, (Suppl. A), 175-186.
- Collares-Pereira, M.J. & Cowx, I.G. (2004) The role of catchment scale environmental management in freshwater fish conservation. *Fisheries Management and Ecology* **11**, 303-312.
- Conrad, J.L., Weinersmith, K.L., Brodin, T., Saltz, J.B. & Sih, A. (2011) Behavioural syndromes in fishes: a review with implications for ecology and fisheries management. *Journal of Fish Biology* **78**, 395–435.
- Conradt, L. & Roper, T.J. (2005) Consensus decision making in animals. *Trends in Ecology & Evolution* **20**, 449-456.
- Cooke, S. (2008) Biotelemetry and biologging in endangered species research and animal conservation: relevance to regional, national, and IUCN Red List threat assessments. *Endangered Species Research* **4**, 165–185.
- Cooke, S.J., Hinch, S.G., Lucas, M.C. & Lutcavage, M. (2012) Chapter 18 - Biotelemetry and biologging. In: *Fisheries Techniques* (3<sup>rd</sup> edition; eds: Zale, A.V., Parrish, A., & Sutton, T.M.) American Fisheries Society, Bethesda, USA 819-860 pp.
- Copp, G. H. & Peñáz, M. (1988) Ecology of fish spawning and nursery zones in the flood plain, using a new sampling technique. *Hydrobiologia* **169**, 209-224.
- Copp, G.H. (1989) The habitat diversity and fish reproduction function of floodplain ecosystems. *Environmental Biology of Fishes* **26**, 1-27.
- Copp, G.H. (1990) Effect of regulation on 0+ fish recruitment in the Great Ouse, a lowland river. *Regulated Rivers* **5**, 251-263.
- Copp, G.H. (1997) Importance of marinas and off channel water bodies as refuges for young fishes in a regulated lowland river. *Regulated Rivers: Research and Management* **13**, 303-307.

- Copp, G.H. & Roche, K. (2003) Range and diet of Eurasian otters *Lutra lutra* (L.) in the catchment of the River Lee (south-east England) since re-introduction. *Aquatic Conservation: Marine and Freshwater Ecosystems* **13**, 65-76.
- Cowx, I. (1983) The biology of bream, *Abramis brama* (L.), and its natural hybrid with roach, *Rutilus rutilus* (L), in the River Exe. *Journal of Fish Biology*, **22** 631-646.
- Cowx, I.G. & Welcomme, R.L. (1998) *Rehabilitation of Rivers for Fish*. Fishing News Books, Blackwell Science, Oxford, 260pp.
- Crawley M.J. (2002) *Statistical Computing: an Introduction to Data Analysis Using S-Plus*. Chichester: John Wiley & Sons Ltd, 772 pp.
- Croft, D. P., Krause, J. & James, R. (2004) Social networks in the guppy (*Poecilia reticulata*). *Proceedings of The Royal Society B* **271** (Suppl.), S516-S519.
- Croft, D.P., James, R., Ward, A.J.W., Botham, M.S., Mawdsley, D. & Krause, J. (2005a) The structure of social networks and association patterns in fish. *Oecologia* **143**, 211-219.
- Croft, D.P., James, R., Ward, A.J.W., Botham, M.S., Mawdsley, D. & Krause, J. (2005b) Assortative interactions and social networks in fish. *Population Ecology* **143**, 211-219
- Croft, D.P., James, R. & Krause, J. (2008) *Exploring animal social networks*. Princeton University Press, Princeton.
- Crook, D.A. (2004) Is the home range concept compatible with the movements of two species of lowland river fish? *Journal of Animal Ecology* **73**, 353-366.
- Danchin, E., Giraldeau, L.A., Valone, T.J. & Wagner, R.H. (2004) Public information: from nosy neighbors to cultural evolution. *Science* **305**, 487-491.
- De Leeuw, J.J. & Winter, H.V. (2008) Migration of rheophilic fish in the large lowland rivers Meuse and Rhine, the Netherlands. *Fisheries Management and Ecology* **15**, 409-415.
- DEFRA (2012) *Waste water treatment in the United Kingdom – 2012; Implementation of the European Union Urban Waste Water Treatment Directive – 91/271/EEC*. Defra, London. [www.defra.gov.uk/publications/files/pb13811-waste-water-2012.pdf](http://www.defra.gov.uk/publications/files/pb13811-waste-water-2012.pdf)
- Duncan, A. & Kubecka, J. (1993) Hydroacoustic methods of fish surveys. Research and Development Note 196, National Rivers Authority, 136 pp.

- Dugatkin, L.A. (1997) *Cooperation among animals: an evolutionary perspective*. Oxford University Press, New York, 221 pp.
- Edmondson, W.T. (1991) *The uses of Ecology: Lake Washington and beyond*. University of Seattle Press, Seattle, USA, 331 pp.
- Ellis, T., James, J. D., Stewart, C. & Scott, A. P. (2004). A non-invasive stress assay based upon measurement of free cortisol released into the water by rainbow trout. *Journal of Fish Biology* **65**, 1233–1252.
- English, K.K., Koski, W.R., Sliwinski, C., Blakley, A., Cass, A. & Woodey, J.C. (2005) Migration timing and river survival of late-run Fraser river sockeye salmon estimated using radio telemetry techniques. *Transaction of the American Fisheries Society* **134**, 1342-1365.
- Environment Agency (undated) *Coarse fish biology and management*. Fishery advisory booklet, Kidderminster, UK. [http://www.environment-agency.gov.uk/static/documents/Leisure/Coarse\\_fish\\_advisory\\_booklet.pdf](http://www.environment-agency.gov.uk/static/documents/Leisure/Coarse_fish_advisory_booklet.pdf)
- Environment Agency (2003) *River habitat survey in Britain and Ireland: field survey guidance manual: 2003 version*. Environment Agency, Bristol. 74 pp.
- Environment Agency (2008) *River Witham Catchment Flood Management Plan: Final Report*. Environment Agency, Bristol, 933 pp.
- Erlinge, S. (1968) Food studies on captive otters *Lutra lutra* L. *Oikos* **19**, 259-270.
- Extence, C.A., Chadd, R. P., England, J., Dunbar, M. J., Wood, P. J. & Taylor E.D. (2011) The assessment of fine sediment accumulation in rivers using macro-invertebrate community response. *River Research and Applications* **29(1)**, 17-55.
- Feunteun, E. & Marion, L. (1994) Assessment of grey heron predation on fish communities: the case of the largest European colony. *Hydrobiologia* **279/280**, 327-344.
- Flemming, G. (2002) *Flood risk management*. Thomas Telford, London, 256 pp.
- Forbes, I. & Wheeler, A. (1997) A comparison of late eighteenth century and present-day fishery surveys on the River Witham, Lincolnshire, UK. *Fisheries Management and Ecology* **4**, 325–335.

- Fowler, J., Cohen, L. & Jarvis, P. (1998) *Practical statistics for field biology*. Wiley, London, 259 pp.
- Fredrich, F., Ohmann, B., Curio, B. & Kirschbaum, F. (2003) Spawning migrations of the chub in the River Spree, Germany. *Journal of Fish Biology* **63**, 710-723.
- Frost, A.J., Winrow-Giffen, A., Ashley, P.J. & Sneddon, L.U. (2007) Plasticity in animal personality traits: does prior experience alter the degree of boldness? *Proceedings of The Royal Society B* **274**, 333-339.
- Gallagher, S.P., Hahn, P.K.J. & Johnson, D.H. (2008) Redd Counts. In: *Salmonid Field Protocols Handbook: Techniques for Assessing Status and Trends in Salmon and Trout Populations*. (eds. David H. Johnson, D.H., Shrier, B.M., O'Neal, J.S., Knutzen, J.A. Augerot, X., O'Neil, T.A. & Pearsons, T.N.) American Fisheries Society, Bethesda, Maryland, USA. 197-234 pp.
- Garner, P., Bass, J.A.B. & Collett, G.D. (1996) The effects of weed cutting upon the biota of a large regulated river. *Aquatic Conservation: Marine and Freshwater Ecosystems* **6**, 21-29.
- Garner P., Bass, J.A.B. & Collett G.D. (1998) The effects of weed cutting upon the biota of a large regulated river. *Aquatic Conservation: Marine and Freshwater Ecosystems* **6**, 21-29.
- Gerking, S.D. (1950) Stability of a stream fish population. *Journal of Wildlife Management* **14**, 193-202.
- Gerking, S.D. (1953) Evidence for the concepts of home range and territory in stream fishes. *Ecology* **34**, 347-365.
- Gerking, S.D. (1959) The restricted movements of fish populations. *Biology Review* **34**, 221-242.
- Giles N. (1994) *Freshwater fish of the British isles*. Shrewsbury: Swan Hill Press, 187 pp.
- Gilroy, D.J., Jensen, O.P. Allen, B.C., Chandra, S. Ganzorig, B., Hogan, Z., Maxted, J.T. & Vander Zanden, M.J. (2010) Home range and seasonal movement of taimen, *Hucho taimen*, in Mongolia. *Ecology of Freshwater Fish* **19**, 545-554.



- Girvan, M. & Newman, M.E.J. (2002) Community structure in social and biological networks. *Proceedings of the National Academy of Sciences of the United States of America* **99**, 7821-7826.
- Goldspink, C.R., & Banks, J.W. (1971) A readily recognizable tag for marking bream *Abramis brama* (L.). *Journal of Fish Biology* **3**, 407-411.
- Goldspink, C.R. (1978) The population density, growth rate and production of bream, *Abramis brama*, in Tjeukemeer, The Netherlands. *Journal of Fish Biology* **13**, 499-517.
- Gosling, S.D. (2008) Personality in Non-human Animals. *Social and Personality Psychology Compass* **2**, 985–1001.
- Gozlan, R.E., Britton, J.R., Cowx, I. & Copp, G.H. (2010) Current knowledge on non-native freshwater fish introductions. *Journal of Fish Biology* **76**, 751-786.
- Gregory, K.J. (2006) The human role in changing river channels. *Geomorphology* **79**, 172-191.
- Griffiths, S.W. (2003) Learned recognition of conspecifics by fishes. In: *Fish are smarter than you think: learning in fishes*. (eds. Brown, C., Laland, K.N., Krause, J.) *Fish Fisheries (Special Edition)* **4**, 256-268.
- Grift R.E., Buijse A.D., van Densen W.L.T. & Klein Breteler J.P.G. (2001) Restoration of the river-floodplain interaction: benefits for the fish community in the River Rhine. *Archiv für Hydrobiologie Supplement* **135** (Large Rivers 12), 173-185.
- Groisman, P.Y., Karl, T.R., Easterling, D.R., Knight, R.W., Jamason, P.F., Hennesy, K.J., Suppiah, R., Page, C.M., Wibig, J., Fortuniak, K., Razuvaev, V.N., Douglas, A., Forland, E. & Zhai, P.M. (1999) Changes in the probability of heavy precipitation: important indicators of climate change. *Climatic Change* **42**, 243–283.
- Groisman, P.Y., Knight, R.W., Easterling, D.R., Karl, T.R., Hegerl, G.C. & Razuvaev, V.N. (2005) Trends in intense precipitation in the climate record. *Journal of Climate* **18**, 1326–1350.
- Hadderingh, R.H. & Bakker, H.D. (1998) Fish mortality due to passage through hydroelectric power stations on the Meuse and Vecht rivers. In: *Fish Migration and Fish Bypasses* (eds: Jungwirth, M., Schmutz, S. & Weiss, S.) Fishing News Books, Blackwell Sciences, London, 315-328 pp.

- Hanneman, R.A. & Riddle, M. (2005) Introduction to social network methods. Riverside, CA: University of California, Riverside. (Published at <http://faculty.uer.edu/~hanneman/>)
- Hansen, E.A. & Closs, G.P. (2005) Diel and home range size in relation to food supply in a drift-feeding stream fish. *Behavioural Ecology* **16**, 640-648.
- Harms, C.A. (2005) Surgery in fish research: common procedures and postoperative care. *Lab Anim* **34**: 28-34.
- Hartley, P.H.T. (1947) The natural history of some British freshwater fish. *Proceedings of the Zoological Society of London* **117**, 129-206.
- Hasler, A. D. & Wisby, W. J. (1958) The return of displaced largemouth bass and green sunfish to a 'home' area. *Ecology* **39**, 289-293.
- Halvorsen, M. & Stabell, O. B. (1990). Homing behaviour of displaced stream-dwelling brown trout. *Animal Behaviour* **39**, 1089–1097.
- Hayne, D.W. (1949) Calculation of size of home range. *Journal of Mammalogy* **30**, 1-18.
- Heape, W. (1931) *Emigration, Migration and Nomadism*. Heffer & Sons Ltd, Cambridge, 369 pp.
- Heermann, L. & Borcherdig, J. (2006) Winter short-distance migration of juvenile fish between two floodplain water bodies of the Lower River Rhine *Ecology of Freshwater Fish* **15**, 161–168.
- Hickley, P. & Dexter, K.F. (1979) A comparative index for quantifying growth in the length of fish. *Aquaculture Research* **10**, 147-151.
- Hickley, P. (1996) Fish population survey methods: a synthesis. In: *Stock Assessment in Inland Fisheries* (ed: Cowx, I.G.), Fishing News Books, Blackwell, Oxford, 3-10 pp.
- Henderson, H. F., Hasler A.D. & Chipman, G.G., (1966) An ultrasonic transmitter for use in studies of movements of fishes. *Transactions of the American Fisheries Society* **95**(4), 350-356.
- Hert, E. (1992) Homing and home-site fidelity in rock-dwelling cichlids (Pisces: Teleostei) of Lake Malawi, Africa. *Environmental Biology of Fishes* **33**, 229-237.

- Hoare, D.J. & Krause, J. (2003) Social organisation, shoal structure and information transfer. *Fish and Fisheries* **4**, 269-279
- Hodder, K.H., Kenward, R.E., Walls, S.S. & Clarke, R.T. (1998) Estimating core ranges: a comparison of techniques using the common buzzard (*Buteo buteo*). *Journal of Raptor Research*, **32**, 82–89.
- Holker, F. (2006) Effects of body size and temperature on metabolism of bream compared to sympatric roach. *Animal Biology* **56**, 23-37.
- Huber, M. & Kirchhofer, A. (1998) Radio telemetry as a tool to study habitat use of nase (*Chondrostoma nasus* L.) in medium-sized rivers. *Hydrobiologia* **371/372**, 309-319.
- Huckstorf, V., Lewin, W & Wolter, C. (2008) Environmental flow methodologies to protect fisheries resources in human-modified large lowland rivers. *River Research and Applications* **24**, 519-527.
- Huet, M. (1959) Profiles and biology of western European streams as related to fish management. *Transactions of the American Fisheries Society* **88**, 155-163.
- Humphries, P., Cook, R.A., Richardson, A.M. & Serafini, L.G. (2006) Creating a disturbance: manipulating slackwaters in a lowland river. *Rivers Research and Applications* **22**, 525-542.
- Huntington, T.G. (2006) Evidence for intensification of the global water cycle: review and synthesis. *Journal of Hydrology* **319**, 83–95.
- Jacobsen, L., Berg, S., Jepsen, N. & Skov, C. (2004) Does roach behaviour differ between shallow lakes of different environmental state? *Journal of Fish Biology* **65**, 135-147.
- Janik, V.M. (2000) Whistle matching in wild bottlenose dolphins (*Tursiops truncatus*). *Science* **289**, 1355-1357.
- Jellyman, D. (2009) A review of radio and acoustic telemetry studies of freshwater fish in New Zealand. *Marine and Freshwater Research* **60**, 321-327.
- Jepsen, N., Koed, A., Thorstad, E. B., & Etienne, B. (2002) Surgical implantation of telemetry transmitters in fish: how much have we learned? *Hydrobiologia* **483**, 239-248.

- Jepsen, N., Schreck, C., Clements, S. & Thorstad, E. B. (2003). A brief discussion on the 2% tag/bodymass rule of thumb. In: *Aquatic Telemetry: Advances and Applications. Proceedings of the Fifth Conference on Fish Telemetry held in Europe* (eds: Sepedicato, M. T., Lembo, G. & Marmulla, G.). Rome, FAO/COISPA. 295 pp.
- Jepsen, N., Mikkelsen, J. S. & Koed, A. (2008). Effects of tag and suture type on survival and growth of brown trout with surgically implanted tags in the wild. *Journal of Fish Biology* **72**, 594-602.
- Johnson, P.B & Hasler, A.D. (1977) Winter aggregations of carp (*Cyprinus carpio*) as revealed by Ultrasonic Tracking. *Transactions of the American Fisheries Society* **106**(6), 556-559.
- Johnson A.C., Acreman, M.C. Dunbar, M.J., Feist, S.W., Giacomello, A.M., Gozlan, R.E., Hinsley, S.A., Ibbotson, A.T., Jarvie, H.P., Jones, J.I., Longshaw, M., Maberly, S.C., Marsh, T.J., Neal, C., Newman, J.R., Nunn, M.A., Pickup, R.W., Reynard, N.S., Sullivan, C.A., Sumpter J.P., Williams, R.J. (2009) The British river of the future: How climate change and human activity might affect two contrasting river ecosystems in England. *Science of the Total Environment* **407**, 4787–4798.
- Junk, W.J., Bayley, P.B. & Sparks, R.E. (1989) The flood pulse concept in river-floodplain systems. *Canadian Special Publication of Fisheries and Aquatic Sciences* **106**, 110-127.
- Jungwirth, M. (1998) River continuum and fish migration – going beyond the longitudinal river corridor in understanding ecological integrity. In: *Fish Migration and Fish Bypasses* (eds: Jungwirth, M., Schmutz, S. & Weiss, S.) Fishing News Books, Blackwell Sciences, London, 3-18 pp.
- Kalpers, J., Palata, K., Clotuche, E., Baras, E., Libois, R.M., Philippart, J.C. & Ruwet, J.C. (1989). Application of radio-tracking to the survey of wild populations of vertebrate species (mammals, birds, fishes) in Belgium. In: *Biotelemetry X* (ed: Amalaner Jr., C. J.) Fayetteville, University of Arkansas, 23–35 pp.
- Kavanagh, R.P, Stanton, M.A. & Brassil, T.E. (2007) Koalas continue to occupy their previous home-ranges after selective logging in *Callitris–Eucalyptus* forest. *Wildlife Research* **34**, 94–107.
- Kell, L. (1991) A comparison of methods used for coarse fish population estimation. In: *Catch Effort Sampling Strategies* (ed: Cowx, I.G.) Fishing News Books, Blackwell, Oxford, 184-200 pp.

- Kennedy, M. & Fitzmaurice, P. (1968) The biology of bream *Abramis brama* (L.) in Irish waters. *Proceedings of the Royal Irish Academy*, Inland Fisheries Trust, Dublin, 72 pp.
- Klefoth, T., Pieterek, T. & Arlinghaus, R. (2012) Impacts of domestication on angling vulnerability of common carp, *Cyprinus carpio*: the role of learning, foraging behaviour and food preferences. *Fisheries Management and Ecology* **20**, 174–186.
- Knight, C.M., Gozlan, R.E. & Lucas, M.C. (2008) Can seasonal home range size in pike *Esox lucius* predict excursion distance? *Journal of Fish Biology* **73**, 1058–1064.
- Knight, C.M., Kenwood, R.E., Gozlan, R.E., Hodder, K.H., Walls, S.S. & Lucas, M.C. (2009) Home-range estimation within complex restricted environments: importance of method selection in detecting seasonal change. *Wildlife Research* **36**, 213–224.
- Knights, B.C. & Lasee, B.A. (1996). Effects of implanted transmitters on adult bluegills at two temperatures. *Transactions of the American Fisheries Society* **125**, 440–449.
- Krause, J (1993) The relationship between foraging and shoal position in a mixed shoal of roach (*Rutilus rutilus*) and chub (*Leuciscus cephalus*): a field study. *Oecologia* **93**, 356–359.
- Krause, J. & Ruxton, G.D. (2002) *Living in groups*. Oxford University Press, Oxford, 213 pp.
- Krause, J., Croft, D.P. & James, R. (2007) Social network theory in the behavioural sciences: potential applications. *Behavioural Ecology and Sociobiology* **62**, 15–27.
- Krause, J., Lusseau, D. & James, R. (2009) Animal social networks: an introduction. *Behavioural Ecology and Sociobiology* **63** 967–973.
- Kubecka, J. (1996) Use of horizontal dual-beam sonar for fish surveys in shallow waters. In: *Stock Assessment in Inland Fisheries* (ed: Cowx, I.G.) Blackwell, Oxford, pp. 165–178.
- Kubecka, J., & Duncan, A., (1998). Diurnal changes in fish behaviour in a lowland river monitored by dual-beam echo sounder. *Fisheries Research* **35**, 55–63.
- Kuliskova, P., Horky, P., Slavik, O. & Jones, J. I. (2009) Factors influencing movement behaviour and home range size in the *Leuciscus idus*. *Journal of Fish Biology* **74**, 1269–1279.
- Lacroix, G.L. & Voegeli, F.A. (2000) Development of automated monitoring systems for ultrasonic transmitters. In: *Advances in Fish Telemetry. Proceedings of the Third*

*Conference on Fish Telemetry in Europe* (eds: Moore, A & Russell, I.) CEFAS, Lowestoft, 37-50 pp.

Lammens, E.H.R.R. (1985) A test of a model for planktivorous filter feeding by bream *Abramis brama*. *Environmental Biology of Fishes* **13**, 289-296.

Lammens, E.H.R.R (1999) The central role of fish in lake restoration and management. *Hydrobiologia* **395/396**, 191-198.

Langler, G.J. & Smith, C. (2001) Effects of habitat enhancement on 0-group fishes in a lowland river. *Regulated Rivers: Research and Management* **17**, 677-686.

Langford, T. E. (1974) Trials with ultrasonic tags for the study of coarse fish behaviour and movements around power station outfalls. *Aquaculture Research* **5**, 3, 61-62.

Langford, T. E., Milner, A. G. P., Foster, D. J. & Fleming, J. M., (1979) The movements and distribution of some common bream (*Abramis brama* L.) in the vicinity of power station intakes and outfalls in British rivers as observed by ultra-sonic tracking. Laboratory note RD/L/N 145/78. Central Electricity Research Laboratories, Fawley: 24 pp.

Langford, T. E. (1981) The movement and distribution of sonic-tagged coarse fish in two British rivers in relation to power station cooling-water outfalls. In: *Proceedings of the 3<sup>rd</sup> International Conference on Biotelemetry* (ed: Lang, F.M.) University of Wyoming, 197-232 pp.

Law, F. (2002) Impacts of climate change and catchment dynamics. In: *Flood risk management* (ed: Flemming, G.) Thomas Telford, London, 169-180 pp.

Laughton, R. (1989) The movements of adult Salmon within the River Spey. *Scottish Fisheries Research Report* 41. ISSN 0308 8022.

Letourneur, Y. (2000) Spatial and temporal variability in territoriality of tropical benthic damselfish on a coral reef (Reunion Island). *Environmental Biology of Fishes* **57**, 377-391.

Lilja, J., Keskinen, T., Marjomaki, T.J., Valkeajarvi, P. & Karjalainen, J. (2003) Upstream migration activity of cyprinids and percids in a channel, monitored by a horizontal split-beam echosounder. *Aquatic Living Resource* **16**, 185-190.

Linfield, R.S.J. (1985) An alternative concept to home range theory with respect to populations of cyprinids in major river systems. *Journal of Fish Biology* **27**, 187-196.

- Lower, N., Moore, A., Scott A. P., Ellis T., James J. D. & Russell I. C. (2005) A non-invasive method to assess the impact of electronic tag insertion on stress levels in fishes. *Journal of Fish Biology* **67**, 1202-1212.
- Lucas, M.C. & Johnstone, A.D.F. (1990) Observations on the retention of intragastric transmitters, and their effects on food consumption, in cod, *Gadus morhua* L. *Journal of Fish Biology* **37**, 647-649.
- Lucas, M.C. & Batley, E. (1996) Seasonal movements and behaviour of adult barbel *Barbus barbus*, a riverine cyprinid fish: implications for river management. *Journal of Applied Ecology* **33**, 1345-1358.
- Lucas, M.C. & Frear, P.A. (1997) Effects of a flow-gauging weir on the migratory behaviour of adult barbel, a riverine cyprinid. *Journal of Fish Biology* **50**, 382-396.
- Lucas, M.C., Mercer, T., Batley, E., Frear, P.A., Peirson, G., Duncan, A. & Kubecka, J. (1998a). Spatio-temporal variations in the distribution and abundance of fish in the Yorkshire Ouse system. *The Science of the Total Environment* **210/211**, 437-455.
- Lucas, M.C., Thom, T.J., Duncan, A. & Slavik, O. (1998b) *Coarse fish migration occurrence, causes and implications*. Technical Report W152. Environment Agency, Bristol, 161 pp.
- Lucas, M.C., Mercer T., Armstrong J.D., McGinty, S. & Rycroft, P. (1999) Use of a flat-bed passive integrated transponder antenna array to study the migration and behaviour of lowland river fishes at a fish pass. *Fisheries Research* **44**, 183-191.
- Lucas, M.C., Mercer, T., Peirson, G. & Frear, P. (2000) Seasonal movements of coarse fish in lowland rivers and their relevance to fisheries management. In: *Management and Ecology of River Fisheries*. (ed: Cowx, I.G.) Oxford: Blackwell Science. 87-100 pp.
- Lucas, M. C. & Baras, E. (2001) *Migration of Freshwater Fishes*. Blackwell Sciences Ltd, Oxford. 420 pp.
- Lucas, M.C., Walker, L., Mercer, T. & Kubecka, J. (2002) A Review of Fish Behaviours Likely to Influence Acoustic Fish Stock Assessment in Shallow Temperate Rivers and Lakes. *R&D Technical Report: W2-063/TR/1*. Environment Agency, Bristol.
- Lusseau, D. (2003) The emergent properties of a dolphin social network. *Proceedings of The Royal Society B: Biological Sciences* **270**, S186-S188.

- Lusseau, D. & Newman, M.E.J. (2004) Identifying the role that animals play in their social networks. *Proceedings of The Royal Society B* (Suppl.) **271**, S477-S481.
- Lusseau, D., Whitehead, H. & Gero, S. (2008) Incorporating uncertainty into the study of animal social networks. *Animal Behaviour* **75**, 1809-1815.
- Lyons, J. & Lucas, M.C. (2002) The combined use of acoustic tracking and echosounding to investigate the movements and distribution of common bream (*Abramis brama*) in the River Trent, England. *Hydrobiologia* **483**: 265-273.
- Lyons, J., Hickley, P., & Gledhill, S. (2002) An evaluation of recreational fishing in England and Wales. In: *Recreational Fisheries: Ecological, Economic & Social Evaluation* (ed: Pitcher, T.), Blackwell Science, Oxford, 144-155 pp.
- Magnhagen, C. & Saffan, F. (2005) Is boldness affected by group composition in young-of-the-year perch (*Perca fluviatilis*)? *Behavioural Ecology and Sociobiology* **57**, 295-303.
- Makagon, M.M., McCowan, B. & Mench, J.A. (2012) How can social network analysis contribute to social behaviour research in applied ethology. *Applied Animal Behaviour Science* **138**, 152-161.
- Maitland, P.S. & Campbell, R.N.B (1992) *Freshwater Fishes of the British Isles*. Harper Collins, London, 368 pp.
- Malinin, L.K. (1971) Home range and actual paths of fish in the river pool of the Rybinsk reservoir. *Fisheries Research Board of Canada Translation Series* **2282**, 26 pp.
- Malmquist, B. & Rundle, S. (2002) Threats to the running water ecosystems of the world. *Environmental Conservation* **29**, 134-153.
- Mann, R.H.K. (1988) Fish and fisheries of regulated rivers in the UK. *Regulated Rivers: Research and Management* **2**, 411-424.
- Marciak, Z. (1972) Relationship of limnological features to growth of bream (*Abramis brama* L.) and its catches in Polish lakes. *Verhandlungen des Internationalen Verein Limnologie* **18**, 1182-1189.



- Masters, J.E.G., Welton, J.S., Beaumont, W.R.C., Hodder, K.H., Pinder, A.C., Gozlan, R.E. & Ladle, M. (2002) Habitat utilisation by pike *Esox lucius* L. during winter floods in a southern English chalk river. *Hydrobiologia* **483**, 185-191.
- Matondo, B.N., Ovidio, M., Phillippart, J. & Poncin, P. (2009) Hybridization behaviour between two common European cyprinid fish species – silver bream, *Blicca bjoerkna* and common bream, *Abramis brama* – in a controlled environment. *Animal Biology* **59**, 97-108.
- May, J. & Van Rossum, M.K. (1995) The Quick and the Dead: Fish Entrainment, Entrapment, and the Implementation and Application of Section 316(b) of the Clean Water Act. *The Vermont Law Review*, **20**, 373.
- McFarlane, G.A., Wydoski, R.S. & Prince E.D (1990) Historical review of the development of external tags and marks. *American Fisheries Society Symposium* **7**, 9-29.
- Mesing, C. L. & Wicker, A. M. (1986) Home range, spawning migrations, and homing of radio-tagged Florida largemouth bass in two central Florida lakes. *Transactions of the American Fisheries Society* **115**, 286–295.
- Milinski, M. (1987) TIT FOR TAT in sticklebacks and the evolution of cooperation. *Nature* **325**, 433-435.
- Molls, F. (1999) New insights into the migration and habitat use by bream and white bream in the floodplain of the River Rhine. *Journal of Fish Biology* **55**, 1187-1200.
- Moore, A., Russell, I.C. & Potter, E.C.E. (1990) The effects of interperitoneally implanted dummy acoustic transmitters on the behaviour and physiology of juvenile Atlantic salmon. *Journal of Fish Biology* **37**, 713-721.
- Moore, A., Cotter, D., Quayle, V., Rogan, G., Poole, R., Lower, N. & Privitera, L. (2008) The impact of a pesticide on the physiology and behaviour of hatchery-reared Atlantic salmon, *Salmo salar*, smolts during the transition from fresh water to the marine environment. *Fisheries Management and Ecology* **15**, 385-392.
- Mouton, A.M., Buysse, D., Stevens, M. Van der Neucker, T. & Coeck, J. (2012) Evaluation of riparian habitat restoration in a lowland river. *Rivers Research and Applications* **28**, 845-857.

- Mulcahy, D.M. (2011) Antibiotic use during the intracoelomic implantation of electronic tags into fish. *Reviews in Fish Biology and Fisheries* **21**, 83-96.
- Muhlfeld, C.C. & Marotz, B. (2005) Seasonal movement and habitat use by subadult bull trout in the Upper Flathead River system, Montana. *North American Journal of Fisheries Management* **25**, 797-810.
- NASCO (2012) *Atlantic salmon at sea: Findings from recent research and their implications for management*. North Atlantic Salmon Conservation Organization (NASCO), Edinburgh. 24pp.
- National Farmers' Union (2008) 'Farming in the East Midlands'  
<http://www.nfuonline.com/x324.xml>
- Neely, B. C. & Steffensen, K. D. (2010) A comparison of gastrically and surgically implanted telemetry transmitters in shovelnose sturgeon. *Fisheries Management and Ecology* **16**, 323-328.
- Nikol'skii, G.V. (1961) *Special Ichthyology*. (Published for the National Science Foundation, Washington). Jerusalem: Israel Program for Scientific Translations, 539 pp.
- Northcote, T.G. (1978) Migratory strategies and production in freshwater fishes. In: *Ecology of Freshwater Fish Production* (ed: Gerking, S.D.) Blackwell, Oxford, UK; 326–359 pp.
- Northcote, T.G. (1984) Mechanisms of fish migration in rivers. In: *Mechanisms of Migration in Fishes* (ed: McCleave, J.D., Arnold, G.P., Dodson, J.J. & Neil, W.H.) New York: Plenum Press, 317-355 pp.
- Northcote, T.G. (1998) migratory behaviour of fish and its significance to movement through riverine fish passage facilities. In: *Fish Migration and Fish Bypasses* (eds: Jungwirth, M., Schmutz, S. & Weiss, S.) Fishing News Books, Blackwell Sciences, London, 3-18 pp.
- Nunn, A.D., Harvey, J.P. & Cowx, I.G. (2007) Benefits to 0+ fishes of connecting man-made waterbodies to the lower River Trent, England. *Rivers Research and Applications* **23**, 361-376.
- Nunn, A.D., Copp, G.H., Vilizzi, L. & Carter, M. (2010) Seasonal and diel patterns in the migrations of fishes between a river and a floodplain tributary. *Ecology of Freshwater Fish* **19**, 153-162.

- O'Sullivan P.E. (1995) Eutrophication. *International Journal of Environmental Studies* **47**, 173-196.
- Odling-Smee, L. & Braithwaite, V.A. (2003) The role of learning in fish orientation. *Fish and Fisheries* **4**, 235–246.
- Ovidio, M, Philippart, J. & Baras, E. (2000) Methodological bias in home range and mobility estimates when locating radio-tagged trout, *Salmo trutta*, at different time intervals. *Aquatic Living Resource* **13**, 449-454.
- Peirson, G., Tingley, D. Spurgeon, J. & Radford, A. (2001) Economic evaluation of inland fisheries in England and Wales. *Fisheries Management and Ecology* **8**, 415-424.
- Penczak, T., Forbes, I., Coles, T. F., Atkins, T. and Hill, T. (1991) Fish community structure in the rivers of Lincolnshire and South Humberside, England. *Hydrobiologia* **211**, 1-9.
- Perrow, M.R, Meijer, M., Dawidowicz, P. & Coops, H. (1997) Biomanipulation in shallow lakes: state of the art. *Hydrobiologia* **342/343**, 355-365.
- Peter, A. (1998) Interruption of the river continuum by barriers and the consequences for migratory fish. . In: *Fish Migration and Fish Bypasses* (eds: Jungwirth, M., Schmutz, S. & Weiss, S.) Fishing News Books, Blackwell Science, 99-112 pp.
- Phillips, R. & Rix, M. (1985) *Freshwater Fish of Britain, Ireland and Europe*. Pan Books Ltd. London, 144 pp.
- Pike, T.W., Samanta, M., Lindstrom, J. & Royle, N.J. (2008) Behavioural phenotype affects social interactions in an animal network. *Proceedings of The Royal Society B: Biological Sciences* **275**, 2515-2520.
- Pilcher, M.W., Copp, G.H. & Szomolai, V. (2004) A comparison of adjacent natural and channelized stretches of a lowland river. *Biologia-Bratislava* **59**, 669-673.
- Pinder, L.C.V. (1997) Research on the Great Ouse: overview and implications for management. *Regulated Rivers: Research and Management* **13**, 309-315.
- Pinder, L.C.V., Marker, A.F.H., Mann, R.H.K., Bass, J.A.B. & Copp, G.H. (1997) The River Great Ouse, a highly eutrophic, slow-flowing, regulated, lowland river in Eastern England. *Regulated Rivers: Research and Management* **13**, 203-219.

- Pitcher, T.J. (1979) Sensory information and the organisation of behaviour in a shoaling cyprinid fish. *Animal Behaviour* **27**, 126-149.
- Pitcher, T.J. (1983) Heuristic definitions of shoaling behaviour. *Animal Behaviour* **31**, 611-613.
- Pitcher, T.J. (1986) Functions of shoaling behaviour in teleosts. In: *The Behaviour of Teleost Fishes* (ed: Pitcher, T.J.), Croom Helm, London, 294-337 pp.
- Pollux, B.J.A., Korosi, A., Verberk, W.C.E.P., Pollux, P.M.J. & van de Velde, G. (2006) Reproduction, growth and migration of fishes in a regulated lowland tributary: potential recruitment to the River Meuse. *Hydrobiologia* **565**, 105-120.
- Poncin, P., Philippart, J.C. & Ruwet, J.C. (1996) Territorial and non-territorial spawning behaviour in the bream. *Journal of Fish Biology* **49**, 622-626.
- Priede, I.G. & Swift, S.M. (1992) *Wildlife telemetry: Remote Monitoring and Tracking of Animals*. Ellis-Horwood, 708 pp.
- Radford, A.F., Hatcher, A. & Whitmarsh, D. (1991) *An economic evaluation of salmon fisheries in Great Britain*. A research report to the Ministry of Agriculture Fisheries and Food, 342 pp.
- Ramstad, K. M. & Woody, C. A. (2003) Radio tag retention and tag-related mortality among adult sockeye salmon. *North American Journal of Fisheries Management* **23**, 978-982.
- Raven, P.J., Holmes, N.T.H., Dawson, F.H., Fox, P.J.A., Everard, M., Fozzard, I.R. & Rouen, K.J. (1998) *River Habitat Quality: the physical character of rivers and streams in the UK and Isle of Man*. Environment Agency, Bristol. 96 pp.
- Reale, D., Reader, S.M., Sol, D., McDougall, P.T. & Dingemanse, N.J. (2007) Integrating animal temperament within ecology and evolution. *Biological Reviews* **82**, 291-318.
- Rice, S.P., Greenwood, M.T. & Joyce, C.B. (2001) Tributaries, sediment sources, and the longitudinal organisation of macroinvertebrate fauna along river systems. *Canadian Journal of Fisheries and Aquatic Science* **58**, 824-840.
- Riis, T. & Hawes, I. (2002) Relationships between water level fluctuations and vegetation diversity in shallow water of New Zealand lakes. *Aquatic Botany* **74**, 133-148.

- Rinchard, J. & Kestemont, P. (1996) Comparative study of reproductive biology in single- and multiple-spawner cyprinid fish: morphological and histological features. *Journal of Fish Biology* **49**, 883-894.
- Ross, L.G. & Ross, B. (2008) *Anaesthetic and Sedative Techniques for Aquatic Animals*, 3<sup>rd</sup> Edition. Blackwell Science, Oxford, UK, 221 pp.
- Schiemer, F. & Waidbacher, H. (1992) Strategies for conservation of a Danubian fish funa. In: *River Conservation and Management* (eds: Boon, P.J., Calow, P. & Petts, G.E.) Wiley, Chichester, 363-382 pp.
- Schiemer, F. (2000) Fish as indicators for the assessment of the ecological integrity of large rivers. *Hydrobiologia* **422/423**, 271-278.
- Schutz, U. & Berg, R. (1987) The migration of ultrasonic-tagged bream, *Abramis brama* (L.), in Lake Constance (Bodensee-Untersee). *Journal of Fish Biology* **31**, 409-414.
- Shikhshabekov, M.M. (1972) Annual cycle of the gonads of the bream *Abramis brama* (L.) in the lakes of the Terek river delta. *Voprosy Ikhtiologii* **12**, 193-197.
- Sibert, J.R., Lutcavage, M.E., Nielsen, A., Brill, R.W. & Wilson, S.G. (2006) Interannual variation in large-scale movement of Atlantic bluefin tuna (*Thunnus thynnus*) determined from pop-up satellite archival tags. *Canadian Journal of Fisheries and Aquatic Sciences* **63**, 2154-2166.
- Sih, A., Hanser, S.F. & McHugh, K.A. (2009) Social network theory: new insights and issues for behavioural ecologists. *Behavioural Ecology and Sociobiology* **63**, 975-988.
- Skov, C. J., Brodersen, J., Bronmark, C., L.-A. Hansson L.-A., P. Hertonsen, P., & Nilsson, P.A. (2005) Evaluation of PIT-tagging in cyprinids. *Journal of Fish Biology* **67**, 1195-1201.
- Smith, R.J.F. (1985) *The Control of Fish Migration*. Springer-Verlag, Berlin. 243 pp.
- Smith, R.J.F. (1991) Social behaviour, homing and migration. In: *Cyprinid Fishes: Systematics, Biology and Exploitation* (eds: Winfield, I.J. & Nelson, J.S.), Chapman and Hall, London, UK, 509-529 pp.

- Smith, G. W., Campbell, R. N. B. & MacLaine, J. S. (1998) Regurgitation rates of intragastric transmitters by adult Atlantic salmon (*Salmo salar* L.) during riverine migration. *Hydrobiologia* **371/372**, 117–121.
- Sneddon, L.U. (2003) The bold and the shy: individual differences in rainbow trout. *Journal of Fish Biology* **62**, 971–975.
- Staas, S. & Newman, D. (1996) The occurrence of larval and juvenile 0+ fish in the Lower River Rhine. *Archiv fur Hydrobiologie Supplement* **113**, 325–332.
- Steinmann, P., Koch, W. & Scheuring, L. (1937) Die Wanderung unserer Subwasserfische. Dargestellt auf Grund von Merkierungsversuchen. *Zeitschrift fur Fischerei und deren Hilfswissenschaften* **35**, 369–467.
- Svendsen, J.C., Koed, A. & Lucas, M.C. (2005) The angle of attack of the body of common bream while swimming at different speeds in a flume tank. *Journal of Fish Biology* **66**, 572–577.
- Svendsen, K.M., Renshaw, C.E., Magilligan, F.J., Nislow, K.H. & Kaste, J.M. (2009) Flow and sediment regimes at tributary junctions on a regulated river: impact on sediment residence time and benthic macroinvertebrate communities. *Hydrological Processes* **23**, 284–296.
- Trefethen-Parker, S. (1956) *Sonic equipment for tracking individual fish*. US Fish and Wildlife Service Special Science Report Fisheries No. 179, 11 pp.
- Turnpenny, A. (1998) Mechanisms of fish damage in low-head turbines: an experimental appraisal. In: *Fish Migration and Fish Bypasses* (eds: Jungwirth, M., Schmutz, S. & Weiss, S.) Fishing News Books, Blackwell Sciences, London, 301–314 pp.
- Tyus, H.M. (1985) Homing behaviour noted for Colorado squawfish. *Copeia* **1985**, 213–215.
- UK TAG (2008) Guidance on the Classification of Ecological Potential for Heavily Modified Water Bodies and Artificial Water Bodies. WFD UK TAG.  
[http://www.wfduk.org/sites/default/files/Media/Classification%20of%20ecological%20potential%20for%20HMWBs%20and%20AWBs\\_Final\\_310308TAG%20guidance.pdf](http://www.wfduk.org/sites/default/files/Media/Classification%20of%20ecological%20potential%20for%20HMWBs%20and%20AWBs_Final_310308TAG%20guidance.pdf)
- United States Environmental Protection Agency (1986) *Phosphorus: Water Quality Standards Criteria Summaries: A compilation of State/Federal Criteria*. Office of Water regulations and Standards, Washington, DC 20460. 24pp.

- Waidbacher, H.G. & Haidvogel, G. (1998) Fish migration and fish passage facilities in the Danube: past and present. In: *Fish Migration and Fish Bypasses* (eds: Jungwirth, M., Schmutz, S. & Weiss, S.) Fishing News Books, Oxford, 85-98 pp.
- Walker, T.J. & Hasler, A.D. (1949) Detection and discrimination of odours of aquatic plants by bluenose minnow (*Hyborhynchus notatus*, Raf.). *Physiological Zoology* **22**, 45-63.
- Wagner, G.N., Cooke, S.J., Brown, R.S. & Deters, K.A. (2011) Surgical implantation techniques for electronic tags in fish. *Reviews in Fish Biology and Fisheries* **21**, 71-81.
- Ward, J. V., Tocker, K. & Schiemer, F. (1999) Biodiversity of floodplain river ecosystems: ecotones and connectivity. *Regulated Rivers: Research and Management* **15**, 125-139
- Ward, J.V. & Stanford, J.A. (1995) Ecological connectivity in alluvial river ecosystems and its disruption by flow regulation. *Regulated Rivers: Research and Management* **10**, 159-168.
- Wieser W. (1991) Physiological energetics and ecophysiology. In: *Cyprinid Fishes: Systematics, Biology and Exploitation* (eds: Winfield, I.J. & Nelson, J.S.) Chapman and Hall, London, 426-455 pp.
- Webster, M.M., Atton, N., Hoppitt, W.J.E. & Laland, K.N. (2013) Environmental complexity influences association network structure and network-based diffusion of foraging information in fish shoals. *The American Naturalist* **181**, 235-244.
- Welcomme, R.L. (1985) *River fisheries*. FAO fisheries technical paper 262, Food and Agriculture organization of the United Nations, Rome.
- Welcomme, R.L. (1994) The status of large river habitats. In: *Rehabilitation of Freshwater Fisheries* (ed: Cowx, I.G.) Fishing News Books, Oxford, 11-20 pp.
- Westerberg, H. & Lagenfelt, I. (2008) Sub-sea powercables and the migration behaviour of the European eel. *Fisheries Management and Ecology* **15**, 369-375.
- Wey, T., Blumstein, D.T., Shen, W. & Jordan, F. (2008) Social network analysis of animal behaviour: a promising tool for the study of sociality. *Animal Behaviour* **75**, 333-344.
- Wheeler, A. (1978) *Key to the fishes of Northern Europe*. Frederick Warne, London, 430 pp.

- Wheeler, W.H. (1990) *A History of the Fens of South Lincolnshire* (ed: Simmons, B.) Stamford: Paul Watkins, 489 pp.
- Whelan, K. F. (1983) Migratory patterns of bream *Abramis brama* L. shoals in the River Suck system. *Irish Fisheries Investigations*, Series A. **23**, 11-15.
- Wieser, W. (1991) Physiological energetics and ecophysiology. In: *Cyprinid Fishes: Systematics, Biology and Exploitation* (eds: Winfield, I.J. & Nelson, J.S.), Chapman and Hall, London, UK, 426-455 pp.
- Winfield, I.J., Peirson, G., Cryer, M. And Townsend, C.R. (1983) The behavioural basis of prey selection by underyearling bream (*Abramis brama* (L.)) and roach (*Rutilus rutilus* (L.)). *Freshwater Biology* **13**, 139-149.
- Winter, J.D. (1983). Underwater biotelemetry. In: *Fisheries Techniques* (eds: Nielson, L.A. & Johnson J.D.) American Fisheries Society, Bethesda, Maryland, 371-395pp.
- Winter J.G. (1996) Advances in underwater biotelemetry. In: *Fisheries Techniques* (2<sup>nd</sup> edition; eds: Murphy, B.R. & Willis, D.W.) Bethesda, MD: American Fisheries Society, 555–590 pp.
- Winter, H.V. & Fredrich, F. (2003) Migratory behaviour of ide: a comparison between the lowland rivers Elbe, Germany, and Vecht, The Netherlands. *Journal of Fish Biology* **63**, 871-880.
- Wilby, R.L., Beven, K.J. & Reynard, N.S. (2008) Climate change and fluvial flood risk in the UK: more of the same? *Hydrological Processes* **22**, 2511–2523.
- Wolter, C. & Vilcinskas, A. (1997) Characterization of the typical fish community of inland waterways of the North-Eastern lowlands in Germany. *Regulated Rivers: Research & Management* **13**, 335-343.
- Wolter, C. (1999) Comparison of intraspecific genetic variability in four common cyprinids, *Abramis brama*, *Abramis bjoerkna*, *Rutilus rutilus* and *Scardinius erthrophthalmus*, within and between lowland river systems. *Hydrobiologia* **364**, 163-177.
- Wolter, C. (2001a) Conservation of fish species diversity in navigable waterways. *Landscape and Urban Planning* **53**, 135-144.



- Wolter, C. (2001b) Rapid changes of fish assemblages in artificial lowland waterways. *Limnologica* **31**, 27-35.
- Wolter, C. & Freyhof, J. (2004) Diel distribution patterns of fishes in a temperate large lowland river. *Journal of Fish Biology* **64**, 632-642.
- Woolnough, D.A., Downing, J.A. & Newton, T.J. (2009) Fish movement and habitat use depends on water body size and shape. *Ecology of Freshwater Fish* **18**, 83-91.
- Worthington, T., Kemp, P.S., Osborne, P.E., Howes, C. and Easton, K. (2011) A review of the historical distribution and status of the burbot (*Lota lota*) in English rivers. *Journal of Applied Ichthyology* **27**, 1-8.
- Volta, P., Jeppesen, E., Leoni, B., Campi, B., Sala, P., Garibaldi, L., Lauridsen, T. L. & Winfield, I. J. (2013) Recent invasion by a non-native cyprinid (common bream *Abramis brama*) is followed by major changes in the ecological quality of a shallow lake in southern Europe. *Biological Invasions*, DOI: 10.1007/s10530-013-0433-z.
- Vostradovsky, J. (1975). Horizontal distribution of individually tagged fish in the Lipno reservoir (English). *EIFAC Technical Papers* **23** (Suppl. 1, V.2), 651–655.
- Young, M. K. (1999) Summer diel activity and movement of adult brown trout in high elevation streams in Wyoming, U.S.A. *Journal of Fish Biology* **54**, 181–189.